Warming and Forecast Skill in Monthly Temperature: Frankfurt am Main, 1981–2025

Björn Hanneke with e2er[‡]

October 20, 2026

1 Introduction

Month-to-month air temperature in Central Europe is dominated by a strong, nearly deterministic annual cycle, on top of which irregular anomalies of roughly two degrees Celsius obscure any slow change in the mean. For planners in municipal heat management, energy supply, and agriculture, this creates a concrete forecasting question: at horizons of up to three years, does a statistical model add anything to the heuristic of reusing the value from the same month one year earlier? We study this question for monthly mean air temperature at 2 metres in Frankfurt am Main, using NASA POWER data (NASA Langley Research Center (LaRC) POWER Project) for the grid cell containing 50.11°N, 8.68°E from January 1981 to December 2025.

This paper addresses two descriptive and predictive questions. First, how has the monthly mean temperature in Frankfurt changed since 1981, in its overall level, across seasons, and in the amplitude and phase of the seasonal cycle? Second, how well do seasonal forecasting models predict the 36 months of a fixed hold-out relative to a seasonal naive baseline? We make no causal claim. All trend estimates are associations with calendar time, and the forecasting comparison evaluates predictive accuracy only.

Our design fixes the forecast setup before any model is fitted. Models are estimated on January 1981 to December 2022 (504 months) and evaluated on the final 36 months, January 2023 to December 2025. We compare two baselines (seasonal naive and naive) with four candidates: additive Holt-Winters exponential smoothing without trend, ETS(A,N,A), and with damped trend, ETS(A,Ad,A); a seasonal ARIMA model with orders chosen on the training window; and a regression on a linear trend and annual harmonics. Accuracy is measured by RMSE, MAE, and MASE,

[*] This paper was produced with e2er. Its dossier lists every step of the run (the specialists, the checks and the researcher's actions), the files the steps produced, and the template, specialists, skills, connectors and AI models used, pinned to the versions the run used where E2ER's repository holds them: `https://e2er.org/d/c985731bebd41662`.

[†] Demonstration. This study was produced with e2er as a demonstration or test run and is published as is. It is not presented as a research contribution, and its author does not vouch for its findings.

and differences against the seasonal naive baseline are tested with the Diebold-Mariano test. We report every declared model, including those that lose to the baseline.

Our findings can be summarised as follows. Annual mean temperature in Frankfurt rose by about 0.50 degrees Celsius per decade over 1981–2022, and the warming appears in all four seasons, with seasonal estimates between 0.38 and 0.62 degrees per decade that differ by less than sampling error. We find no evidence that the amplitude or phase of the seasonal cycle changed. On the hold-out, all four candidate models reduce the root mean squared error of the seasonal naive baseline by roughly 30 percent, and the differences are statistically significant at the 5 percent level, although the short hold-out limits the power of this comparison. The naive last-value forecast is far worse than the seasonal naive baseline. We find no evidence that the four candidates differ from one another.

We contribute in three ways. First, we provide a transparent and reproducible local trend estimate with explicit uncertainty for a reanalysis grid cell in the Rhine-Main region. Second, we quantify how much simple statistical models that average over several years improve on the single-year seasonal naive forecast, with a setup frozen before estimation; because the design contains no climatological-mean benchmark, it does not separate the gain from trend modelling from the gain from averaging out one year of anomaly noise. Third, we report that model-based prediction intervals are about half as wide as those of the baseline while containing 35 of 36 hold-out values; the sample is too small to test calibration.

Related work falls into three groups. Observed warming in Central Europe is documented in national weather service reports and regional climate assessments, and reanalysis products such as MERRA-2 are widely used for trend analysis, although changes in the assimilated observing system can introduce inhomogeneities into reanalysis trends. In forecasting, competition evidence shows that simple benchmarks are hard to beat, which motivates the seasonal naive baseline, whereas dynamical seasonal and decadal prediction systems form a different class that we do not consider. We do not benchmark our trend against published German Weather Service (DWD) figures for Hesse or Germany, and we do not validate the POWER series against stations; the 0.50 degrees per decade estimate therefore has no external benchmark in this study, and the comparison is left to future work.

We describe the data, the empirical strategy, and the results in turn, and then discuss implications and limitations.

2 Data

2.1 NASA POWER and the Frankfurt grid cell

We use the monthly 2 metre air temperature (T2M) from the NASA Prediction of Worldwide Energy Resources (POWER) project (NASA Langley Research Center (LaRC) POWER Project) (API version 2.10.0, accessed 13 October 2026). The MERRA-2 cell measures 0.5° in latitude by 0.625° in longitude, and the grid elevation is 171.6 m. POWER derives its meteorological

parameters from the MERRA-2 reanalysis, which assimilates observations into a global atmospheric model on a regular grid. The series is therefore a grid-cell reanalysis value for the cell containing 50.11°N, 8.68°E, not a station record. Local effects such as the urban heat island of Frankfurt, and any differences between the grid-cell elevation and the elevation of a measuring site, are only partly represented. Absolute levels and trends should consequently be read as properties of the reanalysis cell, and not as those of an individual weather station.

2.2 Sample and variables

The sample contains 540 monthly observations from January 1981 to December 2025. The data end in December 2025 because the POWER monthly load covers whole calendar years; months of 2026 are excluded by design, so the forecasts for 2026 to 2028 remain unevaluated. No observation is missing, no fill values occur, and all 45 years are complete. We split the sample at a fixed origin: the 504 months from 1981-01 to 2022-12 form the training window, and the 36 months from 2023-01 to 2025-12 form the hold-out. The mean monthly temperature is 9.94 degrees Celsius with a standard deviation of 7.54, reflecting the annual cycle. The training mean is 9.83 degrees, whereas the hold-out mean is 11.46 degrees, which already indicates that the hold-out period is warmer than the average of the training window. Anomalies relative to the 1981–2010 monthly climatology have a standard deviation of 1.90 degrees over the full sample and a mean of 0.44 degrees. Decade means of the annual temperature rise from 9.09 degrees in 1981–1990 to 11.19 degrees in 2021–2025. Figure 1 shows the monthly series and Figure 2 the annual means.

3 Empirical Strategy

3.1 Trend description

We describe the long-run change in three ways, all on the training years 1981–2022 and, for comparison, on the full sample. First, we regress annual mean temperature $\bar{y}_s$ on calendar year s by OLS,

$$\bar{y}_s = \alpha + \beta\,(s - s_0) + u_s, \tag{1}$$

with Newey-West (HAC) standard errors based on a Bartlett kernel and bandwidth

$$L = \lfloor 4\,(T/100)^{2/9} \rfloor, \tag{2}$$

where T is the number of observations in the regression. The slope is reported as $10\hat{\beta}$ in degrees Celsius per decade. Second, we repeat this on the deseasonalised monthly series $\tilde{y}_t = y_t - \hat{\mu}_{m(t)}$, where $\hat{\mu}_m$ is the mean of calendar month m in the sample used, regressing $\tilde{y}_t$ on elapsed time in years. Third, we estimate season-specific slopes for December–February, March–May, June–August, and September–November. Because the annual regression is estimated on only 42 observations, we complement it with the nonparametric Mann-Kendall test (9) and Sen's slope (11), including an




Figure 1: Monthly mean 2 m air temperature, Frankfurt am Main grid cell (NASA POWER), 1981–2025.

autocorrelation-corrected variant. We also test whether the amplitude and phase of the annual harmonic drift linearly over time. With elapsed time τ_t in years, we estimate

$$y_t = a + b\,\tau_t + (c_1 + d_1\tau_t)\cos(2\pi\tau_t) + (c_2 + d_2\tau_t)\sin(2\pi\tau_t) + u_t, \tag{3}$$

so that the amplitude at time τ is $A(\tau) = \sqrt{(c_1 + d_1\tau)^2 + (c_2 + d_2\tau)^2}$ and the phase is $\phi(\tau) = \operatorname{atan2}(c_2 + d_2\tau,\, c_1 + d_1\tau)$. The joint Wald test of $H_0\colon d_1 = d_2 = 0$ uses HAC standard errors with the lag from Equation (2). These estimates describe associations with calendar time and do not identify causes of warming.

3.2 Stationarity diagnostics

We apply the Augmented Dickey-Fuller test (2) and the KPSS test (8) to the seasonally adjusted training series. The ADF test has a unit root as null hypothesis, whereas KPSS has stationarity as null. All diagnostics are computed on data up to December 2022 and therefore do not use the hold-out.




Figure 2: Annual mean 2 m air temperature, Frankfurt am Main grid cell (NASA POWER), 1981–2025.

3.3 Forecasting models and evaluation

The forecast setup was fixed before estimation and frozen together with the hold-out. The two baselines are the seasonal naive forecast (the value twelve months earlier) and the naive forecast (the last observed value) (5). The four candidates are ETS(A,N,A) and ETS(A,Ad,A), i.e., additive Holt-Winters exponential smoothing without and with a damped trend; a seasonal ARIMA model (1) whose orders are chosen on the training window by AICc in a stepwise search in the spirit of Hyndman and Khandakar (6); and an OLS regression on a linear trend and annual harmonics, with the number of harmonics chosen by AICc. Model choice uses the training window only. Every model is fitted at the origin December 2022 and forecasts the 36 months of the hold-out in a single pass, so that all hold-out errors are genuine out-of-sample errors (12). Prediction intervals are at the 95 percent level, and we report the share of hold-out values that fall inside them.

We measure accuracy by RMSE and MAE in degrees Celsius and by the mean absolute scaled error (MASE), which scales the MAE by the in-sample MAE of the seasonal naive method on the training window (7). This scale is 2.04 degrees. To test whether a candidate forecasts better than the seasonal naive baseline, we use the Diebold-Mariano test (3) with the small-sample correction of Harvey et al. (4), applied to squared and absolute error differences over the 36 hold-out months. A rolling-origin exercise inside the training window (twenty annual origins from December 2000

Table 1: Seasonal trends, training years 1981–2022, degrees Celsius per decade. HAC (Newey–West) standard errors; associations with calendar time, not causal effects

Season	Slope	HAC s.e.	CI lower	CI upper	N
December–February	0.621	0.195	0.238	1.003	41
March–May	0.377	0.123	0.136	0.619	42
June–August	0.607	0.133	0.347	0.867	42
September–November	0.389	0.105	0.183	0.595	42

to December 2019, each with a 36-month horizon) supplements the single hold-out; because these origins overlap, we use it to assess horizon and season structure and not to select models. Finally, each model is refitted on all 540 months to produce forecasts for January 2026 to December 2028, which are not evaluated.

We stated three expectations before seeing the results. The first (H1) was a positive annual trend of between 0.3 and 0.5 degrees per decade. The second (H2) was that the candidate models improve on the seasonal naive baseline by a modest margin, with a hold-out MASE of 0.75 to 0.95. The third (H3) was that the exponential smoothing and ARIMA models do not clearly beat the harmonic trend regression on a three-year hold-out. We evaluate them in Section 4 without revision.

4 Results

4.1 Trend, seasonality, and stationarity

Annual mean temperature in Frankfurt increased by 0.50 degrees Celsius per decade over 1981–2022 (HAC standard error 0.08, 95 percent confidence interval 0.34 to 0.67). The estimate sits at the upper edge of the range of 0.3 to 0.5 degrees that we stated in advance for H1; the confidence interval overlaps that range but extends to 0.67. The slope from the deseasonalised monthly series is practically identical at 0.50 degrees per decade, and extending the sample to 2025 raises the annual estimate to 0.54 degrees per decade. The Mann-Kendall test rejects the null of no monotone trend in annual means (statistic 4.48, $p < 0.001$) with a Sen slope of 0.50 degrees per decade; the variance adjustment for autocorrelation leaves the statistic unchanged, since the estimated variance inflation factor equals one (Table 2).

Warming appears in every season (Table 1). Over the training years, the slope is 0.62 degrees per decade in winter and 0.61 in summer, compared with 0.38 in spring and 0.39 in autumn. The confidence intervals of the four seasons overlap substantially, so the data do not support a claim that the seasons warm at different rates. We find no evidence that the seasonal cycle changed: the fitted amplitude of the annual harmonic changes by -0.15 degrees and the phase by -1.6 degrees between the start and end of the training window, and the joint Wald test of both drift terms does not reject constancy ($p = 0.76$).

The stationarity tests, computed on the seasonally adjusted training series, point in different

Table 2: Stationarity and trend diagnostics on the training period (sample end 2022-12). KPSS p-value is bounded at 0.01 by the table

Test	Statistic	p-value
ADF, seasonally adjusted	-6.868	0.0000
KPSS (level), seasonally adjusted	1.987	0.0100
Mann–Kendall, annual means	4.476	0.0000

Table 3: Out-of-sample accuracy on the 36-month hold-out, 2023-01 to 2025-12. RMSE and MAE in degrees Celsius; MASE scaled by the in-sample seasonal naive MAE; nominal interval level 95%

Model	RMSE	MAE	MASE	Coverage	Mean width
Seasonal naive (baseline)	1.917	1.640	0.803	1.000	14.21
Naive (baseline)	12.058	9.784	4.790	1.000	71.44
ETS(A,N,A)	1.301	0.976	0.478	0.972	7.34
ETS(A,Ad,A)	1.300	0.976	0.478	0.972	7.34
SARIMA	1.369	1.002	0.490	0.972	7.18
Trend + harmonics	1.362	1.029	0.504	0.972	7.19

directions. The ADF test rejects a unit root (statistic -6.87), whereas the KPSS test rejects level stationarity (statistic 1.99, p-value bounded at 0.01). The ADF rejection argues against a unit root, and a level-KPSS rejection is expected for a series with a deterministic upward drift. We did not compute trend-stationary variants of either test, so we read the pair as consistent with a drift and no unit root, not as a demonstration of trend stationarity.

4.2 Hold-out forecast accuracy

Table 3 reports the out-of-sample accuracy of all six models on the 36 hold-out months. The seasonal naive baseline attains an RMSE of 1.92 degrees Celsius, an MAE of 1.64 degrees, and a MASE of 0.80. The naive last-value forecast performs far worse, with an RMSE of 12.06 degrees and a MASE of 4.79, because it ignores the annual cycle entirely; it loses to the seasonal naive baseline by a wide margin, and its Diebold-Mariano statistics are strongly negative. All four candidates beat the seasonal naive baseline. ETS(A,Ad,A) has the lowest RMSE (1.30 degrees), followed closely by ETS(A,N,A) (1.30), the harmonic trend regression (1.36), and SARIMA (1.37). Their MASE values lie between 0.48 and 0.50, i.e., hold-out errors of about half the in-sample seasonal naive MAE. The RMSE ratio relative to seasonal naive is 0.68 for both exponential smoothing variants and 0.71 for SARIMA and the harmonic regression. The two exponential smoothing variants produce identical accuracy figures at the reported precision, so we do not count them as independent evidence. The direction of H2 holds on the hold-out, but its magnitude does not: the realised MASE of 0.48 to 0.50 lies well below the pre-stated range of 0.75 to 0.95.

Table 4 shows that the improvements are statistically significant but not overwhelming. With the Harvey-Leybourne-Newbold correction and squared-error loss, the Diebold-Mariano statistic

Table 4: Diebold–Mariano tests against the seasonal naive baseline on the hold-out (Harvey–Leybourne–Newbold correction, $t(35)$, truncation 12). Positive statistics mean the model has lower loss than seasonal naive

Model	Squared, stat.	Squared, p	Absolute, stat.	Absolute, p
Naive	-8.298	0.0000	-9.866	0.0000
ETS(A,N,A)	2.746	0.0095	2.589	0.0139
ETS(A,Ad,A)	2.753	0.0093	2.589	0.0139
SARIMA	2.167	0.0372	2.243	0.0314
Trend + harmonics	2.466	0.0187	2.519	0.0165

against seasonal naive is 2.75 for ETS(A,N,A) ($p = 0.009$), 2.75 for ETS(A,Ad,A) ($p = 0.009$), 2.47 for the harmonic regression ($p = 0.019$), and 2.17 for SARIMA ($p = 0.037$). Absolute-error loss gives the same ranking and conclusions. With only 36 monthly errors that are serially dependent, the test has limited power and the p-values should be read as indicative. We apply no multiple-testing adjustment across the four comparisons against one baseline, and the candidate models are nested in related model classes. Pairwise tests among the candidates reveal no significant difference: the smallest p-value is 0.18 (ETS(A,Ad,A) against the harmonic regression), and none of the three pairs is significant at the 5 percent level, which is consistent with H3; with 36 months, this reflects low power and not evidence of equivalence.

4.3 Prediction intervals

The 95 percent intervals of the seasonal naive baseline cover all 36 hold-out values (coverage 100 percent), but they are wide, with a mean width of 14.2 degrees. The candidate models cover 35 of the 36 values (97.2 percent) with intervals that are about half as wide (mean width 7.2 to 7.3 degrees). The intervals of the candidates contain 35 of 36 values and those of the baseline all 36. With 36 serially dependent observations, neither rate can be distinguished from 95 percent, so we draw no conclusion on calibration; the difference in width is the robust finding.

4.4 Horizon and season structure

The error of the seasonal naive baseline is flat across horizons: RMSE is 1.92 degrees in months 1 to 12, 1.94 in months 13 to 24, and 1.89 in months 25 to 36. This is expected because its forecast does not depend on the horizon beyond the interval width. The relative advantage of the candidates is largest in the final year of the hold-out, where their RMSE is about 43 to 46 percent of the baseline's. This pattern coincides with the warm years 2023 to 2025 and is consistent with the level shift that the baseline, anchored on a single past year, cannot track; with only three hold-out years, however, we cannot separate this from chance.

The rolling-origin exercise in the training window gives a less favourable picture for the seasonal naive baseline and a more modest one for the candidates. Across twenty overlapping origins, the baseline's MASE is 1.02, whereas the candidate models reach MASE values between 0.71 and 0.73.

In the rolling exercise, the RMSE ratio at horizons 25 to 36 is between 0.71 and 0.73. Hence the advantage of the candidates over seasonal naive is reproduced outside the final hold-out, and it is smaller than on the 2023–2025 window.

4.5 Forecasts beyond the data

Refitted on all 540 months, the harmonic regression estimates a linear trend of 0.054 degrees per year. We produce forecasts for January 2026 to December 2028 for all six models with 95 percent intervals but do not tabulate them here. Because these forecasts lie beyond the data, they cannot be evaluated here, and we do not interpret them as predictions of any particular year.

5 Discussion and Conclusion

Monthly mean air temperature in the Frankfurt reanalysis grid cell rose by about 0.5 degrees Celsius per decade between 1981 and 2022, in all four seasons and no detectable change in the amplitude or phase of the annual cycle. On a hold-out of 36 months, simple statistical models that combine the seasonal pattern with a slowly moving level or a linear trend reduce the RMSE of the seasonal naive baseline by around 30 percent, and these gains are significant at the 5 percent level but not decisive given the short hold-out. We find no evidence that the choice among exponential smoothing, SARIMA, and a harmonic regression matters for accuracy at this horizon. The naive last-value forecast ignores the annual cycle and performs far worse.

For practitioners, the results suggest that reusing last year's value is a defensible heuristic with a larger error: the seasonal naive RMSE is 1.92 degrees against 1.30 to 1.37 for the candidates, whose intervals are about half as wide. Because the three-year hold-out coincides with a warm period, the gain may exceed that in an average window, although the hold-out RMSE ratio of 0.68 to 0.71 is close to the rolling-origin ratio of 0.71 to 0.73 at horizons 25 to 36, so we cannot establish a difference. For scholars, the study shows that models averaging over several years beat the single-year heuristic; because the design has no climatological-mean benchmark, it does not show that the trend of roughly 0.5 degrees per decade is what drives the gain.

Our analysis is subject to several scope conditions. The data are a MERRA-2 reanalysis grid cell, not a station record, so trends and levels may differ from those measured at a weather station; a comparison with station observations from the German Weather Service (DWD) is a natural extension but is not part of this study, and no DWD data are used here. The hold-out contains only 36 serially dependent months, which limits the power of the Diebold-Mariano comparisons and the precision of the coverage estimates. Reanalysis trends may also be affected by changes in the assimilated observing system, which we did not examine, and the ranking rests on a single forecast origin and a single grid cell. Finally, the trend estimates are associations with calendar time and the analysis makes no claim about the causes of warming.

We leave to future work whether the gain over seasonal naive persists when the models are evaluated against DWD station observations, and whether it exceeds the gain of a climatological-

mean benchmark.

References

[1] Box, G. E. P., Jenkins, G. M., and Reinsel, G. C. (2008). *Time Series Analysis: Forecasting and Control (4th edition)*. Wiley.

[2] Dickey, D. A. and Fuller, W. A. (1979). Distribution of the Estimators for Autoregressive Time Series with a Unit Root. *Journal of the American Statistical Association*, 74(366a):427–431.

[3] Diebold, F. X. and Mariano, R. S. (1995). Comparing Predictive Accuracy. *Journal of Business & Economic Statistics*, 13(3):253–263.

[4] Harvey, D., Leybourne, S., and Newbold, P. (1997). Testing the equality of prediction mean squared errors. *International Journal of Forecasting*, 13(2):281–291.

[5] Hyndman, R. J. and Athanasopoulos, G. (2021). *Forecasting: Principles and Practice (3rd edition)*. OTexts, Melbourne.

[6] Hyndman, R. J. and Khandakar, Y. (2008). Automatic Time Series Forecasting: The forecast Package for R. *Journal of Statistical Software 27(3)*.

[7] Hyndman, R. J. and Koehler, A. B. (2006). Another look at measures of forecast accuracy. *International Journal of Forecasting*, 22(4):679–688.

[8] Kwiatkowski, D., Phillips, P. C., Schmidt, P., and Shin, Y. (1992). Testing the null hypothesis of stationarity against the alternative of a unit root. *Journal of Econometrics*, 54(1-3):159–178.

[9] Mann, H. B. (1945). Nonparametric Tests Against Trend. *Econometrica*, 13(3):245–259.

[NASA Langley Research Center (LaRC) POWER Project] NASA Langley Research Center (LaRC) POWER Project. Prediction of worldwide energy resources (POWER). https://power.larc.nasa.gov/. Funded through the NASA Earth Science Division. Cite the API version and access date of each load (data_sources.json).

[11] Sen, P. K. (1968). Estimates of the Regression Coefficient Based on Kendall's Tau. *Journal of the American Statistical Association*, 63(324):1379–1389.

[12] Tashman, L. J. (2000). Out-of-sample tests of forecasting accuracy: an analysis and review. *International Journal of Forecasting*, 16(4):437–450.