DEPARTMENT OF CLINICAL BIOCHEMISTRY

Patient [redacted]	[redacted]	[redacted]	[redacted]
Admn [redacted] **No**	: [redacted] / UMR9996	[redacted] **By**	: Dr.CHENNA VIJAY MD, DM (NEUROLOG
Bill Date	: 22-Apr-25 08:40 am	**Report Date**	: 22-Apr-25 09:47 am

Parameter	Specimen	Results	Biological Reference	Method
SGPT/ALT		13	0 - 50 U/L	UV WITHOUT P5P
SGOT/AST		26	0 - 50 U/L	UV WITHOUT P5P
Alkaline Phosphatase		**169**	30 - 120 U/L	PNPP AMP BUFFER
TOTAL PROTEIN		**5.8**	6.3 - 8.2 gm/dl	BIURET
ALBUMIN		3.5	3.5 - 5.2 g/dl	BROMOCRESOL GREEN
Globulin		2.3	2.0 - 3.5 gm/dl	Calculation
A/G Ratio		1.52		
CRP		2.6	0 - 6.0 mg/L	LATEX PARTICLE IMMUNOTURBIDIMETRIC
Calcium		**7.5**	8.8 - 10.6 mg/dl	ARSENAZO
Phosphorous		2.9	2.5 - 4.5 mg/dl	PHOSPOMOLYBDATE COMPLEX
Magnesium		2.06	1.8 - 2.6 mg/dl	XYLIDYL BLUE
PRO CALCITONIN				
Procalcitonin		0.861	Low Risk for Sepsis: <0.5 - High risk for sepsis:>2.0 ng/mL	Enhanced Chemiluminescence

Interpretation

1. PCT is the prohormone of calcitonin secreted by different types of cells from numerous organs in response to proinflammatory stimulation, particularly bacterial stimulation.
2. Depending upon clinical background, PCT > 0.1 can indicate clinically relevant bacterial infection, requiring antibiotic treatment.
3. PCT is an important marker enabling specific differentiation between bacterial infection and other causes of inflammatory reactions
4. Resorption of septic infection is accompanied by decreases in PCT concentration which returns to normal with half life of 24 hours
5. In certain situations (newborns, polytrauma, burns,major surgery, prolonged or severe cardiogenic shock etc) PCT elevation may be independent of any infectious aggression. The return to normal values is usually rapid
6. Viral infections, allergies, autoimmune diseases and graft rejection do not lead to significant increase in PCT. Localized bacterial infections can lead to moderate increase in PCT levels.

NOTE.

Patients on Biotin supplement may have interference in some immunoassays. With individuals taking high dose Biotin (more than 5 mg per day) supplements, at least 8-hour wait time before blood draw is recommended.

Low Risk for Sepsis : < 0.5 ng/ml.
High Risk for Sepsis : > 2.0 ng/ml.