

Minimum Risk Bayes Decision Theoretic Classifier

Bayes Theorem: $p(y|x) = p(x|y) \cdot p(y) \rightarrow \text{posterior} \propto \text{likelihood} \times \text{prior}$

$p(x|y)$ is multivariate Gaussian

$$p(x|y) = \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp\left(-\frac{1}{2} (x - \mu)^T \Sigma^{-1} (x - \mu)\right)$$

where μ and Σ are μ_{MLE} and Σ_{MLE} , respectively.

Training

Compute the values of μ_{MLE} and Σ_{MLE} .

$$\mu_{MLE_i}^T = \frac{1}{m} \sum_{j=1}^m x_{train_i}^{(j)}$$

- μ_{MLE_i} is the mean vector of the i^{th} class training set.
- m is the number of samples in the i^{th} class training set.
- $x_{train_i}^{(j)}$ is the j^{th} sample of the i^{th} class training set.

$$\Sigma_{MLE_i} = \frac{1}{m} (X_{train_i} - M_{train_i})^T (X_{train_i} - M_{train_i})$$

- X_{train_i} is the feature matrix of the i^{th} class training set.
- M_{train_i} is the mean matrix of the i^{th} class training set

Classification

Compute the discriminant function for each class to find the final class label.

$$g_i(x_{test}) = -\frac{1}{2} (x_{test} - \mu_{MLE_i})^T \Sigma_{MLE_i}^{-1} (x_{test} - \mu_{MLE_i}) - \frac{d}{2} \log(2\pi) - \frac{1}{2} \log|\Sigma_{MLE_i}| + \log(p(w_i))$$

The term $\frac{d}{2} \log(2\pi)$ can be dropped.

- $p(w_i)$ is the i^{th} class prior probability
- Class label = $\text{argmax}(g_1, g_2, \dots)$

EXAMPLE

2D dataset with 2 classes and equal prior probabilities.

petal_length	petal_width	class
1.5	0.2	1
3.0	1.1	2
1.3	0.2	1
4.9	1.5	2
1.4	0.3	1
5.0	1.7	2
3.7	1.0	2
1.9	0.2	1

Training

Class 1 training data: $\begin{bmatrix} 1.5 & 0.2 \\ 1.3 & 0.2 \\ 1.4 & 0.3 \\ 1.9 & 0.2 \end{bmatrix}$

Class 1 mean vector: $\mu_{MLE1}^T = [1.525 \quad 0.225]$

Class 1 mean matrix: $\begin{bmatrix} 1.525 & 0.225 \\ 1.525 & 0.225 \\ 1.525 & 0.225 \\ 1.525 & 0.225 \end{bmatrix}$

Class 1 covariance:

$$\Sigma_{MLE1} = \frac{1}{4} \begin{bmatrix} 1.5 - 1.525 & 0.2 - 0.225 \\ 1.3 - 1.525 & 0.2 - 0.225 \\ 1.4 - 1.525 & 0.3 - 0.225 \\ 1.9 - 1.525 & 0.2 - 0.225 \end{bmatrix}^T \begin{bmatrix} 1.5 - 1.525 & 0.2 - 0.225 \\ 1.3 - 1.525 & 0.2 - 0.225 \\ 1.4 - 1.525 & 0.3 - 0.225 \\ 1.9 - 1.525 & 0.2 - 0.225 \end{bmatrix} = \begin{bmatrix} 0.0692 & -0.0042 \\ -0.0042 & 0.0025 \end{bmatrix}$$

Class 2 training data: $\begin{bmatrix} 3.0 & 1.1 \\ 4.9 & 1.5 \\ 5.0 & 1.7 \\ 3.7 & 1.0 \end{bmatrix}$

Class 2 mean vector: $\mu_{MLE2}^T = [4.15 \quad 1.325]$

Class 2 covariance: $\Sigma_{MLE2} = \begin{bmatrix} 0.9367 & 0.2850 \\ 0.2850 & 0.1091 \end{bmatrix}$

Classification

Class 1 test data: $x_{test}^T = [1.5 \quad 0.2]$

$$g_1 = -\frac{1}{2} [1.5 - 1.525]^T \begin{bmatrix} 0.0692 & -0.0042 \\ -0.0042 & 0.0025 \end{bmatrix}^{-1} [1.5 - 1.525] - \frac{1}{2} \log \left(\begin{vmatrix} 0.0692 & -0.0042 \\ -0.0042 & 0.0025 \end{vmatrix} \right) + \log \left(\frac{1}{2} \right) = 3.5304$$

$$g_2 = -4.7728$$

Predicted class label: $\hat{y} = \operatorname{argmax} (g_1, g_2) = 1$

Note:

$$A \cdot A^{-1} = I$$

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$