
Ultracold molecule assembly

Yichao Yu


Ni Group/Harvard

Aug 11, 2017


- Strong and tunable interaction
- Rich internal energy levels
- High filling fraction
- Single site detection and manipulation




- Strong and tunable interaction
- Rich internal energy levels
- High filling fraction
- Single site detection and manipulation




- Strong and tunable interaction
- Rich internal energy levels
- High filling fraction
- Single site detection and manipulation



- Strong and tunable interaction
- Rich internal energy levels
- High filling fraction
- Single site detection and manipulation




- Strong and tunable interaction
- Rich internal energy levels
- High filling fraction
- Single site detection and manipulation




Applications

Simulation of many-body system^[1]



[1] B. Yan et al., "Observation of dipolar spin-exchange interactions with lattice-confined polar molecules.", Nature **501**, 521–5 (2013).

Applications




[2] S. F. Yelin et al., "Schemes for robust quantum computation with polar molecules", 4 (2006).

- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!



- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!



- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!





- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!


- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!


- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!

Atom loading and cooling

Single atoms


 85% cround state after Cesium Raman sideband cooling







Atom loading and cooling

- Single atoms
- 85% cround state after Cesium Raman sideband cooling







- High initial temperature $(70\mu K)$
- High Lamb Dicke parameter
- Large light shift (scalar and tensor)
- Trap anharmonicity




- High initial temperature $(70\mu K)$
- High Lamb Dicke parameter
- Large light shift (scalar and tensor)
- Trap anharmonicity




- High initial temperature $(70\mu K)$
- High Lamb Dicke parameter
- Large light shift (scalar and tensor)
- Trap anharmonicity







- High initial temperature $(70\mu K)$
- High Lamb Dicke parameter
- Large light shift (scalar and tensor)
- Trap anharmonicity




- High initial temperature $(70\mu K)$
- High Lamb Dicke parameter
- Large light shift (scalar and tensor)
- Trap anharmonicity



- High initial temperature $(70\mu K)$
- High Lamb Dicke parameter
- Large light shift (scalar and tensor)
- Trap anharmonicity




- High initial temperature $(70\mu K)$
- High Lamb Dicke parameter
- Large light shift (scalar and tensor)
- Trap anharmonicity




- High initial temperature $(70\mu K)$
- High Lamb Dicke parameter
- Large light shift (scalar and tensor)
- Trap anharmonicity


Sequence and simulation





Sequence and simulation




Sequence and simulation


















| AAIS I (GAIGI) | |
|----------------|--------------------------|
| Axis | Ground state probability |
| 1 (Axial) | 93.1(2.5)% |
| 2 (Radial) | 91.9(2.3)% |
| 3 (Radial) | 92.9(2.5)% |


3D ground state: 79.5(3.6)%Loss after cooling: 15%

Total 3D ground state preparation fidelity: 67.6(3.1)%

Rabi flopping (radial)




Rabi flopping (radial)



Good agreement in ground state probability between spectrum and Rabi flopping data.

Rabi flopping (axial)



In progress

