
# Ultracold molecule assembly

Yichao Yu


Ni Group/Harvard

Aug 11, 2017


- Strong and tunable interaction
- Rich internal energy levels
- High filling fraction
- Single site detection and manipulation




- Strong and tunable interaction
- Rich internal energy levels
- High filling fraction
- Single site detection and manipulation




- Strong and tunable interaction
- Rich internal energy levels
- High filling fraction
- Single site detection and manipulation



- Strong and tunable interaction
- Rich internal energy levels
- High filling fraction
- Single site detection and manipulation




- Strong and tunable interaction
- Rich internal energy levels
- High filling fraction
- Single site detection and manipulation



# **Applications**


# Simulation of many-body system<sup>[1]</sup>



$$H \propto \sum V_{ij} \left( S_i^+ S_j^- + S_i^- S_j^+ \right)$$

[1] B. Yan et al., "Observation of dipolar spin-exchange interactions with lattice-confined polar molecules.", Nature **501**, 521–5 (2013).

# **Applications**




[2] S. F. Yelin et al., "Schemes for robust quantum computation with polar molecules", Phys. Rev. A 74, 050301 (2006).

- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!



- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!



- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!

- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!


- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!


- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!

# Atom loading and cooling

# Single atoms


 85% ground state after Cesium Raman sideband cooling








### Atom loading and cooling

- Single atoms
- 85% ground state after Cesium Raman sideband cooling












- High initial temperature  $(70\mu K)$
- High Lamb Dicke parameter
- Large light shift
- Trap anharmonicity
- Off resonance scattering from Raman beams  $\approx 0.2 \sim 0.5 \text{kHz}$




- High initial temperature  $(70\mu K)$
- High Lamb Dicke parameter
- Large light shift
- Trap anharmonicity
- Off resonance scattering from Raman beams  $\approx 0.2 \sim 0.5 \text{kHz}$




- High initial temperature  $(70\mu K)$
- High Lamb Dicke parameter
- Large light shift
- Trap anharmonicity
- Off resonance scattering from Raman beams  $\approx 0.2 \sim 0.5 \text{kHz}$







- High initial temperature  $(70\mu K)$
- High Lamb Dicke parameter
- Large light shift
- Trap anharmonicity
- Off resonance scattering from Raman beams  $\approx 0.2 \sim 0.5 \text{kHz}$




- High initial temperature  $(70\mu K)$
- High Lamb Dicke parameter
- Large light shift
- Trap anharmonicity
- Off resonance scattering from Raman beams
  ≈ 0.2 ~ 0.5kHz



- High initial temperature  $(70\mu K)$
- High Lamb Dicke parameter
- Large light shift
- Trap anharmonicity
- Off resonance scattering from Raman beams  $\approx 0.2 \sim 0.5 \text{kHz}$




- High initial temperature  $(70\mu K)$
- High Lamb Dicke parameter
- Large light shift
- Trap anharmonicity
- Off resonance scattering from Raman beams  $\approx 0.2 \sim 0.5 \text{kHz}$




- High initial temperature  $(70\mu K)$
- High Lamb Dicke parameter
- Large light shift
- Trap anharmonicity
- Off resonance scattering from Raman beams  $\approx 0.2 \sim 0.5 \text{kHz}$


# Sequence and simulation





### Sequence and simulation




# Sequence and simulation


















AAIS I (GAIGI)	
Axis	Ground state probability
1 (Axial)	93.1(2.5)%
2 (Radial)	91.9(2.3)%
3 (Radial)	92.9(2.5)%


3D ground state: 79.5(3.6)%Loss after cooling: 15%

**Total 3D ground state preparation fidelity:** 67.6(3.1)%

### Rabi flopping (radial)




### Rabi flopping (radial)




Good agreement in ground state probability between spectrum and Rabi flopping data.

# Rabi flopping (axial)




# In progress









——— Strong dipole

Weak dipole

# **Quantum computation** Strong dipole Weak dipole

