
Trapping and imaging of single atoms in the presence of light shift




Yichao Yu May 26, 2016 Ni Group/Harvard

Group members
Nicholas Hutzler
Lee Liu
Jessie Zhang
PI
Kang-Kuen Ni


- MOT Loading
- Trapping
- Imaging
- Works for Cs
- Doesn't work for Na




- MOT Loading
- Trapping
- Imaging
- Works for Cs
- Doesn't work for Na



- MOT Loading
- Trapping
- Imaging
- Works for Cs
- Doesn't work for Na




- MOT Loading
- Trapping
- Imaging
- Works for Cs
- Doesn't work for Na




- MOT Loading
- Trapping
- Imaging
- Works for Cs
- Doesn't work for Na

- Inefficient cooling; Heating
- Shift imaging light out of resonance




- Inefficient cooling; Heating
- Shift imaging light out of resonance









- Inefficient cooling; Heating
- Shift imaging light out of resonance









- Inefficient cooling; Heating
- Shift imaging light out of resonance








- Inefficient cooling; Heating
- Shift imaging light out of resonance










- Inefficient cooling; Heating
- Shift imaging light out of resonance

Cs single atom loading

U			
$\lambda_{trap}(nm)$	922	935	970
Loading (%)	0	≈ 50	≈ 50









- Alternate between resonant and trap light
- Switching at $1 \sim 3$ MHz $f_{trap} = 10 \sim 400 \text{ kHz}$ $\Gamma = 2\pi \times (5 \sim 10) \text{ MHz}$
- Being able to load single Na atom

- Alternate between resonant and trap light
- Switching at $1 \sim 3$ MHz $f_{trap} = 10 \sim 400$ kHz $\Gamma = 2\pi \times (5 \sim 10)$ MHz
- Being able to load single Na atom




- Alternate between resonant and trap light
- Switching at $1 \sim 3 \text{MHz}$ $f_{trap} = 10 \sim 400 \text{ kHz}$ $\Gamma = 2\pi \times (5 \sim 10) \text{ MHz}$
- Being able to load single
 Na atom



- Alternate between resonant and trap light
- Switching at $1 \sim 3$ MHz $f_{trap} = 10 \sim 400$ kHz $\Gamma = 2\pi \times (5 \sim 10)$ MHz
- Being able to load single Na atom

Cs single atom loading

		<i>U</i>	
$\lambda_{trap}(nm)$	922	935	970
Loading (%)	≈ 50	≈ 50	≈ 50




- Alternate between resonant and trap light
- Switching at $1 \sim 3 \text{MHz}$ $f_{trap} = 10 \sim 400 \text{ kHz}$ $\Gamma = 2\pi \times (5 \sim 10) \text{ MHz}$

Cs single atom loading

$\lambda_{trap}(nm)$	922	935	970
Loading (%)	≈ 50	≈ 50	≈ 50

Cs single atom imaging




- Alternate between resonant and trap light
- Switching at $1 \sim 3 \text{MHz}$ $f_{trap} = 10 \sim 400 \text{ kHz}$ $\Gamma = 2\pi \times (5 \sim 10) \text{ MHz}$


Cs single atom loading


$\lambda_{trap}(nm)$	922	935	970
Loading (%)	≈ 50	≈ 50	≈ 50

Cs single atom imaging





- Alternate between resonant and trap light
- Switching at $1 \sim 3 \text{MHz}$ $f_{trap} = 10 \sim 400 \text{ kHz}$ $\Gamma = 2\pi \times (5 \sim 10) \text{ MHz}$
- Being able to load single Na atom





- Alternate between resonant and trap light
- Switching at $1 \sim 3 \text{MHz}$ $f_{trap} = 10 \sim 400 \text{ kHz}$ $\Gamma = 2\pi \times (5 \sim 10) \text{ MHz}$
- Being able to load single Na atom





Detuning (MHz)

5000

Conclusion

- Measured the effect of light shift on loading and imaging of single atom
- Overcome the light shift by alternating trapping and resonant light to achieve loading of single Na atom.
- Generalizable to other species

