

# Association of single ultracold molecules in optical tweezers

Yichao Yu, Jonathan Hood, Lee R. Liu, Jessie T. Zhang, Yen-Wei Lin, Kenneth Wang, Remý Vatré, Till Rosenband, Kang-Kuen Ni





Department of Chemistry and Chemical Biology and Department of Physics, Harvard University



#### **Ultracold Molecules**

- NaCs has a large permanent electric dipole moment (4.6 Debye)
- Strong anisotropic dipole-dipole interactions
- Coupled internal degrees of freedom can be used to tune interactions and store information






# Our Approach

- Assemble and trap individual molecules in optical tweezers from laser-cooled atoms
- Raman transition from atoms to weakly-bound molecules
- STIRAP to ground state molecules

## **Advantages**

- Fast cycle time (<1s), small vacuum chamber
- Dynamically configurable trapping geometry
- All optical cooling and state-manipulation



## Acknowledgements








## **Trapping and Cooling of Atoms**

Laser-cooled and trapped single Cs and Na atoms  $< 100 \, \, \text{uK}$  in separate rearrangeable optical tweezers.



Cs Raman spectrum

 $P_{0,3D} = 44(5)\%$ 

 $P_{0,3D} = 96(3)\%$ 

-150 -100 -50 0 50 100 150

 $\delta$ , Detuning from carrier (kHz)

Na axial Raman spectrum


 $\delta$ , Detuning from carrier (kHz)

 $\rightarrow$  z initial

 $\rightarrow$  z cooled


<u>6</u> 0.4

0.8 0.6 0.4



Hutzler, Liu, Yu, Ni, New J. Phys 19, 023007 (2017) L. Liu, Zhang, Yu, Hutzler, Hood, Liu, Rosenband, Ni, arXiv:1701.03121 (2017)


Cooled into motional ground states in the tweezers with Raman sideband cooling. Cooling fidelities are 96% for Cesium and 94% for Sodium.



Yu, Hutzler, Zhang, L. Liu, Rosenband, Ni, arXiv:1708.03296 (2017)


# Merging Tweezers

- Na and Cs tweezers are merged together into a single trap with little heating
- Observe hyperfine spin changing collisions when atoms in same trap



## **Photoassociation**


## Near threshold photoassociation



- First photoassociation of a single molecule in an optical tweezer.
- Observation of new NaCs excited state vibrational lines near threshold.
- Vertical lines are fits to the LeRoy long range dispersion model,

$$\Delta E \propto \frac{1}{\sqrt{C_6}} (v_D - v)^3$$

Liu, Hood, Yu, Zhang, Hutzler Rosenband, Ni, Science 360, 6391 (2018)



## **Coherent molecule formation** (Preliminary)

