

A One-Way Quantum Computer


Yichao Yu

Ni Group

Feb. 15, 2021

- 1D Cluster state
 - ▶ Generation
 - ▶ Properties
- High dimensional cluster state
- Quantum circuit
- Gates and single qubit operations

1D Cluster State



$$H = \sum_{a, a' \in \Gamma} \frac{1 + \sigma_z^{(a)}}{2} \frac{1 - \sigma_z^{(a')}}{2}$$

$$\Gamma = \{(a, a') | a' = a + 1\}$$

$$\mathcal{S} = e^{i\pi H}$$

1D Cluster State



$$H = \sum_{a, a' \in \Gamma} \frac{1 + \sigma_z^{(a)}}{2} \frac{1 - \sigma_z^{(a')}}{2}$$
$$\Gamma = \{(a, a') | a' = a + 1\}$$
$$H = \sum \begin{pmatrix} 0 & & & \\ & 0 & & \\ & & 1 & \\ & & & 0 \end{pmatrix}$$

$$\mathcal{S} = e^{i\pi H}$$
$$\mathcal{S} = \bigotimes \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & -1 & \\ & & & 1 \end{pmatrix}$$

1D Cluster State



$$|\phi_N\rangle = \mathcal{S} \bigotimes_a |+\rangle_a = \frac{1}{2^{N/2}} \bigotimes_a (|0\rangle_a \sigma_z^{a+1} + |1\rangle_a)$$

1D Cluster State



$$|\phi_N\rangle = \mathcal{S} \bigotimes_a |+\rangle_a = \frac{1}{2^{N/2}} \bigotimes_a (|0\rangle_a \sigma_z^{a+1} + |1\rangle_a)$$

$$|\phi_2\rangle = \frac{1}{\sqrt{2}}(|0-\rangle + |1+\rangle)$$

1D Cluster State



$$|\phi_N\rangle = \mathcal{S} \bigotimes_a |+\rangle_a = \frac{1}{2^{N/2}} \bigotimes_a (|0\rangle_a \sigma_z^{a+1} + |1\rangle_a)$$

$$|\phi_2\rangle = \frac{1}{\sqrt{2}}(|0-\rangle + |1+\rangle)$$

$$|\phi_3\rangle = \frac{1}{\sqrt{2}}(|+0-\rangle - |-1+\rangle)$$

1D Cluster State



$$|\phi_N\rangle = \mathcal{S} \bigotimes_a |+\rangle_a = \frac{1}{2^{N/2}} \bigotimes_a (|0\rangle_a \sigma_z^{a+1} + |1\rangle_a)$$

$$|\phi_2\rangle = \frac{1}{\sqrt{2}}(|0-\rangle + |1+\rangle)$$

$$|\phi_3\rangle = \frac{1}{\sqrt{2}}(|+0-\rangle - |-1+\rangle)$$

$$|\phi_4\rangle = \frac{1}{2}(|0-0-\rangle - |0+1+\rangle + |1+0-\rangle - |1-1+\rangle)$$

1D Cluster State



$$|\phi_N\rangle = \mathcal{S} \bigotimes_a |+\rangle_a = \frac{1}{2^{N/2}} \bigotimes_a (|0\rangle_a \sigma_z^{a+1} + |1\rangle_a)$$

$$|\phi_2\rangle = \frac{1}{\sqrt{2}}(|0-\rangle + |1+\rangle)$$

$$|\phi_3\rangle = \frac{1}{\sqrt{2}}(|+0-\rangle - |-1+\rangle)$$

$$|\phi_4\rangle = \frac{1}{2}(|0-0-\rangle - |0+1+\rangle + |1+0-\rangle - |1-1+\rangle)$$

$$|\text{GHZ}_N\rangle = \frac{1}{\sqrt{2}} \left(\bigotimes_a |0\rangle_a + \bigotimes_a |1\rangle_a \right)$$

- Maximum connectedness
Ability to create Bell state by local measurements.
Yes for both GHZ state and cluster state.
- Persistency
Minimum local measurements to destroy all entanglements.
GHZ: $P_e = 1$, cluster: $P_e = \lfloor N/2 \rfloor$


- Maximum connectedness
Ability to create Bell state by local measurements.
Yes for both GHZ state and cluster state.
- Persistency
Minimum local measurements to destroy all entanglements.
GHZ: $P_e = 1$, cluster: $P_e = \lfloor N/2 \rfloor$

- Maximum connectedness
Ability to create Bell state by local measurements.
Yes for both GHZ state and cluster state.
- Persistency
Minimum local measurements to destroy all entanglements.
GHZ: $P_e = 1$, cluster: $P_e = \lfloor N/2 \rfloor$

1D Cluster State

- Maximum connectedness
Ability to create Bell state by local measurements.
Yes for both GHZ state and cluster state.
- Persistency
Minimum local measurements to destroy all entanglements.
GHZ: $P_e = 1$, cluster: $P_e = \lfloor N/2 \rfloor$

High Dimensional Cluster State



$$H = \sum_{a, a' \in \Gamma} \frac{1 + \sigma_z^{(a)}}{2} \frac{1 - \sigma_z^{(a')}}{2}$$

$$\Gamma = \{(a, a') | a' = a + \hat{e}_i\}$$

High Dimensional Cluster State


$$H = \sum_{a,a' \in \Gamma} \frac{1 + \sigma_z^{(a)}}{2} \frac{1 - \sigma_z^{(a')}}{2}$$

$$[H, \sigma_z^{(a)}] = 0, \quad [\mathcal{S}, \sigma_z^{(a)}] = 0$$

High Dimensional Cluster State

$$H = \sum_{a,a' \in \Gamma} \frac{1 + \sigma_z^{(a)}}{2} \frac{1 - \sigma_z^{(a')}}{2}$$


$$[H, \sigma_z^{(a)}] = 0, \quad [\mathcal{S}, \sigma_z^{(a)}] = 0$$




High Dimensional Cluster State

$$H = \sum_{a,a' \in \Gamma} \frac{1 + \sigma_z^{(a)}}{2} \frac{1 - \sigma_z^{(a')}}{2}$$

$$[H, \sigma_z^{(a)}] = 0, \quad [\mathcal{S}, \sigma_z^{(a)}] = 0$$




High Dimensional Cluster State



$$K_a = \sigma_x^{(a)} \bigotimes_{a' \in \Gamma'} \sigma_z^{(a')}$$

$$\Gamma' = \{(a, a') | a' = a \pm \hat{e}_i\}$$

High Dimensional Cluster State




$$K_a = \sigma_x^{(a)} \bigotimes_{a' \in \Gamma'} \sigma_z^{(a')}$$

$$\Gamma' = \{(a, a') | a' = a \pm \hat{e}_i\}$$

- $K_a|\phi_N\rangle = \pm|\phi_N\rangle$
- $\{K_a, \sigma_z^{(a)}\} = 0$
- $[K_a, K_b] = 0$
- $[K_a, \sigma_z^{(b)}] \Big|_{a \neq b} = 0$
- Independent
- Complete
- Equivalent definitions of cluster state
 - ▶ Apply full \mathcal{S} and measure σ_z 's on removed sites
 - ▶ Apply partial \mathcal{S} and apply σ_z 's on remaining sites
 - ▶ Eigenstates of all K_a

High Dimensional Cluster State




$$K_a = \sigma_x^{(a)} \bigotimes_{a' \in \Gamma'} \sigma_z^{(a')}$$

$$\Gamma' = \{(a, a') | a' = a \pm \hat{e}_i\}$$

- $K_a |\phi_N\rangle = \pm |\phi_N\rangle$
- $\{K_a, \sigma_z^{(a)}\} = 0$
- $[K_a, K_b] = 0$
- $[K_a, \sigma_z^{(b)}] \Big|_{a \neq b} = 0$
- Independent
- Complete
- Equivalent definitions of cluster state
 - ▶ Apply full \mathcal{S} and measure σ_z 's on removed sites
 - ▶ Apply partial \mathcal{S} and apply σ_z 's on remaining sites
 - ▶ Eigenstates of all K_a

High Dimensional Cluster State




$$K_a = \sigma_x^{(a)} \bigotimes_{a' \in \Gamma'} \sigma_z^{(a')}$$

$$\Gamma' = \{(a, a') | a' = a \pm \hat{e}_i\}$$

- $K_a |\phi_N\rangle = \pm |\phi_N\rangle$
- $\{K_a, \sigma_z^{(a)}\} = 0$
- $[K_a, K_b] = 0$
- $[K_a, \sigma_z^{(b)}] \Big|_{a \neq b} = 0$
- Independent
- Complete
- Equivalent definitions of cluster state
 - ▶ Apply full \mathcal{S} and measure σ_z 's on removed sites
 - ▶ Apply partial \mathcal{S} and apply σ_z 's on remaining sites
 - ▶ Eigenstates of all K_a

High Dimensional Cluster State




$$K_a = \sigma_x^{(a)} \bigotimes_{a' \in \Gamma'} \sigma_z^{(a')}$$

$$\Gamma' = \{(a, a') | a' = a \pm \hat{e}_i\}$$

- $K_a |\phi_N\rangle = \pm |\phi_N\rangle$
- $\{K_a, \sigma_z^{(a)}\} = 0$
- $[K_a, K_b] = 0$
- $[K_a, \sigma_z^{(b)}] \Big|_{a \neq b} = 0$
- Independent
- Complete
- Equivalent definitions of cluster state
 - ▶ Apply full \mathcal{S} and measure σ_z 's on removed sites
 - ▶ Apply partial \mathcal{S} and apply σ_z 's on remaining sites
 - ▶ Eigenstates of all K_a

High Dimensional Cluster State




$$K_a = \sigma_x^{(a)} \bigotimes_{a' \in \Gamma'} \sigma_z^{(a')}$$

$$\Gamma' = \{(a, a') | a' = a \pm \hat{e}_i\}$$

- $K_a |\phi_N\rangle = \pm |\phi_N\rangle$
- $\{K_a, \sigma_z^{(a)}\} = 0$
- $[K_a, K_b] = 0$
- $[K_a, \sigma_z^{(b)}] \Big|_{a \neq b} = 0$
- Independent
- Complete
- Equivalent definitions of cluster state
 - ▶ Apply full \mathcal{S} and measure σ_z 's on removed sites
 - ▶ Apply partial \mathcal{S} and apply σ_z 's on remaining sites
 - ▶ Eigenstates of all K_a

High Dimensional Cluster State




$$K_a = \sigma_x^{(a)} \bigotimes_{a' \in \Gamma'} \sigma_z^{(a')}$$

$$\Gamma' = \{(a, a') | a' = a \pm \hat{e}_i\}$$

- $K_a |\phi_N\rangle = \pm |\phi_N\rangle$
- $\{K_a, \sigma_z^{(a)}\} = 0$
- $[K_a, K_b] = 0$
- $[K_a, \sigma_z^{(b)}] \Big|_{a \neq b} = 0$
- Independent
- Complete
- Equivalent definitions of cluster state
 - ▶ Apply full \mathcal{S} and measure σ_z 's on removed sites
 - ▶ Apply partial \mathcal{S} and apply σ_z 's on remaining sites
 - ▶ Eigenstates of all K_a

High Dimensional Cluster State




$$K_a = \sigma_x^{(a)} \bigotimes_{a' \in \Gamma'} \sigma_z^{(a')}$$

$$\Gamma' = \{(a, a') | a' = a \pm \hat{e}_i\}$$

- $K_a |\phi_N\rangle = \pm |\phi_N\rangle$
- $\{K_a, \sigma_z^{(a)}\} = 0$
- $[K_a, K_b] = 0$
- $[K_a, \sigma_z^{(b)}] \Big|_{a \neq b} = 0$
- Independent
- Complete
- Equivalent definitions of cluster state
 - ▶ Apply full \mathcal{S} and measure σ_z 's on removed sites
 - ▶ Apply partial \mathcal{S} and apply σ_z 's on remaining sites
 - ▶ Eigenstates of all K_a

High Dimensional Cluster State




$$K_a = \sigma_x^{(a)} \bigotimes_{a' \in \Gamma'} \sigma_z^{(a')}$$

$$\Gamma' = \{(a, a') | a' = a \pm \hat{e}_i\}$$

- $K_a |\phi_N\rangle = \pm |\phi_N\rangle$
- $\{K_a, \sigma_z^{(a)}\} = 0$
- $[K_a, K_b] = 0$
- $[K_a, \sigma_z^{(b)}] \Big|_{a \neq b} = 0$
- Independent
- Complete
- Equivalent definitions of cluster state
 - ▶ Apply full \mathcal{S} and measure σ_z 's on removed sites
 - ▶ Apply partial \mathcal{S} and apply σ_z 's on remaining sites
 - ▶ Eigenstates of all K_a

High Dimensional Cluster State




$$K_a = \sigma_x^{(a)} \bigotimes_{a' \in \Gamma'} \sigma_z^{(a')}$$

$$\Gamma' = \{(a, a') | a' = a \pm \hat{e}_i\}$$

- $K_a |\phi_N\rangle = \pm |\phi_N\rangle$
- $\{K_a, \sigma_z^{(a)}\} = 0$
- $[K_a, K_b] = 0$
- $[K_a, \sigma_z^{(b)}] \Big|_{a \neq b} = 0$
- Independent
- Complete
- Equivalent definitions of cluster state
 - ▶ Apply full \mathcal{S} and measure σ_z 's on removed sites
 - ▶ Apply partial \mathcal{S} and apply σ_z 's on remaining sites
 - ▶ Eigenstates of all K_a

High Dimensional Cluster State




$$K_a = \sigma_x^{(a)} \bigotimes_{a' \in \Gamma'} \sigma_z^{(a')}$$


$$\Gamma' = \{(a, a') | a' = a \pm \hat{e}_i\}$$

- $K_a |\phi_N\rangle = \pm |\phi_N\rangle$
- $\{K_a, \sigma_z^{(a)}\} = 0$
- $[K_a, K_b] = 0$
- $[K_a, \sigma_z^{(b)}] \Big|_{a \neq b} = 0$
- Independent
- Complete
- Equivalent definitions of cluster state
 - ▶ Apply full \mathcal{S} and measure σ_z 's on removed sites
 - ▶ Apply partial \mathcal{S} and apply σ_z 's on remaining sites
 - ▶ Eigenstates of all K_a


Quantum Circuit on Cluster State




Quantum Circuit on Cluster State



Quantum Circuit on Cluster State



Quantum Circuit on Cluster State




Gates and Single Qubit Operations (Single Qubit)

Propagation: Measure σ_x

Gates and Single Qubit Operations (Single Qubit)


Propagation: Measure σ_x



Step 1


Gates and Single Qubit Operations (Single Qubit)

Propagation: Measure σ_x



Gates and Single Qubit Operations (Single Qubit)


Propagation: Measure σ_x




$$\begin{aligned} \mathcal{S}(a|0\rangle_1 + b|1\rangle_1)|+\rangle_2 &= |+\rangle_1(a|-\rangle_2 + b|+\rangle_2) \\ &\quad + |-\rangle_1(a|-\rangle_2 - b|+\rangle_2) \end{aligned}$$

Gates and Single Qubit Operations (Single Qubit)

Propagation: Measure σ_x




$$\begin{aligned} \mathcal{S}(a|0\rangle_1 + b|1\rangle_1)|+\rangle_2 &= |+\rangle_1(a|-\rangle_2 + b|+\rangle_2) \\ &\quad + |-\rangle_1(a|-\rangle_2 - b|+\rangle_2) \end{aligned}$$




Gates and Single Qubit Operations (Single Qubit)

Propagation: Measure σ_x




$$\begin{aligned} \mathcal{S}(a|0\rangle_1 + b|1\rangle_1)|+\rangle_2 = & |+\rangle_1(a|-\rangle_2 + b|+\rangle_2) \\ & + |-\rangle_1(a|-\rangle_2 - b|+\rangle_2) \end{aligned}$$




Rotation: Measure $\sigma_x \cos \theta + \sigma_y \sin \theta$



Gates and Single Qubit Operations (CNOT)



Gates and Single Qubit Operations (CNOT)




Gates and Single Qubit Operations (CNOT)



$$S = \bigotimes \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & -1 & \\ & & & 1 \end{pmatrix}$$

Gates and Single Qubit Operations (CNOT)



$$S = \bigotimes \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & -1 & \\ & & & 1 \end{pmatrix}$$

Questions?