

Ultracold molecule assembly

Yichao Yu


Ni Group/Harvard

Aug 11, 2017

Molecules in optical tweezer

Features


- Strong and tunable interaction
- Rich internal energy levels
- High filling fraction
- Single site detection and manipulation



Molecules in optical tweezer

Features


- Strong and tunable interaction
- Rich internal energy levels
- High filling fraction
- Single site detection and manipulation



Molecules in optical tweezer

Features


- Strong and tunable interaction
- Rich internal energy levels
- High filling fraction
- Single site detection and manipulation



Molecules in optical tweezer

Features


- Strong and tunable interaction
- Rich internal energy levels
- High filling fraction
- Single site detection and manipulation




Molecules in optical tweezer

Features

- Strong and tunable interaction
- Rich internal energy levels
- High filling fraction
- Single site detection and manipulation




Simulation of many-body system^[1]



$$H \propto \sum_{ij} V_{ij} (S_i^+ S_j^- + S_i^- S_j^+)$$

[1] B. Yan et al., "Observation of dipolar spin-exchange interactions with lattice-confined polar molecules.", *Nature* **501**, 521–5 (2013).

Quantum computation^[2]



[2] S. F. Yelin et al., “Schemes for robust quantum computation with polar molecules”, *Phys. Rev. A* **74**, 050301 (2006).


Making molecules from atoms

- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!



Making molecules from atoms

- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!



Making molecules from atoms

- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!



Making molecules from atoms

- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!



Making molecules from atoms

- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!



Making molecules from atoms

- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!




Atom loading and cooling

- Single atoms
- 85% ground state after Cesium Raman sideband cooling




Atom loading and cooling


- Single atoms
- 85% ground state after Cesium Raman sideband cooling



Raman sideband cooling





Raman sideband cooling





Raman sideband cooling



$3^2P_{3/2}$





Raman sideband cooling



Raman sideband cooling




Raman sideband cooling




- High initial temperature ($70\mu K$)
- High Lamb Dicke parameter
 $\eta \equiv kz_0$
- Large light shift
- Trap anharmonicity
- Off resonance scattering
 $\approx 3 \sim 15\text{kHz}$

Motional state bran




- High initial temperature ($70\mu K$)
- High Lamb Dicke parameter
 $\eta \equiv kz_0$
- Large light shift
- Trap anharmonicity
- Off resonance scattering
 $\approx 3 \sim 15\text{kHz}$

Motional state branching ratios




- High initial temperature ($70\mu K$)
- High Lamb Dicke parameter
 $\eta \equiv kz_0$
- Large light shift
- Trap anharmonicity
- Off resonance scattering
 $\approx 3 \sim 15\text{kHz}$

Motional state branching ratios





Raman matrix elements




- High initial temperature ($70\mu K$)
- High Lamb Dicke parameter
 $\eta \equiv kz_0$
- Large light shift
- Trap anharmonicity
- Off resonance scattering
 $\approx 3 \sim 15\text{kHz}$



Motional state branching ratios





Raman matrix elements





- High initial temperature ($70\mu K$)
- High Lamb Dicke parameter
 $\eta \equiv kz_0$
- Large light shift
- Trap anharmonicity
- Off resonance scattering
 $\approx 3 \sim 15\text{kHz}$

Raman sideband cooling





- High initial temperature ($70\mu K$)
- High Lamb Dicke parameter
 $\eta \equiv kz_0$
- Large light shift
- Trap anharmonicity
- Off resonance scattering
 $\approx 3 \sim 15\text{kHz}$

Raman sideband cooling




- High initial temperature ($70\mu K$)
- High Lamb Dicke parameter
 $\eta \equiv kz_0$
- Large light shift
- Trap anharmonicity
- Off resonance scattering
 $\approx 3 \sim 15\text{kHz}$

Raman sideband cooling




- High initial temperature ($70\mu K$)
- High Lamb Dicke parameter
 $\eta \equiv kz_0$
- Large light shift
- Trap anharmonicity
- Off resonance scattering
 $\approx 3 \sim 15\text{kHz}$


Raman matrix elements




Raman matrix elements




Sequence and simulation





Sequence and simulation




Sequence and simulation












Raman sidebands




Raman sidebands





Raman sidebands





Raman sidebands



Rabi flopping (radial)





Rabi flopping (radial)



Good agreement in ground state probability
between spectrum and Rabi flopping data.

Rabi flopping (axial)



Decoherence caused by technical noise.
E.g. 1.5 mG of magnetic field noise.


Conclusion


- Trapping of Na and Cs atoms
- Ground state cooling of Na^[3] and Cs





In progress

- Merge trap
- Photoassociation spectroscopy
- Make molecules

[3] Y. Yu et al., “Motional ground state cooling outside the lamb-dicke regime”, [arXiv 1708.03296](#) (2017).









Quantum computation


—— Strong dipole

$|1\rangle$ ——— $|1\rangle$
 $|0\rangle$ ——— $|0\rangle$
Weak dipole

Quantum computation



Quantum computation




Quantum computation



$$\begin{pmatrix} E & \frac{V}{r^3} \\ \frac{V}{r^3} & E \end{pmatrix}$$


Quantum computation






The diagram shows two horizontal lines representing energy levels. The top line has two red dots, and the bottom line has two green dots. This represents a system with two degenerate states, each with a sub-level splitting.

$$\begin{pmatrix} E & \frac{V}{r^3} \\ \frac{V}{r^3} & E \end{pmatrix} \rightarrow \begin{pmatrix} E - \frac{V}{r^3} & \\ & E + \frac{V}{r^3} \end{pmatrix}$$


Before cooling



After cooling



Merge trap



Making molecule

