


# Trapping and imaging of single atoms in the presence of light shift

Yichao Yu

May 26, 2016

Ni Group/Harvard



## Group members

Nicholas Hutzler

Lee Liu

Jessie Zhang

PI

Kang-Kuen Ni



BECKMAN  
FOUNDATION

## Procedure


- MOT Loading
- Trapping
- Imaging
- Works for Cs
- Doesn't work for Na



## Procedure


- MOT Loading
- Trapping
- Imaging
- Works for Cs
- Doesn't work for Na

Tweezer




## Procedure

- MOT Loading
- Trapping
- Imaging
- Works for Cs
- Doesn't work for Na



## Procedure

- MOT Loading
- Trapping
- Imaging
- Works for Cs
- Doesn't work for Na



## Procedure

- MOT Loading
- Trapping
- Imaging
- Works for Cs
- Doesn't work for Na



## Light shift

- Inefficient cooling;  
Heating
- Shift imaging light out of  
resonance



## Light shift


- Inefficient cooling;  
Heating
- Shift imaging light out of  
resonance






## Light shift

- Inefficient cooling;  
Heating
- Shift imaging light out of resonance





## Light shift

- Inefficient cooling;  
Heating
- Shift imaging light out of resonance





## Light shift

- Inefficient cooling;  
Heating
- Shift imaging light out of resonance





## Light shift

- Inefficient cooling;  
Heating
- Shift imaging light out of resonance




## Light shift

- Inefficient cooling;  
Heating
- Shift imaging light out of resonance



## Light shift


- Inefficient cooling;  
Heating
- Shift imaging light out of resonance



## Cs single atom loading

| $\lambda_{\text{trap}}$ (nm) | 922 | 935          | 970          |
|------------------------------|-----|--------------|--------------|
| Loading (%)                  | 0   | $\approx 50$ | $\approx 50$ |

## Cs single atom imaging




## Trap modulation

Alternate between trap and  
resonant (cooling and imaging)  
light at 1 ~ 3 MHz

$$f_{\text{trap}} = 10 \sim 400 \text{ kHz}$$

$$\Gamma = 2\pi \times (5 \sim 10) \text{ MHz}$$




## Trap modulation

Alternate between trap and resonant (cooling and imaging) light at 1 ~ 3 MHz

$$f_{\text{trap}} = 10 \sim 400 \text{ kHz}$$

$$\Gamma = 2\pi \times (5 \sim 10) \text{ MHz}$$



## Cs single atom loading

|                                    |              |              |              |
|------------------------------------|--------------|--------------|--------------|
| $\lambda_{\text{trap}}(\text{nm})$ | 922          | 935          | 970          |
| Loading (%)                        | $\approx 50$ | $\approx 50$ | $\approx 50$ |




## Trap modulation

Alternate between trap and resonant (cooling and imaging) light at 1 ~ 3 MHz

$$f_{\text{trap}} = 10 \sim 400 \text{ kHz}$$


$$\Gamma = 2\pi \times (5 \sim 10) \text{ MHz}$$



## Cs single atom loading

| $\lambda_{\text{trap}}(\text{nm})$ | 922          | 935          | 970          |
|------------------------------------|--------------|--------------|--------------|
| Loading (%)                        | $\approx 50$ | $\approx 50$ | $\approx 50$ |

## Cs single atom imaging




## Trap modulation

Alternate between trap and resonant (cooling and imaging) light at 1 ~ 3 MHz

$$f_{\text{trap}} = 10 \sim 400 \text{ kHz}$$


$$\Gamma = 2\pi \times (5 \sim 10) \text{ MHz}$$



## Cs single atom loading

| $\lambda_{\text{trap}}(\text{nm})$ | 922          | 935          | 970          |
|------------------------------------|--------------|--------------|--------------|
| Loading (%)                        | $\approx 50$ | $\approx 50$ | $\approx 50$ |

## Cs single atom imaging





## Trap modulation

Alternate between trap and  
resonant (cooling and imaging)  
light at 1 ~ 3 MHz

$$f_{\text{trap}} = 10 \sim 400 \text{ kHz}$$

$$\Gamma = 2\pi \times (5 \sim 10) \text{ MHz}$$





Na Single Atom Loaded!!

## Trap modulation

Alternate between trap and resonant (cooling and imaging) light at 1 ~ 3 MHz

$$f_{\text{trap}} = 10 \sim 400 \text{ kHz}$$

$$\Gamma = 2\pi \times (5 \sim 10) \text{ MHz}$$



Na Single Atom Loaded!!

## Conclusion

- Measured the effect of light shift on loading and imaging of single atom
- Overcome the light shift by alternating trapping and resonant light to achieve loading of single Na atom.
- Generalizable to other species

