
Ultracold molecule assembly

Yichao Yu


Ni Group/Harvard

Aug 11, 2017


- Strong and tunable interaction
- Rich internal energy levels
- High filling fraction
- Single site detection and manipulation




- Strong and tunable interaction
- Rich internal energy levels
- High filling fraction
- Single site detection and manipulation




- Strong and tunable interaction
- Rich internal energy levels
- High filling fraction
- Single site detection and manipulation



- Strong and tunable interaction
- Rich internal energy levels
- High filling fraction
- Single site detection and manipulation




- Strong and tunable interaction
- Rich internal energy levels
- High filling fraction
- Single site detection and manipulation



Applications


Simulation of many-body system^[1]



$$H \propto \sum V_{ij} \left(S_i^+ S_j^- + S_i^- S_j^+ \right)$$

[1] B. Yan et al., "Observation of dipolar spin-exchange interactions with lattice-confined polar molecules.", Nature **501**, 521–5 (2013).

Applications




[2] S. F. Yelin et al., "Schemes for robust quantum computation with polar molecules", Phys. Rev. A 74, 050301 (2006).

- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!



- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!



- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!

- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!


- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!


- MOT (Na + Cs)
- Loading single atoms
- Raman sideband cooling
- Merge traps
- Make molecules!

Atom loading and cooling

Single atoms


 85% ground state after Cesium Raman sideband cooling







Atom loading and cooling

- Single atoms
- 85% ground state after Cesium Raman sideband cooling










6/22


$$3^2P_{3/2}$$





$$3^{2}S_{1/2}$$
 $|2, -2; n\rangle$
 $|2, -2; n-1\rangle$


















- High initial temperature $(70\mu K)$
 - High Lamb Dicke parameter $\eta \equiv kz_0$
- Large light shift
- Trap anharmonicity
- $|1, -1; n-1\rangle$ Off resonance scattering $\approx 0.2 \sim 0.5 \text{kHz}$







- High initial temperature $(70\mu K)$
- High Lamb Dicke parameter $\eta \equiv kz_0$
- Large light shift
- Trap anharmonicity
- Off resonance scattering $\approx 0.2 \sim 0.5 \text{kHz}$

Motional state branching ratios $n_{init} = 0$ Probability $0.6 \cdot n_{init} = 1$ 30 60 70 Motional state n





- High initial temperature $(70\mu K)$
- High Lamb Dicke parameter $\eta \equiv kz_0$
- Trap anharmonicity
- Off resonance scattering






- High initial temperature $(70\mu K)$
- High Lamb Dicke parameter $\eta \equiv kz_0$
- Large light shift
- Trap anharmonicity
- Off resonance scattering $\approx 0.2 \sim 0.5 \text{kHz}$













- High initial temperature $(70\mu K)$
- High Lamb Dicke parameter $\eta \equiv kz_0$
- Large light shift
- Trap anharmonicity
- Off resonance scattering $\approx 0.2 \sim 0.5 \text{kHz}$







- High initial temperature $(70\mu K)$
- High Lamb Dicke parameter $\eta \equiv kz_0$
- Large light shift
- Trap anharmonicity
- $|1, -1; n-1\rangle$ Off resonance scattering $\approx 0.2 \sim 0.5 \text{kHz}$







- High initial temperature $(70\mu K)$
- High Lamb Dicke parameter $\eta \equiv kz_0$
- Large light shift
- Trap anharmonicity
- $|1, -1; n-1\rangle$ Off resonance scattering $\approx 0.2 \sim 0.5 \text{kHz}$







- High initial temperature $(70\mu K)$
- High Lamb Dicke parameter $\eta \equiv kz_0$
- Large light shift
- Trap anharmonicity
- $|1,-1;n-1\rangle$ Off resonance scattering $\approx 0.2 \sim 0.5 \text{kHz}$












Raman sidebands




Raman sidebands




Raman sidebands










Axis	Ground state probability
1 (Axial)	93.1(2.5)%
2 (Radial)	91.9(2.3)%
3 (Radial)	92.9(2.5)%


3D ground state: 79.5(3.6)%Loss after cooling: 15%

Total 3D ground state preparation fidelity: 67.6(3.1)%

Rabi flopping (radial)




Rabi flopping (radial)



Good agreement in ground state probability between spectrum and Rabi flopping data.

Rabi flopping (axial)




In progress

Aug 11, 2017









——— Strong dipole


Weak dipole

Quantum computation Strong dipole Weak dipole



Aug 11, 2017





Merge trap

