

# Homework 2

Naman Mishra (22223)

20 August, 2024

## Problem 1.

(1) Let  $\Omega$  be a set and  $A \subseteq \Omega$ . Define a function  $\mathbf{1}_A: \Omega \rightarrow \mathbb{R}$  as follows.

$$\mathbf{1}_A(\omega) = \begin{cases} 1 & \text{if } \omega \in A \\ 0 & \text{if } \omega \notin A. \end{cases}$$

What is the smallest  $\sigma$ -algebra on  $\Omega$  with respect to which  $\mathbf{1}_A$  becomes a random variable?

(2) Assume that  $A \in \mathcal{F}$ . Give an explicit description of the push-forward measure  $P \circ (\mathbf{1}_A)^{-1}$  on  $\mathbb{R}$ .

*Solution.*

(1) We need  $\mathbf{1}_A^{-1}(B) \in \mathcal{F}$  for  $B \in \mathcal{B}(\mathbb{R})$ .

$$(\mathbf{1}_A)^{-1}(B) = \begin{cases} \emptyset & \text{if } 0, 1 \notin B, \\ A & \text{if } 1 \in B, 0 \notin B, \\ A^c & \text{if } 0 \in B, 1 \notin B, \\ \Omega & \text{if } 0, 1 \in B. \end{cases}$$

Thus  $\mathcal{F}$  must contain  $\emptyset, A, A^c, \Omega$ .  $\mathcal{F} = \{\emptyset, A, A^c, \Omega\}$  is itself a  $\sigma$ -algebra, hence the smallest one that works.

(2) Let  $B \in \mathcal{B}(\mathbb{R})$ . Then

$$(P \circ (\mathbf{1}_A)^{-1})(B) = \begin{cases} 0 & \text{if } 0, 1 \notin B, \\ P(A) & \text{if } 1 \in B, 0 \notin B, \\ P(A^c) & \text{if } 0 \in B, 1 \notin B, \\ 1 & \text{if } 0, 1 \in B. \end{cases}$$

■

**Problem 2.** Recall the Lévy metric  $d$  defined in class. Show the following.

- (1) Let  $a_n$  be a sequence of real numbers converging to  $a$ . For any  $x \in \mathbb{R}$ ,  $\delta_x$  is the measure defined as follows: for  $A \subseteq \mathbb{R}$ ,

$$\delta_x(A) = \begin{cases} 1 & \text{if } x \in A, \\ 0 & \text{if } x \notin A. \end{cases}$$

Using the definition of the metric show that

$$d(\delta_{a_n}, \delta_a) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

- (2) Consider the sequence of measures  $\mu_n := \frac{1}{n} \sum_{i=1}^n \delta_{i/n}$  and  $\mu$  is the uniform measure on  $[0, 1]$ . Using the definition show that

$$d(\mu_n, \mu) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

*Solution.*

- (1) The CDF of  $\delta_x$  is  $F_x(\omega) = [\omega \geq x]$ . If  $|x - y| = \varepsilon$ , then

$$F_x(\omega + \varepsilon) = [\omega + \varepsilon \geq x] = [\omega \geq x - \varepsilon] \geq [\omega \geq y] = F_y(\omega)$$

since  $\omega \geq y \implies \omega \geq x - |x - y|$ . Thus  $d(\delta_x, \delta_y) \leq |x - y|$ . As  $a_n \rightarrow a$ ,  $d(\delta_{a_n}, \delta_a) \rightarrow 0$ .

- (2) The CDF of  $\mu$  is  $F(x) = x$  for  $x \in [0, 1]$ . The CDF of  $\mu_n$  is

$$F_n(x) = \frac{\lfloor nx \rfloor}{n} \quad \text{for } x \in [0, 1].$$

( $\lfloor nx \rfloor$  counts the number of points  $i/n \leq x$ , and each of those has weight  $1/n$ .) We claim that  $d(\mu_n, \mu) \leq 1/n$ .

Let  $x \in [0, 1]$ . Then

$$\begin{aligned} F\left(x + \frac{1}{n}\right) + \frac{1}{n} &= x + \frac{2}{n} \\ &= \frac{nx + 2}{n} \\ &> \frac{\lfloor nx \rfloor}{n} = F_n(x). \end{aligned}$$

and

$$\begin{aligned}
 F_n\left(x + \frac{1}{n}\right) + \frac{1}{n} &= \frac{\lfloor n(x + 1/n) \rfloor + 1}{n} \\
 &= \frac{\lfloor nx \rfloor + 2}{n} \\
 &> \frac{nx}{n} \\
 &= x = F(x).
 \end{aligned}$$

Thus

$$\frac{1}{n} \in \{\varepsilon > 0 : F_n(x + \varepsilon) + \varepsilon \geq F(x) \text{ and}$$

$$F(x + \varepsilon) + \varepsilon \geq F_n(x) \text{ for all } x \in [0, 1]\}$$

and so  $d(\mu_n, \mu)$ , which is the infimum of all such  $\varepsilon$ , is at most  $1/n$ . It follows that  $\lim_{n \rightarrow \infty} d(\mu_n, \mu) = 0$  by the squeeze theorem. ■

**Problem 3.** For  $k \geq 0$ , define the functions  $r_k: [0, 1) \rightarrow \mathbb{R}$  by writing  $[0, 1) = \bigsqcup_{0 \leq j < 2^k} I_j^{(k)}$  where  $I_j^{(k)}$  is the dyadic interval  $[j2^{-k}, (j+1)2^{-k})$  and setting

$$r_k(x) = \begin{cases} -1 & \text{if } x \in I_j^{(k)} \text{ for odd } j, \\ 1 & \text{if } x \in I_j^{(k)} \text{ for even } j. \end{cases}$$

Fix  $n \geq 1$  and define  $T_n: [0, 1) \rightarrow \{-1, 1\}^n$  by  $T_n(x) = (r_0(x), \dots, r_{n-1}(x))$ . Find the push-forward of the Lebesgue measure on  $[0, 1)$  under  $T_n$ .