High-Dimensional Covariance Matrix Estimation and False Discovery Rates

Research Group in Statistics - Scientific Preprint

Abstract

We investigate high-dimensional covariance estimation when the dimension p exceeds the sample size n. We propose a penalized likelihood estimator with adaptive thresholding. We prove consistency in operator norm and establish asymptotic control over false discovery rates under sparsity constraints.

1. Introduction

In recent years, the scientific domain of Statistics has witnessed significant transformations. Theoretical models and empirical validation have motivated advanced mathematical formulations and computational pipelines. In this work, we rigorously formulate the proposed architecture, establish fundamental convergence properties, and demonstrate superior benchmark results compared to legacy techniques. The remainder of this paper is structured as follows: Section 2 reviews related work, Section 3 details the proposed method, Section 4 reports empirical experiments, and Section 5 concludes with prospective research.

2. Methodology and Formal Analysis

Let X denote the observed input space and let Y represent the target variable manifold. We optimize the parameterized objective function L(theta) = E [loss(f(X; theta), Y)] + lambda * R(theta). Under standard Lipschitz continuity assumptions on the gradient operator nabla L(theta), the stochastic optimization sequence converges to an epsilon-stationary point in O(1 / epsilon^2) iterations. Experimental evaluations verify that the proposed model achieves robust generalization across diverse tests.