- News & Perspective
- · Drugs & Diseases
- CME & Education
- Specialty Oncology
  - Allergy & Immunology
  - Anesthesiology
  - Business of Medicine
  - Cardiology
  - Critical Care
  - Dermatology
  - Diabetes & Endocrinology
  - Emergency Medicine
  - Family Medicine
  - Gastroenterology
  - General Surgery
  - Hematology-Oncology
  - HIV/AIDS
  - Infectious Diseases
  - Internal Medicine
  - Multispecialty
  - Nephrology
  - Neurology
  - o Ob/Gyn & Women's Health
  - Oncology
  - Ophthalmology
  - Orthopedics
  - Pathology & Lab Medicine
  - Pediatrics
  - Plastic Surgery
  - Psychiatry
  - Public Health
  - Pulmonary Medicine
  - Radiology
  - Rheumatology
  - Transplantation
  - Urology
    - Medical Students
      - Nurses
      - Pharmacists
      - Today on Medscape
- · Edition English
  - English
  - Français
  - Deutsch
- · My Account Dr. A Garg
- Log Out

#### close

Please confirm that you would like to log out of Medscape. If you log out, you will be required to enter your username and password the next time you visit. Log out Cancel

Search Medscape



**News & Perspective Oncology** 

Reuters Health Information

# Treatment Delays Common in Head and Neck Cancer Patients

By Will Boggs MD December 02, 2015



NEW YORK (Reuters Health) - About 25% of patients with head and neck squamous cell cancer (HNSCC) encounter delays in treatment initiation that could worsen their chances of survival, according to results from the National Cancer Data Base (NCDB).

Time to treatment initiation (TTI) may be increased to allow for sophisticated and personalized therapy, but whether such delays adversely affect outcomes is not known.

Dr. Thomas J. Galloway and colleagues from Fox Chase Cancer Center, Philadelphia, used NCDB data to estimate the impact of increasing TTI on survival in a study of 51,655 patients with HNSCC.

Overall, TTI was between 0 and 30 days for 60% of patients, between 31 and 60 days for 30% of patients, between 61 and 90 days for 7% of patients, and more than 90 days for 3% of patients, according to a report released November 30 by the Journal of Clinical Oncology and scheduled for publication online.

Repeated simulations identified two optimal TTI thresholds with ranges of 46 to 52 days and 62 to 67 days, with survival decreasing significantly beyond those ranges.

Median overall survival was 73.5 months for TTI of 46 to 52 days, 61 months for TTI of 53 to 67 days, and 46.6 months for TTI beyond 67 days (p<0.001).

In 2011, 9.6% of patients had TTI greater than 67 days, and 25% had TTI greater than 46 days. The percentages were even higher for patients treated at academic centers (29.1% of those patients had TTI greater than 46 days, and TTI exceeded 46 days for 40% of patients who received primary chemoradiotherapy).

The effect of TTI was more pronounced in early-stage than advanced-stage disease. Despite more patients with delayed TTI, academic facilities had slightly better overall survival than did community hospitals (as did comprehensive community centers).

"Risk associated with increasing TTI currently affects a substantial percentage of patients with HNSCC (25%) in the United States," the researchers conclude. "These results suggest that efforts should be made to ensure that patients with HNSCC initiate appropriate treatment in a timely manner to avoid potentially detrimental deferral of treatment. Patients undergoing a transition in care may require greater coordination to reduce TTI."

"Recently piloted programs offering next-day appointments with cancer specialists address this reversible predictor of mortality and may partially alleviate increasing TTI," they add. "Without such reforms, it is conceivable that outcomes will continue to worsen because of prolonged TTI."

Dr. Ezra E. Cohen, from the University of California, San Diego, Moores Cancer Center, La Jolla, who was not involved in this study, told Reuters Health by email, "It is an important paper that really underscores what many of us have been saying for years. In a setting where we are attempting to cure patients, the sooner we start treatment, the better."

"Coordination of care for head and neck cancer (HNC) patients is complex and needs to be accomplished efficiently and effectively," Dr. Cohen said. "Care for HNC patients requires a multidisciplinary team that includes surgeons, radiation oncologists, medical oncologists, speech/swallow therapists, radiologists, dentists, pharmacists, nurses, nutritionists, and more.

Optimal care for HNC patients should really be delivered in a center that has these facilities and personnel. It is an example of a disease whose treatment probably should be centralized if we are to achieve the best outcomes, i.e., cures."

Dr. Galloway could not be reached for comment.

Four coauthors reported relevant relationships.

SOURCE: http://bit.ly/1ekfGZD

J Clin Oncol 2015.

2 comments

Post as:

Dr. apurva garg

## **Post Comment**

Comments on Medscape are moderated and should be professional in tone and on topic. Please see our Commenting Guide for further information. We reserve the right to remove posts at our sole discretion.

# Latest in Hematology-Oncology

Chemotherapy for Germ Cell Tumor May Spur CML

New Survivorship Care Plan Template for Radiation Oncology

Anthracyclines Linked to Memory Problems in Breast Cancer

**Up-front Genotyping Reduces Fluoropyrimidine Toxicity** 

Controversial UK Cancer Drug Fund Will Be 'Reformed'

Reuters Health Information © 2015

Cite this article: Treatment Delays Common in Head and Neck Cancer Patients. Medscape. Nov 30, 2015.

- 0
- \_
- \_
- o more
- About
  - About Medscape
  - Privacy Policy
  - Terms of Use
  - Help Center
- Membership

- Email Newsletters
- Manage My Account

## Apps

- Medscape
- MedPulse News
- CME & Education
- Business of Medicine

## • WebMD Network

- WebMD
- MedicineNet
- eMedicineHealth
- RxList
- WebMD Corporate

## Editions

- English
- Français
- Deutsch

All material on this website is protected by copyright, Copyright © 1994-2015 by WebMD LLC. This website also contains material copyrighted by 3rd parties.

This website uses cookies to deliver its services as described in our Cookie Policy. By using this website, you agree to the use of cookies.

close