

# First Order Logic

Subhajit Roy  
`subhajit@cse.iitk.ac.in`

Computer Science and Engineering,  
Indian Institute of Technology, Kanpur  
<http://web.cse.iitk.ac.in/~subhajit/cs202/>

# Where are we?

## 1 Syntax

# First-Order Formula

First order term:  $t := c \mid x$

# First-Order Formula

First order term:  $t := c \mid x$

First order formula:

$\Phi := p(t_1, t_2, \dots, t_n) \mid \neg\Phi \mid \Phi \wedge \Phi \mid \Phi \vee \Phi \mid \exists x_1, x_2, \dots, x_n. \Phi \mid \forall x_1, x_2, \dots, x_n. \Phi$

# First-Order Formula

First order term:  $t := c \mid x$

First order formula:

$\Phi := p(t_1, t_2, \dots, t_n) \mid \neg\Phi \mid \Phi \wedge \Phi \mid \Phi \vee \Phi \mid \exists x_1, x_2, \dots, x_n. \Phi \mid \forall x_1, x_2, \dots, x_n. \Phi$

Generally we always add a predicate for  $=$ (equality)

# Predicates

Predicates are used to represent:

- Relations ( $R$ ):
  - $p(x_1, x_2, \dots, x_n) = T$  iff  $(x_1, x_2, \dots, x_n) \in \mathcal{R}$

# Predicates

Predicates are used to represent:

- Relations ( $R$ ):
  - $p(x_1, x_2, \dots, x_n) = T$  iff  $(x_1, x_2, \dots, x_n) \in \mathcal{R}$
- Functions ( $F$ ): relation in which each input has only one output.
  - $p(x_1, x_2, \dots, x_n) = T$  iff  $(x_1, x_2, \dots, x_n) \in \mathcal{F}$

# Predicates

Predicates are used to represent:

- Relations ( $R$ ):
  - $p(x_1, x_2, \dots, x_n) = T$  iff  $(x_1, x_2, \dots, x_n) \in \mathcal{R}$
- Functions ( $F$ ): relation in which each input has only one output.
  - $p(x_1, x_2, \dots, x_n) = T$  iff  $(x_1, x_2, \dots, x_n) \in \mathcal{F}$

The encoding is much nicer with relations and function than with raw predicates.



# First-Order Formula (modified)

## First order term

$$t := c \mid x \mid f(t_1, t_2, \dots, t_n)$$

# First-Order Formula (modified)

## First order term

$$t := c \mid x \mid f(t_1, t_2, \dots, t_n)$$

## Atomic formula

$$\alpha := r(t_1, t_2, \dots, t_n) \mid t_1 = t_2$$

# First-Order Formula (modified)

## First order term

$$t := c \mid x \mid f(t_1, t_2, \dots, t_n)$$

## Atomic formula

$$\alpha := r(t_1, t_2, \dots, t_n) \mid t_1 = t_2$$

## First order formula

$$\Phi := \alpha \mid \neg \Phi \mid \Phi \wedge \Phi \mid \Phi \vee \Phi \mid \exists x_1, x_2, \dots, x_n. \Phi \mid \forall x_1, x_2, \dots, x_n. \Phi$$

# First-Order Formula (modified)

## First order term

$$t := c \mid x \mid f(t_1, t_2, \dots, t_n)$$

## Atomic formula

$$\alpha := r(t_1, t_2, \dots, t_n) \mid t_1 = t_2$$

## First order formula

$$\Phi := \alpha \mid \neg \Phi \mid \Phi \wedge \Phi \mid \Phi \vee \Phi \mid \exists x_1, x_2, \dots, x_n. \Phi \mid \forall x_1, x_2, \dots, x_n. \Phi$$

Then,  $(R, F, C)$  is a **first order language** where  $r \in R$ ,  $f \in F$  and  $c \in C$ .

# Universe (Domain) of Discourse

- Set of elements are we are discussing about
- The quantifiers run over this set
- **Intentional** versus **Extensional** description of the domain:
  - Intentional: what properties that this domain holds (necessary and sufficient conditions)
  - Extensional: listing of all the elements in the set or a enumerative description that shows how the set can be constructed

# Examples

- Not all birds can fly

# Examples

- Not all birds can fly
  - $\neg(\forall x. B(x) \rightarrow F(x))$

# Examples

- Not all birds can fly
  - $\neg(\forall x. B(x) \rightarrow F(x))$
  - $\exists x. B(x) \wedge \neg F(x)$
  - Universe of discourse: all creatures



# Examples

- Not all birds can fly
  - $\neg(\forall x. B(x) \rightarrow F(x))$
  - $\exists x. B(x) \wedge \neg F(x)$
  - Universe of discourse: all creatures
- Andy and Paul have the same paternal grandfather.

# Examples

- Not all birds can fly
  - $\neg(\forall x. B(x) \rightarrow F(x))$
  - $\exists x. B(x) \wedge \neg F(x)$
  - Universe of discourse: all creatures
- Andy and Paul have the same paternal grandfather.
  - $\forall x \forall y \forall u \forall v. (F(x, y) \wedge F(y, a) \wedge F(u, v) \wedge F(v, p) \rightarrow x = u)$  [Note the relation  $F(x, y)$  has no knowledge that  $y$  is the only father of  $x$ ; so the following formulation says that “all grandfathers of Andy and Paul are same; using  $\exists$  would say that at least one grandfather of Andy and Paul are same: both formulations are correct with the implicit knowledge of a unique father.)

# Examples

- Not all birds can fly
  - $\neg(\forall x. B(x) \rightarrow F(x))$
  - $\exists x. B(x) \wedge \neg F(x)$
  - Universe of discourse: all creatures
- Andy and Paul have the same paternal grandfather.
  - $\forall x \forall y \forall u \forall v. (F(x, y) \wedge F(y, a) \wedge F(u, v) \wedge F(v, p) \rightarrow x = u)$  [Note the relation  $F(x, y)$  has no knowledge that  $y$  is the only father of  $x$ ; so the following formulation says that “all grandfathers of Andy and Paul are same; using  $\exists$  would say that at least one grandfather of Andy and Paul are same: both formulations are correct with the implicit knowledge of a unique father.)
  - $f(f(a)) = f(f(p))$  (with functions, captures the knowledge that one has only one father)
  - Universe of discourse: all people

# Examples

- Not all birds can fly
  - $\neg(\forall x. B(x) \rightarrow F(x))$
  - $\exists x. B(x) \wedge \neg F(x)$
  - Universe of discourse: all creatures
- Andy and Paul have the same paternal grandfather.
  - $\forall x \forall y \forall u \forall v. (F(x, y) \wedge F(y, a) \wedge F(u, v) \wedge F(v, p) \rightarrow x = u)$  [Note the relation  $F(x, y)$  has no knowledge that  $y$  is the only father of  $x$ ; so the following formulation says that “all grandfathers of Andy and Paul are same; using  $\exists$  would say that at least one grandfather of Andy and Paul are same: both formulations are correct with the implicit knowledge of a unique father.)
  - $f(f(a)) = f(f(p))$  (with functions, captures the knowledge that one has only one father)
  - Universe of discourse: all people
- Ann likes Mary's brothers

# Examples

- Not all birds can fly
  - $\neg(\forall x. B(x) \rightarrow F(x))$
  - $\exists x. B(x) \wedge \neg F(x)$
  - Universe of discourse: all creatures
- Andy and Paul have the same paternal grandfather.
  - $\forall x \forall y \forall u \forall v. (F(x, y) \wedge F(y, a) \wedge F(u, v) \wedge F(v, p) \rightarrow x = u)$  [Note the relation  $F(x, y)$  has no knowledge that  $y$  is the only father of  $x$ ; so the following formulation says that “all grandfathers of Andy and Paul are same; using  $\exists$  would say that at least one grandfather of Andy and Paul are same: both formulations are correct with the implicit knowledge of a unique father.)
  - $f(f(a)) = f(f(p))$  (with functions, captures the knowledge that one has only one father)
  - Universe of discourse: all people
- Ann likes Mary's brothers
  - Ambiguous! two possibilities:

# Examples

- Not all birds can fly
  - $\neg(\forall x. B(x) \rightarrow F(x))$
  - $\exists x. B(x) \wedge \neg F(x)$
  - Universe of discourse: all creatures
- Andy and Paul have the same paternal grandfather.
  - $\forall x \forall y \forall u \forall v. (F(x, y) \wedge F(y, a) \wedge F(u, v) \wedge F(v, p) \rightarrow x = u)$  [Note the relation  $F(x, y)$  has no knowledge that  $y$  is the only father of  $x$ ; so the following formulation says that “all grandfathers of Andy and Paul are same; using  $\exists$  would say that at least one grandfather of Andy and Paul are same: both formulations are correct with the implicit knowledge of a unique father.)
  - $f(f(a)) = f(f(p))$  (with functions, captures the knowledge that one has only one father)
  - Universe of discourse: all people
- Ann likes Mary's brothers
  - Ambiguous! two possibilities:
  - $\exists x. (B(x, m) \wedge L(a, x))$  (Mary likes one of several brothers)

# Groups

- We can use first-order logic to describe the universe of discourse, like it forms a group under some operator

# Groups

- We can use first-order logic to describe the universe of discourse, like it forms a group under some operator
- Group is a structure  $(G, op, \epsilon)$ 
  - $op$  is associative;
  - existence of a right-identity  $\epsilon$
  - existence of an right-inverse, i.e. for all  $x \in G$  there exists a right inverse  $y \in G$



# Groups

- We can use first-order logic to describe the universe of discourse, like it forms a group under some operator
- Group is a structure  $(G, op, \epsilon)$ 
  - $op$  is associative;
  - existence of a right-identity  $\epsilon$
  - existence of an right-inverse, i.e. for all  $x \in G$  there exists a right inverse  $y \in G$
- $\forall x, y, z. op(op(x, y), z) = \forall x, y, z. op(x, op(y, z))$

# Groups


- We can use first-order logic to describe the universe of discourse, like it forms a group under some operator
- Group is a structure  $(G, op, \epsilon)$ 
  - $op$  is associative;
  - existence of a right-identity  $\epsilon$
  - existence of an right-inverse, i.e. for all  $x \in G$  there exists a right inverse  $y \in G$
- $\forall x, y, z. op(op(x, y), z) = \forall x, y, z. op(x, op(y, z))$
- $\forall x. op(x, \epsilon) = x$

# Groups

- We can use first-order logic to describe the universe of discourse, like it forms a group under some operator
- Group is a structure  $(G, op, \epsilon)$ 
  - $op$  is associative;
  - existence of a right-identity  $\epsilon$
  - existence of an right-inverse, i.e. for all  $x \in G$  there exists a right inverse  $y \in G$
- $\forall x, y, z. op(op(x, y), z) = \forall x, y, z. op(x, op(y, z))$
- $\forall x. op(x, \epsilon) = x$
- $\forall x \exists y. op(x, y) = \epsilon$

# Precedence and associativity

- $\neg$ ,  $\forall y$  and  $\exists y$  bind most tightly;
- then  $\vee$  and  $\wedge$ ;
- then  $\rightarrow$ , which is right-associative.

Syntax Tree for  $\forall x(\neg \exists y p(x, y) \vee \neg \exists y p(y, x))$ 

# Free and Bound variables

## Definition

Let  $\phi$  be a formula in predicate logic.

- occurrence of  $x$  in  $\phi$  is **free** (in  $\phi$ ) if
  - it is a leaf node in the parse tree of  $\phi$ , such that
  - there is no path upwards from that node  $x$  to a node  $\forall x$  or  $\exists x$ .
- Otherwise, that occurrence of  $x$  is called bound.

# Free and Bound variables

## Definition

Let  $\phi$  be a formula in predicate logic.

- occurrence of  $x$  in  $\phi$  is **free** (in  $\phi$ ) if
    - it is a leaf node in the parse tree of  $\phi$ , such that
    - there is no path upwards from that node  $x$  to a node  $\forall x$  or  $\exists x$ .
  - Otherwise, that occurrence of  $x$  is called bound.
- 
- For  $\forall x \phi$ , or  $\exists x \phi$ , we say that  $\phi$  — minus any of  $\psi$ 's subformulas  $\exists x \psi$ , or  $\forall x \psi$  — is the **scope** of  $\forall x$ , respectively  $\exists x$ .

# Free and Bound variables

## Definition

Let  $\phi$  be a formula in predicate logic.

- occurrence of  $x$  in  $\phi$  is **free** (in  $\phi$ ) if
    - it is a leaf node in the parse tree of  $\phi$ , such that
    - there is no path upwards from that node  $x$  to a node  $\forall x$  or  $\exists x$ .
  - Otherwise, that occurrence of  $x$  is called bound.
- 
- For  $\forall x \phi$ , or  $\exists x \phi$ , we say that  $\phi$  — minus any of  $\psi$ 's subformulas  $\exists x \psi$ , or  $\forall x \psi$  — is the **scope** of  $\forall x$ , respectively  $\exists x$ .
  - A formula that has no free variables is referred to as a **closed** formula.



# Free and Bound variables

## Definition

Let  $\phi$  be a formula in predicate logic.

- occurrence of  $x$  in  $\phi$  is **free** (in  $\phi$ ) if
    - it is a leaf node in the parse tree of  $\phi$ , such that
    - there is no path upwards from that node  $x$  to a node  $\forall x$  or  $\exists x$ .
  - Otherwise, that occurrence of  $x$  is called bound.
- 
- For  $\forall x \phi$ , or  $\exists x \phi$ , we say that  $\phi$  — minus any of  $\psi$ 's subformulas  $\exists x \psi$ , or  $\forall x \psi$  — is the **scope** of  $\forall x$ , respectively  $\exists x$ .
  - A formula that has no free variables is referred to as a **closed** formula.

$$(\forall x(P(x) \wedge Q(x))) \rightarrow (\neg P(x) \vee Q(y))$$

# Substitution

## Substitution

Given a variable  $x$ , a term  $t$  and a formula  $\psi$  we define  $\psi[t/x]$  to be the formula obtained by replacing each *free* occurrence of variable  $x$  in  $\psi$  with  $t$ .


# Substitution

## Substitution

Given a variable  $x$ , a term  $t$  and a formula  $\psi$  we define  $\psi[t/x]$  to be the formula obtained by replacing each *free* occurrence of variable  $x$  in  $\psi$  with  $t$ .

- Incorrect failure can lead to (binding) capture.

# Substitution



the term  $f(y, y)$  is  
not free for  $x$  in  
this formula

# Natural Deduction

## Equality

$$\frac{}{t = t} = i \qquad \frac{t_1 = t_2 \quad \phi[t_1/x]}{\phi[t_2/x]} = e$$

# Natural Deduction

## Equality

$$\frac{}{t = t} = i \qquad \frac{t_1 = t_2 \quad \phi[t_1/x]}{\phi[t_2/x]} = e$$

Here equality does not mean syntactic, or intensional, equality, but equality in terms of computation results

# Natural Deduction

## Universal Quantification

$$\frac{\forall x. \phi}{\phi[t/x]} \forall xe \qquad \frac{\boxed{\begin{array}{c} x_0 \text{ fresh} \\ \dots \\ \phi[x_0/x] \end{array}}}{\forall x. \phi} \forall i$$

# Natural Deduction

## Universal Quantification

$$\frac{\forall x. \phi}{\phi[t/x]} \forall xe \quad \frac{\boxed{\begin{array}{c} x_0 \text{ fresh} \\ \dots \\ \phi[x_0/x] \end{array}}}{\forall x. \phi} \forall i$$

Note the *scope* (shown by the box): the scope contains *local assumptions* and fresh variables, and hence the derived formulas and fresh variables are not valid/occur outside the proof



# Natural Deduction

## Existential Quantification

$$\frac{\phi[t/x]}{\exists x \phi} \exists xi \quad \frac{\exists x. \phi \quad \boxed{\begin{array}{c} x_0 \text{ fresh, } \phi[x_0/x] \\ \dots \\ \gamma \end{array}}}{\gamma} \exists e$$

# Natural Deduction

## Existential Quantification

$$\begin{array}{c}
 \frac{\phi[t/x]}{\exists x \phi} \exists xi \quad \frac{\begin{array}{c} \exists x. \phi \\ \boxed{\begin{array}{c} x_0 \text{ fresh, } \phi[x_0/x] \\ \dots \\ \gamma \end{array}} \\ \gamma \end{array}}{\exists e}
 \end{array}$$

Note the *scope* (shown by the box): the scope contains *local assumptions* and fresh variables, and hence the derived formulas and fresh variables are not valid/occur outside the proof

# Natural Deduction

- Proof system
  - Easy to prove something is valid (show a proof for it)
  - Difficult to prove that something is not valid (maybe you are just not able to construct a proof)
  - For satisfiability, easy to prove contradictions (give proof of  $\neg\phi$ )
- Semantics
  - Easy to prove something is not valid (show a falsifiable interpretation)
  - Difficult to prove that something is valid (need to show all interpretations are satisfiable)
  - For satisfiability, easy to prove satisfiability (show the satisfiable interpretations)

# Natural Deduction

- Proof system
  - Easy to prove something is valid (show a proof for it)
  - Difficult to prove that something is not valid (maybe you are just not able to construct a proof)
  - For satisfiability, easy to prove contradictions (give proof of  $\neg\phi$ )
- Semantics
  - Easy to prove something is not valid (show a falsifiable interpretation)
  - Difficult to prove that something is valid (need to show all interpretations are satisfiable)
  - For satisfiability, easy to prove satisfiability (show the satisfiable interpretations)

So, both are important!

# Semantics

## Interpretation

Given a set of formulas  $U$  with a set of relations  $R$ , set of functions  $F$  and a set of constants  $C$ , an interpretation  $\mathcal{I}$  is a four tuple

$(D, r_1^{\mathcal{I}}, r_2^{\mathcal{I}}, \dots, r_n^{\mathcal{I}} \in R^{\mathcal{I}}, f_1^{\mathcal{I}}, f_2^{\mathcal{I}}, \dots, f_m^{\mathcal{I}} \in F^{\mathcal{I}}, c_1^{\mathcal{I}}, c_2^{\mathcal{I}}, \dots, c_l^{\mathcal{I}} \in C^{\mathcal{I}})$ , where

- $D$  is the domain of discourse
- For each relation symbol,  $r_i \in R$ , there exists a concrete relation  $r_i^{\mathcal{I}} \in R^{\mathcal{I}}$  (of same arity)
- For each function symbol,  $f_i \in F$ , there exists a concrete function  $f_i^{\mathcal{I}} \in F^{\mathcal{I}}$  (of same arity)
- For each function symbol,  $c_i \in C$ , there exists a concrete constant  $c_i^{\mathcal{I}} \in C^{\mathcal{I}}$

# Valuations

- Valuations/satisfact of a first-order formula on an interpretation  $\mathcal{I}$  only makes sense for a closed formula:  $\mathcal{I} \models \phi$

# Valuations

- Valuations/satisfact of a first-order formula on an interpretation  $\mathcal{I}$  only makes sense for a closed formula:  $\mathcal{I} \models \phi$
- For open formula, we also need to consider an *environment*  $\Gamma$ , where  $\Gamma : x \mapsto v$  (mapping of free variables to values in the universe of discourse)
  - $\mathcal{I} \models_I \phi, I \in \Gamma$  can now be questioned.

# Valuations

- Valuations/satisfact of a first-order formula on an interpretation  $\mathcal{I}$  only makes sense for a closed formula:  $\mathcal{I} \models \phi$
- For open formula, we also need to consider an *environment*  $\Gamma$ , where  $\Gamma : x \mapsto v$  (mapping of free variables to values in the universe of discourse)
  - $\mathcal{I} \models_I \phi$ ,  $I \in \Gamma$  can now be questioned.
- For simplicity, we will only restrict ourselves to closed formulas



## Example: State Transition System

Given  $\mathcal{S} =$

$(\{a, b, c\}, \{\{(a, a), (a, b), (a, c), (b, c), (c, c)\}, \{b, c\}\}, \{(a, b), (b, c), (c, a)\}, \{$

# Example: State Transition System

Given  $\mathcal{J} =$

$(\{a, b, c\}, \{\{(a, a), (a, b), (a, c), (b, c), (c, c)\}, \{b, c\}\}, \{(a, b), (b, c), (c, a)\}, \{$

- $D = \{a, b, c\}$  (states)
- There are two relations,  $R^{\mathcal{J}} = \{r_1^{\mathcal{J}}, r_2^{\mathcal{J}}\}$ :
  - $r_1^{\mathcal{J}} = \{(a, a), (a, b), (a, c), (b, c), (c, c)\}$  (transition relation)
  - $r_2^{\mathcal{J}} = \{b, c\}$  (a unary relation, i.e. a set) (final states)
- There is one function,  $F^{\mathcal{J}} = \{f_1^{\mathcal{J}}\}$  (additional “failure” function)
  - $f_1^{\mathcal{J}}(a) = b; f_1^{\mathcal{J}}(b) = c; f_1^{\mathcal{J}}(c) = a;$
- There are two constants:  $C^{\mathcal{J}} = \{a, b\}$  (initial states)

Check on  $\mathcal{I}$ 

- $\exists y. R(i, y)$
- $\neg F(i)$
- $\forall x, y, z (R(x, y) \rightarrow R(x, z) \rightarrow y = z)$
- $\forall x \exists y. R(x, y)$

# Ground Terms

- A ground term is a term which does not contain any variables.
- A ground atomic formula is an atomic formula, all of whose terms are ground.
- A ground literal is a ground atomic formula or the negation of one.
- A ground formula is a quantifier-free formula, all of whose atomic formula are ground.
- $A$  is a ground instance of a quantifier free formula  $A'$  iff it can be obtained from  $A'$  by substituting ground terms for the (free) variables in  $A'$ .

Theorem: The set of ground terms is countable.

# Proving Semantic Consequence

Harder than propositional logic (building truth tables)—needs argument on sets (1-ary relations), relations and functions

# Proving Semantic Consequence

Harder than propositional logic (building truth tables)—needs argument on sets (1-ary relations), relations and functions

$$\forall x(P(x) \rightarrow Q(x)) \models \forall xP(x) \rightarrow \forall xQ(x)$$

- Let  $\mathcal{M}$  be a model of  $\forall x(P(x) \rightarrow Q(x))$ ; need to show it is a model of  $\forall xP(x) \rightarrow \forall xQ(x)$
- Case I: all  $x \in P$ 
  - then, all  $x \in Q$  (from  $\forall x(P(x) \rightarrow Q(x))$ )
- Case II: let some  $x_0 \notin P$ 
  - then,  $\forall xP(x)$  is false, so  $\mathcal{M}$  satisfies.

# Proving Semantic Consequence

- $\forall x P(x) \rightarrow \forall x Q(x) \models \forall x (P(x) \rightarrow Q(x))$ 
  - Let  $\mathcal{M}$  be a model of  $\forall x P(x) \rightarrow \forall x Q(x)$ ; need to show it is a model of  $\forall x (P(x) \rightarrow Q(x))$
  - Case I: if all  $x_0 \in P$ , then for all  $x_0 \in Q$ . Consequent holds.
  - Case II: if there is some  $x \notin P$ , the premise is vacuously true asserting no constraints, allowing sets  $P$  and  $Q$  to be arbitrarily. Let us create counterexample:  $(\{a, b\}, \{\{a\}, \{b\}\}, \{\}, \{a, b\})$ .

# Soundness and Completeness

Natural Deduction is sound and complete with respect to first order semantics as described above.



# Compactness

## Compactness Theorem

Let  $\Gamma$  be a set of sentences in predicate logic. If all finite subsets of  $\Gamma$  are satisfiable, then so is  $\Gamma$ .

## Proof

- proof by contradiction
- let all finite subsets of  $\Gamma$  are satisfiable and  $\Gamma$  is not satisfiable
- then,  $\Gamma \models \perp$  ( $\Gamma$  can have infinite premises)
- by completeness,  $\Gamma \vdash \perp$
- so there exists a proof for  $\Gamma \vdash \perp$
- a proof implies that it can use only a finite premises, say some  $\Delta$  ( $\Delta$  is finite)
- so,  $\Delta \vdash \perp$
- by soundness,  $\Delta \models \perp$

# Application of Compactness

## Reachability

Reachability is not expressible in predicate logic

## Proof

- Let there be such a first order formula  $\psi$
- $\Phi_0 \stackrel{\text{def}}{=} c = c'; \phi_1 = R(c, c');$   
 $\Phi_n \stackrel{\text{def}}{=} \exists x_1, x_2, \dots, x_{n-1} (R(c, x_1) \wedge R(x_1, x_2) \wedge \dots \wedge R(x_{n-1}, c'))$
- here, interpretations are graphs
- $\Phi_n$ : there exists a path of length  $n$
- Let  $\Delta = \{\neg\Phi_i \mid i \geq 0\} \cup \{\psi[c/u][c'/v]\}$
- $\Delta$ : there does not exist a path of length 1, 2, ... but a finite path from  $c$  to  $c'$
- But, every finite subset is satisfiable
- $\Delta$  is unsatisfiable

# Application of Compactness

## Reachability

Reachability is not expressible in predicate logic

## Proof

- Let there be such a first order formula  $\psi$
- $\Phi_0 \stackrel{\text{def}}{=} c = c'; \phi_1 = R(c, c');$   
 $\Phi_n \stackrel{\text{def}}{=} \exists x_1, x_2, \dots, x_{n-1} (R(c, x_1) \wedge R(x_1, x_2) \wedge \dots \wedge R(x_{n-1}, c'))$
- here, interpretations are graphs
- $\Phi_n$ : there exists a path of length  $n$
- Let  $\Delta = \{\neg\Phi_i \mid i \geq 0\} \cup \{\psi[c/u][c'/v]\}$
- $\Delta$ : there does not exist a path of length 1, 2, ... but a finite path from  $c$  to  $c'$
- But, every finite subset is satisfiable
- $\Delta$  is unsatisfiable

# Normal Forms

## PCNF (prenex conjunctive normal form)

$$Q_1x_1 Q_2x_2 \dots Q_nx_n M$$

- $Q_1x_1 Q_2x_2 \dots Q_nx_n$  is prefix
- $M$  is the matrix

## Clausal form

$$Q_1x_1 Q_2x_2 \dots Q_nx_n M$$

- If all  $Q_i$  in prefix is universal quantification,
- $M$  is written as CNF

then formula can be written as a list of clauses.

# Skolem's Theorem

## Skolem's Theorem

For a closed formula  $A$ , there exists a clausal form such that  $A \approx A'$  ( $\approx$  is equisat)

# Skolem's Theorem

## Skolem's Theorem

For a closed formula  $A$ , there exists a clausal form such that  $A \approx A'$  ( $\approx$  is equisat)

## Skolemization

For a closed formula  $\exists x A(x, y)$ ,  $\exists x A = A[f(y)/x]$

- $\forall x \exists y p(x, y)$ : for all  $x$ , produce a  $y$  such that  $p(x, y)$  holds
- $f(y)$ : produces one such value of  $x$  (for a given  $y$ ) for which  $p(x, y)$  holds
- so, equisat (not all interpretations are retained) but not equivalent

# Skolemization Algorithm

$$\forall x(p(x) \rightarrow q(x)) \rightarrow (\forall x p(x) \rightarrow \forall x q(x))$$

- Rename bound variables

$$\forall x(p(x) \rightarrow q(x)) \rightarrow (\forall y p(y) \rightarrow \forall z q(z))$$

- Transform to only  $\vee$  and  $\wedge$

$$\neg \forall x(\neg p(x) \vee q(x)) \vee (\neg \forall y p(y) \vee \forall z q(z))$$

- Push negations inside

$$\exists x(p(x) \wedge \neg q(x)) \vee \exists y \neg p(y) \vee \forall z q(z)$$

- Extract quantifiers from matrix (out to in)

$$\exists y \exists x \forall z (p(x) \wedge \neg q(x)) \vee \neg p(y) \vee q(z)$$

- Skolemization (add functions with arguments for universal quantifier outside it)

- $\exists y \exists x \forall z (p(x) \wedge \neg q(x)) \vee \neg p(y) \vee q(z)$

- No, universal quantifier outside existentials:

- $\forall z (p(a) \wedge \neg q(a)) \vee \neg p(b) \vee q(z)$

- $\forall z \exists y \exists x (p(x) \wedge \neg q(x)) \vee \neg p(y) \vee q(z)$

- universal quantifier outside existentials is z:

- $\forall z (p(f(z)) \wedge \neg q(f(z))) \vee \neg p(g(z)) \vee q(z)$

# Herbrand Models

- canonical interpretations for set of models
- if a set of clauses has a model, it has a Herbrand model.



# Herbrand Models

- canonical interpretations for set of models
- if a set of clauses has a model, it has a Herbrand model.

## Herbrand Universe

- $a_i \in C$ , then  $a_i \in H_s$
- $f_i \in F, t_j \in H_s$ , then  $f_i(t_1, t_2, \dots, t_n) \in H_s$

# Herbrand Models

- canonical interpretations for set of models
- if a set of clauses has a model, it has a Herbrand model.

## Herbrand Universe

- $a_i \in C$ , then  $a_i \in H_s$
- $f_i \in F, t_j \in H_s$ , then  $f_i(t_1, t_2, \dots, t_n) \in H_s$

## Herbrand Base

set of ground atomic formulae that can be formed from predicate symbols in  $S$  and terms in  $H_s$

A relation over Herbrand universe is simply a subset of Herbrand base.

# Herbrand's Theorem

A set of clauses  $S$  is unsatisfiable if and only if a finite set of ground instances of clauses of  $S$  is unsatisfiable.

# Herbrand's Theorem

A set of clauses  $S$  is unsatisfiable if and only if a finite set of ground instances of clauses of  $S$  is unsatisfiable.

Gives a semi-decision procedure to solve first order satisfiability/validity.