


Catalan number

From Wikipedia, the free encyclopedia

For names of numbers in Catalan, see List of numbers in various languages#Occitano-Romance.

In combinatorial mathematics, the **Catalan numbers** form a sequence of natural numbers that occur in various counting problems, often involving recursively-defined objects. They are named after the Belgian mathematician Eugène Charles Catalan (1814–1894).

Using zero-based numbering, the *n*th Catalan number is given directly in terms of binomial coefficients by



The $C_5 = 42$ noncrossing partitions of a 5-element set (below, the other 10 of the 52 partitions)

$$C_n = \frac{1}{n+1} \binom{2n}{n} = \frac{(2n)!}{(n+1)!n!} = \prod_{k=2}^n \frac{n+k}{k} \quad \text{for } n \geq 0.$$

The first Catalan numbers for $n = 0, 1, 2, 3, \dots$ are

1, 1, 2, 5, 14, 42, 132, 429, 1430, 4862, 16796, 58786, 208012, 742900, 2674440, 9694845, 35357670, 129644790, 477638700, 1767263190, 6564120420, 24466267020, 91482563640, 343059613650, 1289904147324, 4861946401452, ... (sequence A000108 in OEIS).

Contents

- 1 Properties
- 2 Applications in combinatorics

- 3 Proof of the formula
 - 3.1 First proof
 - 3.2 Second proof
 - 3.3 Third proof
 - 3.4 Fourth proof
 - 3.5 Fifth proof
- 4 Hankel matrix
- 5 History
- 6 Generalizations
- 7 See also
- 8 Notes
- 9 References
- 10 External links

Properties

An alternative expression for C_n is

$$C_n = \binom{2n}{n} - \binom{2n}{n+1} = \frac{1}{n+1} \binom{2n}{n} \quad \text{for } n \geq 0,$$

which is equivalent to the expression given above because $\binom{2n}{n+1} = \frac{n}{n+1} \binom{2n}{n}$. This shows that C_n is an integer, which is not immediately obvious from the first formula given. This expression forms the basis for a proof of the correctness of the formula.

The Catalan numbers satisfy the recurrence relation

$$C_0 = 1 \quad \text{and} \quad C_{n+1} = \sum_{i=0}^n C_i C_{n-i} \quad \text{for } n \geq 0;$$

moreover,

$$C_n = \frac{1}{n+1} \sum_{i=0}^n \binom{n}{i}^2.$$

This is because $\binom{2n}{n} = \sum_{i=0}^n \binom{n}{i}^2$, since choosing n numbers from a $2n$ set of numbers can be uniquely divided into 2 parts: choosing i numbers out of the first n numbers and then choosing $n-i$ numbers from the remaining n numbers.

They also satisfy:

$$C_0 = 1 \quad \text{and} \quad C_{n+1} = \frac{2(2n+1)}{n+2} C_n,$$

which can be a more efficient way to calculate them.

Asymptotically, the Catalan numbers grow as

$$C_n \sim \frac{4^n}{n^{3/2}\sqrt{\pi}}$$

in the sense that the quotient of the n th Catalan number and the expression on the right tends towards 1 as $n \rightarrow +\infty$. Some sources use just $C_n \sim \frac{4^n}{n^{3/2}}$.^[1] (This can be proved by using Stirling's approximation for $n!$.)

The only Catalan numbers C_n that are odd are those for which $n = 2^k - 1$. All others are even.

The only prime Catalan numbers are $C_2 = 2$ and $C_3 = 5$.

The Catalan numbers have an integral representation

$$C_n = \int_0^4 x^n \rho(x) dx$$

where $\rho(x) = \frac{1}{2\pi} \sqrt{\frac{4-x}{x}}$. This means that the Catalan numbers are a solution of the Hausdorff moment problem on the interval $[0, 4]$ instead of $[0, 1]$. The orthogonal polynomials having the weight function $\rho(x)$ on $[0, 4]$ are

$$H_n(x) = \sum_{k=0}^n \binom{n+k}{n-k} (-x)^k.$$

Applications in combinatorics

There are many counting problems in combinatorics whose solution is given by the Catalan numbers. The book *Enumerative Combinatorics: Volume 2* by combinatorialist Richard P. Stanley contains a set of exercises which describe 66 different interpretations of the Catalan numbers. Following are some examples, with illustrations of the cases $C_3 = 5$ and $C_4 = 14$.

- C_n is the number of **Dyck words**^[2] of length $2n$. A Dyck word is a string consisting of n X's and n Y's such that no initial segment of the string has more Y's than X's (see also Dyck language). For example, the following are the Dyck words of length 6:

XXXYYY XYXXYY XYXYXY XXYYXY XXYYXY.


- Re-interpreting the symbol X as an open parenthesis and Y as a close parenthesis, C_n counts the number of expressions containing n pairs of parentheses which are correctly matched:

((())) ()(()) ()()() (())() (())()


- C_n is the number of different ways $n + 1$ factors can be completely parenthesized (or the number of ways of associating n applications of a binary operator). For $n = 3$, for example, we have the following five different parenthesizations of four factors:

((ab)c)d (a(bc))d (ab)(cd) a((bc)d) a(b(cd))


- Successive applications of a binary operator can be represented in terms of a full binary tree. (A rooted binary tree is *full* if every vertex has either two children or no children.) It follows that C_n is the number of full binary trees with $n + 1$ leaves:



Lattice of the 14 Dyck words of length 8 - (and) interpreted as *up* and *down*



The associahedron of order 4 with the $C_4=14$ full binary trees with 5 leaves




- C_n is the number of non-isomorphic ordered trees with n vertices. (An ordered tree is a rooted tree

in which the children of each vertex are given a fixed left-to-right order.)^[3]

- C_n is the number of monotonic lattice paths along the edges of a grid with $n \times n$ square cells, which do not pass above the diagonal. A monotonic path is one which starts in the lower left corner, finishes in the upper right corner, and consists entirely of edges pointing rightwards or upwards. Counting such paths is equivalent to counting Dyck words: X stands for "move right" and Y stands for "move up".


The following diagrams show the case $n = 4$:




This can be succinctly represented by listing the Catalan elements by column height:^[4]

[0,0,0,0][0,0,0,1][0,0,0,2][0,0,1,1]
 [0,1,1,1] [0,0,1,2] [0,0,0,3] [0,1,1,2][0,0,2,2][0,0,1,3]
 [0,0,2,3][0,1,1,3] [0,1,2,2][0,1,2,3]

- C_n is the number of different ways a convex polygon with $n + 2$ sides can be cut into triangles by connecting vertices with straight lines (a form of Polygon triangulation). The following hexagons illustrate the case $n = 4$:




- C_n is the number of stack-sortable permutations of $\{1, \dots, n\}$. A permutation w is called stack-sortable if $S(w) = (1, \dots, n)$, where $S(w)$ is defined recursively as follows: write $w = unv$ where n is the largest element in w and u and v are shorter sequences, and set $S(w) = S(u)S(v)n$, with S being the identity for one-element sequences. These are the permutations that avoid the pattern 231.




The triangles correspond to nodes of the binary trees.

- 9/23/2015
- Catalan number - Wikipedia, the free encyclopedia
- C_n is the number of permutations of $\{1, \dots, n\}$ that avoid the pattern 123 (or any of the other patterns of length 3); that is, the number of permutations with no three-term increasing subsequence. For $n = 3$, these permutations are 132, 213, 231, 312 and 321. For $n = 4$, they are 1432, 2143, 2413, 2431, 3142, 3214, 3241, 3412, 3421, 4132, 4213, 4231, 4312 and 4321.
 - C_n is the number of noncrossing partitions of the set $\{1, \dots, n\}$. *A fortiori*, C_n never exceeds the n th Bell number. C_n is also the number of noncrossing partitions of the set $\{1, \dots, 2n\}$ in which every block is of size 2. The conjunction of these two facts may be used in a proof by mathematical induction that all of the *free* cumulants of degree more than 2 of the Wigner semicircle law are zero. This law is important in free probability theory and the theory of random matrices.
 - C_n is the number of ways to tile a stairstep shape of height n with n rectangles. The following figure illustrates the case $n = 4$:



- C_n is the number of rooted binary trees with n internal nodes ($n + 1$ leaves or external nodes). Illustrated in following Figure are the trees corresponding to $n = 0,1,2$ and 3. There are 1, 1, 2, and 5 respectively. Here, we consider as binary trees those in which each node has zero or two children, and the internal nodes are those that have children.



- C_n is the number of ways to form a “mountain ranges” with n upstrokes and n down-strokes that all stay above the original line. The mountain range interpretation is that the mountains will never go below the horizon.

$n = 0$:	*	1 way
$n = 1$:	/\	1 way
$n = 2$:	<div> <div>/\</div> <div>/\</div> </div>	2 ways
$n = 3$:	<div> <div> <div>/\</div> <div>/\</div> </div> <div> <div>/\</div> <div>/\</div> </div> <div> <div>/\</div> <div>/\</div> </div> <div> <div>/\</div> <div>/\</div> </div> </div>	5 ways

Mountain Ranges

- C_n is the number of standard Young tableaux whose diagram is a 2-by- n rectangle. In other words, it is the number of ways the numbers 1, 2, ..., $2n$ can be arranged in a 2-by- n rectangle so that each

row and each column is increasing. As such, the formula can be derived as a special case of the hook-length formula.

- C_n is the number of ways that the vertices of a convex $2n$ -gon can be paired so that the line segments joining paired vertices do not intersect. This is precisely the condition that guarantees that the paired edges can be identified (sewn together) to form a closed surface of genus zero (a topological 2-sphere).
- C_n is the number of semiorders on n unlabeled items.^[5]

Proof of the formula

There are several ways of explaining why the formula

$$C_n = \frac{1}{n+1} \binom{2n}{n}$$

solves the combinatorial problems listed above. The first proof below uses a generating function. The other proofs are examples of bijective proofs; they involve literally counting a collection of some kind of object to arrive at the correct formula.

First proof

We first observe that all of the combinatorial problems listed above satisfy Segner's^[6] recurrence relation

$$C_0 = 1 \quad \text{and} \quad C_{n+1} = \sum_{i=0}^n C_i C_{n-i} \quad \text{for } n \geq 0.$$

For example, every Dyck word w of length ≥ 2 can be written in a unique way in the form

$$w = Xw_1Yw_2$$

with (possibly empty) Dyck words w_1 and w_2 .

The generating function for the Catalan numbers is defined by

$$c(x) = \sum_{n=0}^{\infty} C_n x^n.$$

The two recurrence relations together can then be summarized in generating function form by the relation

$$c(x) = 1 + xc(x)^2;$$

in other words, this equation follows from the recurrence relations by expanding both sides into power series. On the one hand, the recurrence relations uniquely determine the Catalan numbers; on the other hand, the generating function solution

$$c(x) = \frac{1 - \sqrt{1 - 4x}}{2x} = \frac{2}{1 + \sqrt{1 - 4x}}$$

has a power series at 0 and its coefficients must therefore be the Catalan numbers. The chosen solution satisfies the following condition.

$$\lim_{x \rightarrow 0^+} c(x) = C_0 = 1$$

The other solution has a pole at 0 and this reasoning doesn't apply to it.

The square root term can be expanded as a power series using the identity

$$\sqrt{1 + y} = \sum_{n=0}^{\infty} \binom{\frac{1}{2}}{n} y^n = 1 - 2 \sum_{n=1}^{\infty} \binom{2n-2}{n-1} \left(\frac{-1}{4}\right)^n \frac{y^n}{n}.$$

This is a special case of Newton's generalized binomial theorem; as with the general theorem, it can be proved by computing derivatives to produce its Taylor series. Setting $y = -4x$ and substituting this power series into the expression for $c(x)$ and shifting the summation index n by 1, the expansion simplifies to

$$c(x) = \sum_{n=0}^{\infty} \binom{2n}{n} \frac{x^n}{n+1}.$$

The coefficients are now the desired formula for C_n .

Another way to get $c(x)$ is to solve for $xc(x)$ and observe that $\int_0^x t^n dt$ appears in each term of the power series.

Second proof

This proof depends on a trick known as André's reflection method, which was originally used in connection with Bertrand's ballot theorem. (The reflection principle has been widely attributed to Désiré André, but his method did not actually use reflections; and the reflection method is a variation due to Aebly and Mirimanoff.^[7]) We count the paths which start and end on the diagonal of the $n \times n$ grid. All such paths have n rightward and n upward steps. Since we can choose which of the $2n$ steps are upward (or, equivalently, rightward) ones, there are $\binom{2n}{n}$ total monotonic paths of this type. A *bad* path will cross the main diagonal and touch the next higher (*fatal*) diagonal (depicted red in the illustration). We flip the portion of the path occurring after that touch about that fatal diagonal, as illustrated; this geometric operation amounts to interchanging all the rightward and upward steps after that touch. In the section of the path that is not reflected, there is one more upward step than rightward steps, so the remaining section of the bad path has one more rightward than upward step (because it ends on the main diagonal). When this portion of the path is reflected, it will also have one more upward step than rightward steps. Since there are still $2n$ steps, there must now be $n + 1$ upward steps and $n - 1$ rightward steps. So, instead of reaching the target (n, n) , all bad paths (after the portion of the path is reflected) will end in location $(n - 1, n + 1)$. As any monotonic path




Figure 1. The invalid portion of the path is flipped. Bad paths reach $(n - 1, n + 1)$ instead of (n, n) .




Figure 3. The green and red portions are being exchanged.

Notice that the exceedance has dropped from three to two. In fact, the algorithm will cause the exceedance to decrease by one, for any path that we feed it, because the first vertical step starting on the diagonal (at the point marked with a black dot) is the unique vertical edge that under the operation passes from above the diagonal to below it; all other vertical edges stay on the same side of the diagonal.

It is also not difficult to see that this process is *reversible*: given any path P whose exceedance is less than n , there is exactly one path which yields P when the algorithm is applied to it. Indeed, the (black) edge X , which originally was the first horizontal step ending on the diagonal, has become the *last* horizontal step *starting* on the diagonal.

This implies that the number of paths of exceedance n is equal to the number of paths of exceedance $n - 1$, which is equal to the number of paths of exceedance $n - 2$, and so on, down to zero. In other words, we have split up the set of *all* monotonic paths into $n + 1$ equally sized classes, corresponding to the possible exceedances between 0 and n . Since there are

$$\binom{2n}{n}$$

monotonic paths, we obtain the desired formula

$$C_n = \frac{1}{n+1} \binom{2n}{n}.$$

Figure 4 illustrates the situation for $n = 3$. Each of the 20 possible monotonic paths appears somewhere in the table. The first column shows all paths of exceedance three, which lie entirely above the diagonal. The columns to the right show the result of successive applications of the algorithm, with the exceedance decreasing one unit at a time. There are five rows, that is, $C_3 = 5$.




Figure 4. All monotonic paths in a 3×3 grid, illustrating the exceedance-decreasing algorithm.

Fourth proof

This proof uses the triangulation definition of Catalan numbers to establish a relation between C_n and C_{n+1} . Given a polygon P with $n+2$ sides, first mark one of its sides as the base. If P is then triangulated, we can further choose and orient one of its $2n+1$ edges. There are $(4n+2)C_n$ such decorated triangulations. Now given a polygon Q with $n+3$ sides, again mark one of its sides as the base. If Q is triangulated, we can further mark one of the sides other than the base side. There are $(n+2)C_{n+1}$ such decorated triangulations. Then there is a simple bijection between these two kinds of decorated triangulations: We can either collapse the triangle in Q whose side is marked, or in reverse expand the oriented edge in P to a triangle and mark its new side. Thus

$$(4n+2)C_n = (n+2)C_{n+1}.$$

The binomial formula for C_n follows immediately from this relation and the initial condition $C_1 = 1$.

Fifth proof

This proof is based on the Dyck words interpretation of the Catalan numbers, so C_n is the number of ways to correctly match n pairs of brackets. We denote a (possibly empty) *correct* string with c and its inverse (where "[" and "]" are exchanged) with c^+ . Since any c can be uniquely decomposed into $c = [c_1]c_2$, summing over the possible spots to place the closing bracket immediately gives the recursive definition

$$C_0 = 1 \quad \text{and} \quad C_{n+1} = \sum_{i=0}^n C_i C_{n-i} \quad \text{for } n \geq 0.$$

Now let b stand for a *balanced* string of length $2n$ —that is, containing an equal number of "[" and "]"—and $B_n = \binom{2n}{n} = d_n C_n$ with some factor $d_n \geq 1$. As above, any balanced string can be uniquely decomposed into either $[c]b$ or $c^+[b]$, so

$$B_{n+1} = 2 \sum_{i=0}^n B_i C_{n-i}.$$

Also, any incorrect balanced string starts with $c]$, so

$$B_{n+1} - C_{n+1} = \sum_{i=0}^n \binom{2i+1}{i} C_{n-i} = \sum_{i=0}^n \frac{2i+1}{i+1} B_i C_{n-i}.$$

Subtracting the above equations and using $B_i = d_i C_i$ gives

$$C_{n+1} = 2 \sum_{i=0}^n d_i C_i C_{n-i} - \sum_{i=0}^n \frac{2i+1}{i+1} d_i C_i C_{n-i} = \sum_{i=0}^n \frac{d_i}{i+1} C_i C_{n-i}.$$

Comparing coefficients with the original recursion formula for C_n gives $d_i = i+1$, so

$$C_n = \frac{1}{n+1} \binom{2n}{n}.$$

Hankel matrix

The $n \times n$ Hankel matrix whose (i, j) entry is the Catalan number C_{i+j-2} has determinant 1, regardless of the value of n . For example, for $n = 4$ we have

$$\det \begin{bmatrix} 1 & 1 & 2 & 5 \\ 1 & 2 & 5 & 14 \\ 2 & 5 & 14 & 42 \\ 5 & 14 & 42 & 132 \end{bmatrix} = 1.$$

Moreover, if the indexing is "shifted" so that the (i, j) entry is filled with the Catalan number C_{i+j-1} then the determinant is still 1, regardless of the value of n . For example, for $n = 4$ we have

$$\det \begin{bmatrix} 1 & 2 & 5 & 14 \\ 2 & 5 & 14 & 42 \\ 5 & 14 & 42 & 132 \\ 14 & 42 & 132 & 429 \end{bmatrix} = 1.$$

Taken together, these two conditions uniquely define the Catalan numbers.


History

The Catalan sequence was described in the 18th century by Leonhard Euler, who was interested in the number of different ways of dividing a polygon into triangles. The sequence is named after Eugène Charles Catalan, who discovered the connection to parenthesized expressions during his exploration of the Towers of Hanoi puzzle. The counting trick for Dyck words was found by D. André in 1887.

In 1988, it came to light that the Catalan number sequence had been used in China by the Mongolian mathematician Mingantu by 1730.^{[9][10]} That is when he started to write his book *Ge Yuan Mi Lu Jie Fa*, which was completed by his student Chen Jixin in 1774 but published sixty years later. P.J. Larcombe (1999) sketched some of the features of the work of Mingantu, including the stimulus of Pierre Jartoux, who brought three infinite series to China early in the 1700s.

For instance, Ming used the Catalan sequence to express series expansions of $\sin(2\alpha)$ and $\sin(4\alpha)$ in terms of $\sin(\alpha)$.

Generalizations




The two-parameter sequence of non-negative integers $\frac{(2m)!(2n)!}{(m+n)!m!n!}$ is a generalization of the

Catalan numbers. These are named **super-Catalan numbers**, by Ira Gessel. These number should not confused with the Schröder–Hipparchus numbers, which sometimes are also called super-Catalan numbers.

For $m = 1$, this is just two times the ordinary Catalan numbers, and for $m = n$, the numbers have an easy combinatorial description. However, other combinatorial descriptions are only known^[11] for $m = 2$ and $m = 3$, and it is an open problem to find a general combinatorial interpretation.

See also

- Associahedron
- Bertrand's ballot theorem
- Binomial transform
-  Partition related number triangles
- Catalan's triangle
- Catalan–Mersenne number
- Fuss–Catalan number
- List of factorial and binomial topics
- Lobb numbers
- Narayana number
- Schröder–Hipparchus number
- Tamari lattice
- Wedderburn–Etherington number

Notes

1. Cormen, Thomas H.; Leiserson, Charles E.; Rivest, Ronald L. (1990). "Dynamic Programming". *Introduction to Algorithms*. Cambridge, Massachusetts: The MIT Press. p. 304. ISBN 0262031418.
2. Equivalent definitions of Dyck paths (<http://www.FindStat.org/DyckPaths>)
3. Stanley p.221 example (e)
4. ˇCrepinšek, Matej; Luka Mernik (2009). "AN EFFICIENT REPRESENTATION FOR SOLVING CATALAN NUMBER RELATED PROBLEMS" (<http://www.ijpam.eu/contents/2009-56-4/11/11.pdf>) (PDF). *International Journal of Pure and Applied Mathematics* **56** (4).
5. Kim, K. H.; Roush, F. W. (1978), "Enumeration of isomorphism classes of semiorders", *Journal of Combinatorics, Information & System Sciences* **3** (2): 58–61, MR 538212 (<https://www.ams.org/mathscinet-getitem?mr=538212>).
6. A. de Segner, Enumeratio modorum, quibus figurae planae rectilineae per diagonales dividuntur in triangua. *Novi commentarii academiae scientiarum Petropolitanae* **7** (1758/59) 203–209.
7. Renault, Marc, Lost (and found) in translation: André's actual method and its application to the generalized ballot problem. Amer. Math. Monthly 115 (2008), no. 4, 358--363. (<http://webSPACE.ship.edu/msrenault/ballotproblem/monthly358-363-renault.pdf>)
8. Rukavicka Josef (2011), *On Generalized Dyck Paths*, *Electronic Journal of Combinatorics online* (<http://www.combinatorics.org/ojs/index.php/eljc/article/view/v18i1p40/pdf>)
9. The 18th century Chinese discovery of the Catalan numbers (http://www.math.ucla.edu/~pak/lectures/Cat/Larcombe-The_18th_century_Chinese_discovery_of_the_Catalan_numbers.pdf)
10. Ming Antu, the First Inventor of Catalan Numbers in the World (http://en.cnki.com.cn/Article_en/CJFDTOTAL-NMGX198802004.htm)
11. Chen, Xin. "The super Catalan numbers S(m,m+s) for s≤3 and some integer factorial ratios" (<http://www-users.math.umn.edu/~reiner/REU/ChenWang2012.pdf>) (PDF). www-users.math.umn.edu/~reiner/REU/ChenWang2012.pdf. Retrieved 26 September 2014.

References

- Conway and Guy (1996) *The Book of Numbers*. New York: Copernicus, pp. 96–106.
- Gardner, Martin (1988), *Time Travel and Other Mathematical Bewilderments*, New York: W.H. Freeman and Company, pp. 253–266 (Ch. 20), ISBN 0-7167-1924-X
- Koshy, Thomas (2008), *Catalan Numbers with Applications* (http://www.amazon.com/Thomas-Koshy/e/B001H6NZT4/ref=ntt_athr_dp_pel_1), Oxford University Press, ISBN 0-19-533454-X
- Koshy, Thomas & Zhenguang Gao (2011) "Some divisibility properties of Catalan numbers", *Mathematical Gazette* 95:96–102.
- Larcombe, P.J. (1999) "The 18th century Chinese discovery of the Catalan numbers (http://www.math.ucla.edu/~pak/lectures/Cat/Larcombe-The_18th_century_Chinese_discovery_of_the_Catalan_numbers.pdf)", *Mathematical Spectrum* 32:5–7.
- Stanley, Richard P. (1999), *Enumerative combinatorics. Vol. 2* (<http://www-math.mit.edu/~rstan/ec/>), Cambridge Studies in Advanced Mathematics **62**, Cambridge University Press, ISBN 978-0-521-56069-6, MR 1676282 (<https://www.ams.org/mathscinet-getitem?mr=1676282>)
- Egecioglu, Omer (2009), *A Catalan-Hankel Determinant Evaluation* (http://www.cs.ucsb.edu/~omer/DOWNLOADABLE/catalan_hankel09.pdf) (PDF)

External links

- Stanley, Richard P. (1998), *Catalan addendum to Enumerative Combinatorics, Volume 2* (<http://www-math.mit.edu/~rstan/ec/catadd.pdf>) (PDF)
- Weisstein, Eric W., "Catalan Number" (<http://mathworld.wolfram.com/CatalanNumber.html>), *MathWorld*.
- Dickau, Robert M.: Catalan numbers (<http://mathforum.org/advanced/robertd/catalan.html>) Further examples.
- Davis, Tom: Catalan numbers (<http://mathcircle.berkeley.edu/BMC6/pdf0607/catalan.pdf>). Still more examples.
- Schmidhammer, Jürgen: Catalan-Zahlen (<http://www.bnv-bamberg.de/home/ba2636/catalanz.pdf>) Zulassungsarbeit zum Staatsexamen (PDF-File; 7,05 MB)
- "Equivalence of Three Catalan Number Interpretations" from The Wolfram Demonstrations Project [1] (<http://demonstrations.wolfram.com/EquivalenceOfThreeCatalanNumberInterpretations/>)
- Dyck paths (<http://www.findstat.org/DyckPaths>) and binary trees (<http://www.findstat.org/BinaryTrees>) in the FindStat (<http://www.findstat.org/>) database
- <https://www.youtube.com/watch?v=pEJo0DJhYvU> (Movie: generating all correct strings of parentheses with backtracking method in C language)

Retrieved from "https://en.wikipedia.org/w/index.php?title=Catalan_number&oldid=681350090"

Categories: Integer sequences | Factorial and binomial topics | Enumerative combinatorics

-
- This page was last modified on 16 September 2015, at 18:19.
 - Text is available under the Creative Commons Attribution-ShareAlike License; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.