Search Wikipedia

Edit | View history

Read

Q



WikipediA The Free Encyclopedia

Article Talk

Main page Contents Current events Random article About Wikipedia Contact us Donate

Contribute Help Learn to edit Community portal Recent changes

Upload file Tools What links here Related changes Special pages Permanent link

Cite this page Wikidata item Print/export Download as PDF

Page information

Printable version Languages

O

বাংলা Deutsch Español Français 한국어 Italiano Português Русский l agalog 文 11 more

Edit links

Multinomial distribution From Wikipedia, the free encyclopedia

In probability theory, the **multinomial distribution** is a generalization of the binomial distribution. For example, it models the probability of counts for each side of a k-sided die rolled n times. For n independent trials each of which leads to a success for exactly one of k categories, with each category having a given fixed success probability, the multinomial distribution gives the probability of any particular combination of numbers of successes for the various categories. When k is 2 and n is 1, the multinomial distribution is the Bernoulli distribution. When k is 2 and n is


bigger than 1, it is the binomial distribution. When k is bigger than 2 and n is 1, it is the categorical distribution. The Bernoulli distribution models the outcome of a single Bernoulli trial. In other words, it models whether flipping a (possibly biased) coin one time will result in either a success (obtaining a head)

or failure (obtaining a tail). The binomial distribution generalizes this to the number of heads from performing n independent flips (Bernoulli trials) of the same coin. The multinomial distribution models the outcome of n experiments, where the outcome of each trial has a categorical distribution, such as rolling a k-sided die n times. Let k be a fixed finite number. Mathematically, we have k possible mutually exclusive outcomes,

with corresponding probabilities  $p_1, ..., p_k$ , and n independent trials. Since the k outcomes are mutually exclusive and one must occur we have  $p_i \ge 0$  for i = 1, ..., k and  $\sum p_i = 1$ . Then if the

random variables  $X_i$  indicate the number of times outcome number i is observed over the n trials, the vector  $X = (X_1, ..., X_k)$  follows a multinomial distribution with parameters n and **p**, where

containing a 1 and all other elements containing a 0) rather than as an integer in the range 1...K; in this form, a categorical distribution is equivalent to a multinomial distribution over a single trial. Contents [hide] 1 Specification



 $f(x_1,\ldots,x_k;n,p_1,\ldots,p_k)=\Pr(X_1=x_1 \text{ and }\ldots \text{ and } X_k=x_k)$ 

Suppose one does an experiment of extracting n balls of k different colors from a bag, replacing the extracted balls after each draw. Balls of the same color are equivalent. Denote the

for non-negative integers 
$$x_1,...,x_k$$
. The probability mass function can be expressed using the gamma function as: 
$$f(x_1,\ldots,x_k;p_1,\ldots,p_k) = \frac{\Gamma(\sum_i x_i+1)}{\prod_i \Gamma(x_i+1)} \prod_{i=1}^k p_i^{x_i}.$$

Just like one can interpret the binomial distribution as (normalized) one-dimensional (1D) slices of Pascal's triangle, so too can one interpret the multinomial distribution as 2D (triangular)

As slices of generalized Pascal's triangle [edit]

As polynomial coefficients [edit] Similarly, just like one can interpret the binomial distribution as the polynomial coefficients of  $(p+(1-p))^n$  when expanded, one can interpret the multinomial distribution as the coefficients of  $(p_1+p_2+p_3+\cdots+p_k)^n$  when expanded. (Note that just like the binomial distribution, the coefficients must sum to 1.) This is the origin of the name "multinomial"

# $E(X_i) = np_i$ .

distribution".

The covariance matrix is as follows. Each diagonal entry is the variance of a binomially distributed random variable, and is therefore  $\operatorname{Var}(X_i) = np_i(1-p_i).$ 

 $Cov(X_i, X_i) = -np_ip_i$ 

for i, j distinct.

k = n and where the  $p_i$  are all equal, the covariance matrix is the centering matrix.

 $\rho(X_i, X_i) = 1.$ 

Note that the sample size drops out of this expression.

The entries of the corresponding correlation matrix are

Each of the 
$$k$$
 components separately has a binomial distribution with parameters  $n$  and  $p_i$ , for the appropriate value of the subscript  $i$ . The support of the multinomial distribution is the set  $\{(n_1,\ldots,n_k)\in\mathbb{N}^k\mid n_1+\cdots+n_k=n\}.$ 

Its number of elements is

 $E(\mathbf{X}) = n\mathbf{p},$ and

Example [edit] Suppose that in a three-way election for a large country, candidate A received 20% of the votes, candidate B received 30% of the votes, and candidate C received 50% of the votes. If six

large.

Example

Cells:

Αi

 $\Pr(A=1,B=2,C=3) = \frac{6!}{1!2!3!}(0.2^1)(0.3^2)(0.5^3) = 0.135$ 

First, reorder the parameters  $p_1, \ldots, p_k$  such that they are sorted in descending order (this is only to speed up computation and not strictly necessary). Now, for each trial, draw an auxiliary variable X from a uniform (0, 1) distribution. The resulting outcome is the component

If we have: 2 3 Categories

**Probabilities** 

Then, with software such as Excel, we may use the following recipe:

Sampling from a multinomial distribution [edit]

multinomial distribution with n equal to the number of such repetitions.

After that, we will use functions such as SumIf to accumulate the observed results by category and to calculate the estimated covariance matrix for each simulated sample. Ar ln dis

Equivalence tests for multinomial distributions [edit] The goal of equivalence testing is to establish the agreement between a theoretical multinomial distribution and observed counting frequencies. The theoretical distribution may be a fully

significance level. The equivalence test for Euclidean distance can be found in text book of Wellek (2010).[2] The equivalence test for the total variation distance is developed in Ostrovski (2017).[3] The exact equivalence test for the specific cumulative distance is proposed in Frey (2009).[4] The distance between the true underlying distribution p and a family of the multinomial distributions  $\mathcal{M}$  is defined by  $d(p,\mathcal{M}) = \min_{h \in \mathcal{M}} d(p,h)$ . Then the equivalence test problem is given by  $H_0=\{d(p,\mathcal{M})\geq arepsilon\}$  and  $H_1=\{d(p,\mathcal{M})<arepsilon\}$ . The distance  $d(p,\mathcal{M})$  is usually computed using numerical optimization. The tests for this case are developed recently

 Negative multinomial distribution • Hardy–Weinberg principle (it is a trinomial distribution with probabilities  $(\theta^2, 2\theta(1-\theta), (1-\theta)^2)$ )

3. A Ostrovski, Vladimir (May 2017). "Testing equivalence of multinomial distributions". Statistics & Probability Letters. 124: 77–82. doi:10.1016/j.spl.2017.01.004 . S2CID 126293429 . Official

### Citations [edit] 1. ^ "1.7 - The Multinomial Distribution | STAT 504" ☑. onlinecourses.science.psu.edu. Retrieved 2016-09-11. Wellek, Stefan (2010). Testing statistical hypotheses of equivalence and noninferiority. Chapman and Hall/CRC. ISBN 978-1439808184.

This page was last edited on 4 May 2021, at 18:35 (UTC).

References [edit]

V\*T\*E

Related distributions [edit]

in Ostrovski (2018).[5]

- web link (subscription required) @. Alternate, free web link @. 4. ^ Frey, Jesse (March 2009). "An exact multinomial test for equivalence". The Canadian Journal of Statistics. 37: 47–59. doi:10.1002/cjs.10000 ₺.Official web link (subscription required) ₺. Ostrovski, Vladimir (March 2018). "Testing equivalence to families of multinomial distributions with application to the independence model". Statistics & Probability Letters. 139: 61–66. doi:10.1016/j.spl.2018.03.014 년. S2CID 126261081 년.Official web link (subscription required) 년. Alternate, free web link 년.
- Sources [edit] • Evans, Morton; Hastings, Nicholas; Peacock, Brian (2000). Statistical Distributions (3rd ed.). New York: Wiley. pp. 134 4-136. ISBN 0-471-37124-6.
  - Discrete univariate Benford · Bernoulli · beta-binomial · binomial · categorical · hypergeometric · Poisson binomial · Rademacher · soliton · discrete uniform · Zipf · Zipf-Mandelbrot with finite support Discrete univariate beta negative binomial · Borel · Conway-Maxwell-Poisson · discrete phase-type · Delaporte · extended negative binomial · Flory-Schulz · Gauss-Kuzmin · geometric · with infinite support logarithmic · negative binomial · Panjer · parabolic fractal · Poisson · Skellam · Yule-Simon · zeta Continuous univariate arcsine · ARGUS · Balding-Nichols · Bates · beta · beta rectangular · continuous Bernoulli · Irwin-Hall · Kumaraswamy · logit-normal · noncentral beta · PERT ·

Probability distributions (List)

Cauchy · exponential power · Fisher's z · Gaussian q · generalized normal · generalized hyperbolic · geometric stable · Gumbel · Holtsmark · hyperbolic secant ·

- Continuous univariate hyper-Erlang · hyperexponential · hypoexponential · inverse chi-squared (scaled inverse chi-squared) · inverse Gaussian · inverse gamma · Kolmogorov · Lévy · supported on a semi-infinite interval log-Cauchy · log-Laplace · log-logistic · log-normal · Lomax · matrix-exponential · Maxwell-Boltzmann · Maxwell-Jüttner · Mittag-Leffler · Nakagami ·
- noncentral chi-squared · noncentral F · Pareto · phase-type · poly-Weibull · Rayleigh · relativistic Breit-Wigner · Rice · shifted Gompertz · truncated normal · type-2 Gumbel · Weibull (discrete Weibull) · Wilks's lambda
- Mixed continuous-discrete univariate rectified Gaussian Multivariate (joint) matrix gamma · normal-inverse-Wishart · normal-Wishart · Wishart
- Categories: Discrete distributions | Multivariate discrete distributions | Factorial and binomial topics | Exponential family distributions

Text is available under the Creative Commons Attribution-ShareAlike License; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.

Privacy policy About Wikipedia Disclaimers Contact Wikipedia Mobile view Developers Statistics Cookie statement


WIKIMEDIA

MediaWiki

[hide]

Multinomial  $|p_1,\ldots,p_k|$  event probabilities ( $\Sigma p_i=1$ )  $ig|x_i \in \{0,\ldots,n\}, \;\; i \in \{1,\ldots,k\}, ext{ with } \sum x_i = n$  $\frac{n!}{x_1!\cdots x_k!}p_1^{x_1}\cdots p_k^{x_k}$  $\mathrm{E}(X_i) = np_i$  $\mathrm{Var}(X_i) = np_i(1-p_i)$  $\mathrm{Cov}(X_i,X_j) = -np_ip_j \ \ (i 
eq j)$  $-\log(n!) - n \sum_{i=1}^k p_i \log(p_i) + \sum_{i=1}^k \sum_{x_i=0}^n \binom{n}{x_i} p_i^{x_i} (1-p_i)^{n-x_i} \log(x_i!)$  $\Big(\sum^{\kappa}p_{i}e^{t_{i}}\Big)$ 

**Parameters** n>0 number of trials (integer) Support **PMF** Mean Variance Entropy MGF CF **PGF**  $\left(\sum_{i=1}^{k}p_{i}z_{i}
ight)^{n} ext{ for }(z_{1},\ldots,z_{k})\in\mathbb{C}^{k}$ 



# 1.1 Probability mass function 2.1 As slices of generalized Pascal's triangle

# variable which is the number of extracted balls of color i (i = 1, ..., k) as $X_i$ , and denote as $p_i$ the probability that a given extraction will be in color i. The probability mass function of this multinomial distribution is:

Probability mass function [edit]

 $= \left\{ egin{array}{ll} rac{n!}{x_1! \cdots x_k!} p_1^{x_1} imes \cdots imes p_k^{x_k}, & ext{when } \sum_{i=1}^k x_i = n \ 0 & ext{otherwise}. \end{array} 
ight.$ for non-negative integers  $x_1, ..., x_k$ .

$$f(x_1,\ldots,x_k;p_1,\ldots,p_k)=\frac{\Gamma(\sum_i x_i+1)}{\prod_i \Gamma(x_i+1)}\prod_{i=1}^k p_i^{x_i}.$$
 This form shows its resemblance to the Dirichlet distribution, which is its conjugate prior. 
$$\textbf{Visualization} \quad \texttt{[edit]}$$

### slices of Pascal's pyramid, or 3D/4D/+ (pyramid-shaped) slices of higher-dimensional analogs of Pascal's triangle. This reveals an interpretation of the range of the distribution: discretized equilateral "pyramids" in arbitrary dimension—i.e. a simplex with a grid. [citation needed]

Properties [edit] The expected number of times the outcome i was observed over n trials is

The off-diagonal entries are the covariances:

All covariances are negative because for fixed n, an increase in one component of a multinomial vector requires a decrease in another component. When these expressions are combined into a matrix with i, j element  $cov(X_i, X_j)$ , the result is a  $k \times k$  positive-semidefinite covariance matrix of rank k-1. In the special case where

 $\rho(X_i, X_j) = \frac{\text{Cov}(X_i, X_j)}{\sqrt{\text{Var}(X_i) \text{Var}(X_j)}} = \frac{-p_i p_j}{\sqrt{p_i (1 - p_i) p_j (1 - p_j)}} = -\sqrt{\frac{p_i p_j}{(1 - p_i) (1 - p_j)}}.$ 

Matrix notation [edit] In matrix notation,

 $Var(\mathbf{X}) = n\{diag(\mathbf{p}) - \mathbf{p}\mathbf{p}^{\mathrm{T}}\},\$ with  $\mathbf{p}^{\mathrm{T}}$  = the row vector transpose of the column vector  $\mathbf{p}$ .

voters are selected randomly, what is the probability that there will be exactly one supporter for candidate A, two supporters for candidate B and three supporters for candidate C in the sample? Note: Since we're assuming that the voting population is large, it is reasonable and permissible to think of the probabilities as unchanging once a voter is selected for the sample. Technically speaking this is sampling without replacement, so the correct distribution is the multivariate hypergeometric distribution, but the distributions converge as the population grows

 $j=\min\left\{j'\in\{1,\ldots,k\}:\left(\sum_{i=1}^{j}p_i
ight)-X\geq 0
ight\}.$ 

To simulate from a multinomial distribution [edit] Various methods may be used to simulate from a multinomial distribution. A very simple solution is to use a uniform pseudo-random number generator on (0,1). First, we divide the (0,1) interval in k subintervals equal in length to the probabilities of the k categories. Then, we generate n independent pseudo-random numbers to determine in which of the k intervals they occur and count the number of occurrences in each interval.

 $\{X_i = 1, X_k = 0 \text{ for } k \neq j\}$  is one observation from the multinomial distribution with  $p_1, \ldots, p_k$  and n = 1. A sum of independent repetitions of this experiment is an observation from a

## 0.15 | 0.20 | 0.30 | 0.16 | 0.12 | 0.07 Superior limits of subintervals | 0.15 | 0.35 | 0.65 | 0.81 | 0.93 | 1.00

Formulae: Rand() = If(\$Ai<0.15;1;0) = If(And(\$Ai>=0.15;\$Ai<0.35);1;0)

specified multinomial distribution or a parametric family of multinomial distributions.

Another way is to use a discrete random number generator. In that case, the categories must be labeled or relabeled with numeric values.
In the two cases, the result is a multinomial distribution with k categories. This is equivalent, with a continuous random distribution, to simulate k independent standardized normal
distributions, or a multinormal distribution $N(0,I)$ having $k$ components identically distributed and statistically independent.
Since the counts of all categories have to sum to the number of trials, the counts of the categories are always negatively correlated. <sup>[1]</sup>

Let q denote a theoretical multinomial distribution and let p be a true underlying distribution. The distributions p and q are considered equivalent if  $d(p,q)<\varepsilon$  for a distance d and a tolerance parameter  $\varepsilon>0$ . The equivalence test problem is  $H_0=\{d(p,q)\geq\varepsilon\}$  versus  $H_1=\{d(p,q)<\varepsilon\}$ . The true underlying distribution p is unknown. Instead, the counting frequencies  $p_n$  are observed, where n is a sample size. An equivalence test uses  $p_n$  to reject  $H_0$ . If  $H_0$  can be rejected then the equivalence between p and q is shown at a given

• When k = 2, the multinomial distribution is the binomial distribution. • Categorical distribution, the distribution of each trial; for k = 2, this is the Bernoulli distribution. The Dirichlet distribution is the conjugate prior of the multinomial in Bayesian statistics. Dirichlet-multinomial distribution. Beta-binomial model.

- supported on a bounded interval raised cosine · reciprocal · triangular · U-quadratic · uniform · Wigner semicircle Benini · Benktander 1st kind · Benktander 2nd kind · beta prime · Burr · chi-squared · chi · Dagum · Davis · exponential-logarithmic · Erlang · exponential · F · folded normal · Fréchet · gamma · gamma/Gompertz · generalized gamma · generalized inverse Gaussian · Gompertz · half-logistic · half-normal · Hotelling's T-squared ·