Study	logOR [95%CI]	P-value	
ShiuanKidneyPD-(L)1, n = 13	-2.16 [-4.58; 0.26]	0.08	
PuchMelanomaPD-(L)1, n = 49	-1.63 [-2.78; -0.49]	0.01	←■
GideMelanomaPD-(L)1, n = 38	-1.18 [-2.11; -0.25]	0.01	-
KimGastricPD-(L)1, n = 31	-0.87 [-1.77; 0.03]	0.06	
JungLungPD-(L)1, n = 27	-0.46 [-1.38; 0.47]	0.34	_
MariathasanBladderPD-(L)1, n = 133	3 -0.41 [-0.79; -0.04]	0.03	
MariathasanKidneyPD-(L)1, n = 46	-0.30 [-1.09; 0.48]	0.45	-
SnyderUreteralPD-(L)1, n = 18	-0.29 [-1.30; 0.71]	0.56	-
HugoMelanomaPD-(L)1, n = 27	-0.28 [-1.06; 0.51]	0.49	-
BraunKidneyPD-(L)1, n = 142	0.02 [-0.35; 0.39]	0.92	
RiazMelanomaPD-(L)1, n = 16	0.10 [-1.14; 1.33]	0.88	
Limagne2LungPD-(L)1, n = 22	0.25 [-0.67; 1.17]	0.59	
Miao1KidneyPD-(L)1, n = 28	0.55 [-0.34; 1.44]	0.22	
MariathasanUreteralPD-(L)1, n = 18	0.78 [-0.40; 1.97]	0.20	-
Total	-0.30 [-0.59; -0.01]		

-2-1 0 1 2 logOR estimate

Heterogeneity: χ^2_{13} = 24.06 (P = .03), I^2 = 46% [0%; 71%] Test for overall effect: z = -2.05 (P = .04)