

# Sample title

Brandon Hosley

University of Illinois - Springfield

September 3, 2020

# Overview

1 Q1: Bias–variance tradeoff

2 Q2: Hastie and Tibshirani

# Bias-Variance Trade-Off

Bias Error

- Also called 'Overfitting'

Variance

# Bias-Variance Trade-Off

## Bias Error

- Also called 'Overfitting'

## Variance

- Also called 'Underfitting'


# Bias-Variance Trade-Off

Bias Error

- Also called 'Overfitting'

Variance

- Also called 'Underfitting'




# Bias-Variance Trade-Off

## Bias Error

- Also called 'Overfitting'
- Predicts test data too well

## Variance

- Also called 'Underfitting'




# Bias-Variance Trade-Off

## Bias Error

- Also called 'Overfitting'
- Predicts test data too well

## Variance


- Also called 'Underfitting'
- Generalizes too well



# Bias-Variance Trade-Off

- Aiming for the lowest total error typically means finding a "middle-ground"
- A common technique for this is determining the minimum *mean squared error*.

$$\begin{aligned}\text{MSE} &= \left( E \left[ \hat{f}(x) \right] - f(x) \right)^2 + E \left[ \left( \hat{f}(x) - E \left[ \hat{f}(x) \right] \right)^2 \right] + \sigma_e^2 \\ &= \text{Bias}^2 + \text{Variance} + \text{Irreducible Error}\end{aligned}$$





# Hastie Lectures

- Statistical Learning and Regression
- Dimensionality and Parametric Models
- Assessing Model Accuracy and Bias-Variance Trade-Off
- Classification Problems and K-Nearest Neighbors

# Statistical Learning and Regression

- Predicting values through various *means*

# Statistical Learning and Regression

- Predicting values through various *means*
- The simplest method of regression may be to use:

$$\hat{f}(x_n) = \bar{x}_n$$

# Statistical Learning and Regression

- Predicting values through various *means*
- The simplest method of regression may be to use:

$$\hat{f}(x_n) = \bar{x}_n$$

- This type of model may be improved by extending the values contributing to the average to:

$$\hat{f}(x_n) = \bar{x}_{n \pm i}$$

# Statistical Learning and Regression

- Predicting values through various *means*
- The simplest method of regression may be to use:

$$\hat{f}(x_n) = \bar{x}_n$$

- This type of model may be improved by extending the values contributing to the average to:

$$\hat{f}(x_n) = \bar{x}_{n \pm i}$$

- Or the range may be adjusted such that it includes the  $k$ -nearest neighbors.

# Statistical Learning and Regression


- Predicting values through various *means*
- The simplest method of regression may be to use:

$$\hat{f}(x_n) = \bar{x}_n$$

- This type of model may be improved by extending the values contributing to the average to:

$$\hat{f}(x_n) = \bar{x}_{n \pm i}$$

- Or the range may be adjusted such that it includes the  $k$ -nearest neighbors.




# Dimensionality and Parametric Models

- Approximate proportion of a data's range needed to capture a certain fraction of the data scales very quickly with added dimensions

# Dimensionality and Parametric Models

- Approximate proportion of a data's range needed to capture a certain fraction of the data scales very quickly with added dimensions





# Dimensionality and Parametric Models

- Approximate proportion of a data's range needed to capture a certain fraction of the data scales very quickly with added dimensions
- Structured models may help address these scaling problems

# Dimensionality and Parametric Models

- Approximate proportion of a data's range needed to capture a certain fraction of the data scales very quickly with added dimensions
- Structured models may help address these scaling problems

The simplest structured model is a linear model:

$$f_L(X) = \beta_0 + \beta_1 X_1 + \cdots + \beta_n X_n$$

# Dimensionality and Parametric Models

- Approximate proportion of a data's range needed to capture a certain fraction of the data scales very quickly with added dimensions

- Structured models may help address these scaling problems

The simplest structured model is a linear model:

$$f_L(X) = \beta_0 + \beta_1 X_1 + \cdots + \beta_n X_n$$

- Structured models allowed allow flexible adjustment to obtain preferred level of fitting

# Assessing Model Accuracy and Bias-Variance Trade-Off

# Classification Problems and K-Nearest Neighbors