

Statistical Theory Homework 1

Bohao Tang

August 31, 2017

1.
 - (a) $\{(0, g), (0, f), (0, s), (1, g), (1, f), (1, s)\}$
 - (b) $\{(0, s), (1, s)\}$
 - (c) $\{(0, g), (0, f), (0, s)\}$
 - (d) $\{(0, s), (1, s), (1, g), (1, f)\}$
 - (e) $\{(1, s)\}$
2.
 - (a) $A \cap B^c \cap C^c$
 - (b) $A \cap B \cap C^c$
 - (c) $A \cap B \cap C$
 - (d) $A \cup B \cup C$
 - (e) $(A \cap B \cap C^c) \cup (A \cap B^c \cap C) \cup (A^c \cap B \cap C)$
 - (f) $A^c \cap B^c \cap C^c$
3.
 - (a) $\mathbf{P}(\text{ at least a 6 }) = \frac{6+6-1}{6 \times 6} = \frac{11}{36}$
 - (b) $\mathbf{P}(\text{ same number }) = \frac{6}{6 \times 6} = \frac{1}{6}$
4. It's the same number as to choose 2 people out of 20 to shake hands, and that's $\binom{20}{2} = 190$.
5.
 - (a) $\frac{3 \times 4}{\binom{5}{2} \times 2!} = \frac{3}{5}$
 - (b) $\frac{2 \times 3 + 3 \times 2}{\binom{5}{2} \times 2!} = \frac{3}{5}$
 - (c) $\frac{\binom{3}{2} \times 2!}{\binom{5}{2} \times 2!} = \frac{3}{10}$
6. I think it's to choose n ordered character from a-z , where n is the sum of the numbers of one's last and given name. Then:
 - (a) $26 \times 26 \times 26 = 17576$
 - (b) $26 \times 26 + 26 \times 26 \times 26 = 18252$
 - (c) $26 \times 26 + 26 \times 26 \times 26 + 26 \times 26 \times 26 \times 26 = 475228$
7. We can deal $\{1, 2, 3\}$ as one number and arrange the $n - 2$ numbers in order first, then we arrange 1, 2, 3 in order. Therefore: $\mathbf{P} = \frac{(n-2)! \times 3!}{n!} = \frac{6}{n(n-1)}$, where n must no less than 3.

8. It's equivalent to choose 7 children from 10 to have one and only one gift, therefore the number of results is $\binom{10}{3} = 120$.
9. It's the sum of ways to choose t elements in (x_1, x_2, \dots, x_n) to be 1, where t ranges from k to n . And that's $\sum_{t=k}^n \binom{n}{t}$
10. $\mathbf{P} = \frac{|\text{choose } r \text{ spaces occupied out of the rest } N-2 \text{ spaces}|}{|\text{choose } r \text{ occupied out of } N|} = \frac{\binom{N-2}{r}}{\binom{N}{r}} = \frac{(N-r)(N-r-1)}{N(N-1)}$
11. I'm a little confused about the meaning of "double" here, I think it means 2 dices are in same number.
- (a) $\mathbf{P} = \frac{2}{6 \times 6} / \frac{6}{6 \times 6} = \frac{1}{3}$
- (b) $\mathbf{P} = \frac{5+5}{6 \times 6} / \frac{6 \times 6 - 6}{6 \times 6} = \frac{1}{3}$
12. $\mathbf{P}(\text{coin with heads and tails} | \text{show up heads}) =$
 $\mathbf{P}(\text{choosed third coin and tossed to be heads}) / \mathbf{P}(\text{tossed to be heads}) =$
 $\frac{1/3 \times 1/2}{1/3 \times 1/2 + 1/3 \times 1 + 1/3 \times 0} = \frac{1}{3}$
13. if the n^{th} stamp is of type i , then it's new means the first $n-1$ stamps are of types other than i , whose probability is $(1-p_i)^{n-1}$. Therefore,

$$\mathbf{P}(\text{new}) = \sum_1^m \mathbf{P}(\text{new} | n^{\text{th}} \text{ of type } i) \mathbf{P}(n^{\text{th}} \text{ of type } i) = \sum_1^m p_i (1-p_i)^{n-1}$$

Extra Credit:

Denote $A_{n,r,k}$ to be