

Sleep Hygiene and Mental Health

How sleep affects well-being and evidence-based habits for better rest

Introduction

Sleep is essential for both physical and mental health, yet many people struggle to get enough good-quality rest. This document explains why sleep matters, how sleep and mental health are connected, common sleep problems, and evidence-based habits, often called sleep hygiene, that can help improve sleep quality.

What is Sleep Hygiene?

While sleeping, the brain and body remain active. Sleep is a natural process that helps the body restore energy, supports learning and memory, and keeps people healthy. During sleep, the brain cycles through different stages of non-rapid eye movement (non-REM) sleep and rapid eye movement (REM) sleep, during which brain waves, breathing, heart rate, and body temperature change throughout the night. All stages of sleep are needed for healthy, restorative rest.

(Source: MedlinePlus)

Sleep needs vary by age, lifestyle, and health, but general guidance suggests most adults need about seven to eight hours of sleep a night for good health and mental functioning, with some adults needing up to nine hours. It is a common misconception that adults need less sleep as they age; in fact, older adults often simply get less sleep or spend less time in deep, restful stages of sleep and wake more easily.

(Source: MedlinePlus)

The Mental Health Foundation (UK) describes sleep and mental health as closely related: living with a mental health condition can affect sleep, and poor sleep can, in turn, affect mental health. Quality sleep supports balanced emotions, focus, and overall well-being, while poor sleep can contribute to or worsen conditions such as anxiety and depression.

(Source: Mental Health Foundation UK)

How Mental Health Conditions Can Affect Sleep

- Anxiety can cause racing thoughts, which can make it hard to fall asleep.
- Depression can lead to oversleeping, such as sleeping late or a lot during the day, but it can also cause insomnia if troubling thoughts interfere with sleep.
- Post-traumatic stress disorder (PTSD) can cause nightmares or night terrors that disturb sleep and create anxiety about falling asleep again.

- Mania can cause a person to feel elated or energetic, along with racing thoughts, making it hard to feel tired or want to sleep.
- Certain medications used to treat mental health conditions, or coming off them, can cause side effects such as insomnia, nightmares, or oversleeping.

(Source: Mental Health Foundation UK)

Common Sleep Problems

There are many different types of sleep disorders, which are conditions that disturb normal sleep patterns.

(Source: MedlinePlus)

- Insomnia: trouble falling asleep or staying asleep, or both; it can be acute (short-term), often caused by stress, or chronic, lasting a month or longer.
- Sleep-related breathing disorders, such as sleep apnea, in which breathing stops for ten seconds or more during sleep.
- Sleep-related movement disorders, such as restless leg syndrome, which involves uncomfortable sensations in the legs and a strong urge to move them.
- Hypersomnia: extreme daytime sleepiness, including conditions such as narcolepsy.
- Circadian rhythm sleep-wake disorders: problems falling asleep or waking up at appropriate times, often related to shift work or jet lag.
- Parasomnia: unusual behaviors during sleep or while falling asleep or waking up, such as sleepwalking or talking during sleep.

(Source: MedlinePlus)

Signs that a person may have a sleep disorder include regularly taking more than 30 minutes to fall asleep, waking up several times with trouble falling back to sleep, waking up too early, feeling sleepy during the day, or a bed partner noticing loud snoring, gasping, or pauses in breathing during sleep.

(Source: MedlinePlus)

Causes of Poor Sleep

- Other health conditions, such as heart disease, lung disease, nerve disorders, or chronic pain.
- Mental health conditions, including depression and anxiety.
- Stimulants such as caffeine, nicotine, and alcohol, especially when consumed close to bedtime.
- Irregular sleep schedules, such as those caused by shift work or frequent travel across time zones.
- Poor sleep habits established over time, since sleep habits learned in childhood can carry into adulthood.

(Source: MedlinePlus)

The Mental Health Foundation (UK) notes that life circumstances also shape sleep quality; for example, financial or social strain can reduce both sleep quality and duration, and caring for young children is strongly associated with disrupted sleep.

(Source: Mental Health Foundation UK)

Common Symptoms of Poor Sleep

- Difficulty falling asleep, staying asleep, or waking up too early.
- Feeling tired, unfocused, or irritable during the day.
- Difficulty concentrating or making decisions.
- Low mood or reduced ability to regulate emotions.
- Reduced immune function and increased vulnerability to illness.

(Source: MedlinePlus, Mental Health Foundation UK)

Diagnosis

To diagnose a sleep disorder, a health care provider will review medical and sleep history and perform a physical exam. A sleep study, called a polysomnogram, may be used to monitor and record data about the body during a full night of sleep, including brain activity, breathing, and movement. Other sleep studies may assess how quickly a person falls asleep during daytime naps or how well they can stay awake during the day.

(Source: MedlinePlus)

Evidence-Based Treatment Options

Treatment for sleep problems depends on the specific cause or disorder.

(Source: MedlinePlus)

- Practicing good sleep habits and lifestyle changes, such as eating a healthy diet and getting regular exercise.
- Cognitive behavioral therapy for insomnia (CBT-I): a structured talk therapy that helps address anxiety about sleep and unhelpful thoughts and behaviors related to sleep.
- Relaxation techniques to reduce anxiety around bedtime.
- Continuous positive airway pressure (CPAP) devices for sleep apnea.
- Treating an underlying medical or mental health condition, such as depression, which often improves sleep.

- Sleep medicines, when used, are generally intended for short-term use, since many can become habit-forming and long-term changes to lifestyle and sleep habits are considered the best approach for ongoing sleep problems.

(Source: MedlinePlus)

Healthy Lifestyle Recommendations

Good sleep hygiene refers to healthy habits that support consistent, high-quality sleep.

(Source: WHO)

- Be consistent: go to bed at the same time each night and get up at the same time each morning, including on weekends.
- Make the sleeping area quiet, dark, relaxing, and at a comfortable temperature.
- Limit the use of electronic devices, such as TVs, computers, and smartphones, before sleeping.
- Avoid large meals, caffeine, and alcohol close to bedtime.
- Get regular physical activity during the day, since being active can help you fall asleep more easily at night.

(Source: WHO)

Self-Help Techniques

- Establish a regular, relaxing bedtime routine that signals to the brain that it is time to sleep.
- If you cannot sleep, do not lie in bed worrying; get up and do something relaxing until you feel sleepy, then return to bed.
- Keep a sleep diary to track sleep patterns and notice which habits or circumstances affect sleep quality.
- Address worries before bed, for example by writing down concerns or a to-do list earlier in the evening rather than at bedtime.
- Reserve the bed primarily for sleep, so the brain associates it with rest.

(Source: Mental Health Foundation UK, MedlinePlus)

When to Seek Professional Help

Most sleep problems resolve within about a month, but longer stretches of poor sleep can seriously affect daily life and mental health.

(Source: Mental Health Foundation UK)

- Sleep problems that persist for a month or longer.
- Chronic insomnia lasting a month or more, especially if it is a symptom of another underlying condition.

- Loud snoring, gasping, or pauses in breathing during sleep, which may indicate sleep apnea and requires medical evaluation.
- Excessive daytime sleepiness that interferes with safety or daily functioning, such as difficulty staying awake while driving.
- Sleep problems occurring alongside symptoms of depression, anxiety, or another mental health condition.

(Source: MedlinePlus, Mental Health Foundation UK)

A health care provider can help identify whether a sleep problem is linked to another medical or mental health condition and can recommend appropriate treatment, which may include therapy, lifestyle changes, or, in some cases, medical devices or short-term medication.

(Source: MedlinePlus)

Myths vs Facts

- Myth: Adults need less sleep as they get older. Fact: There is no strong evidence that older adults need less sleep; they often simply get less sleep or spend less time in deep, restful stages.
- Myth: You can "catch up" on lost sleep over the weekend. Fact: Consistent sleep and wake times support the body's internal clock better than irregular sleep patterns, including sleeping in on weekends.
- Myth: Alcohol helps you sleep better. Fact: While alcohol may help some people fall asleep faster, it disrupts the structure of sleep and can lead to more fragmented, lower-quality rest.
- Myth: Sleep problems are only a physical health issue. Fact: Sleep and mental health are closely linked in both directions; mental health conditions can disrupt sleep, and poor sleep can worsen mental health.

(Source: MedlinePlus, Mental Health Foundation UK)

Frequently Asked Questions

Q: How much sleep do adults need?

A: Most adults need about seven to eight hours of sleep a night for good health and mental functioning, though individual needs can vary.

(Source: MedlinePlus)

Q: Can poor sleep cause anxiety or depression?

A: Poor sleep and mental health conditions have a two-way relationship: mental health conditions such as anxiety and depression can disrupt sleep, and ongoing poor sleep can, in turn, worsen mood and mental health.

(Source: Mental Health Foundation UK)

Q: What should I do if I can't fall asleep?

A: Rather than lying in bed worrying, it can help to get up and do something relaxing until you feel sleepy, then return to bed; addressing worries earlier in the evening, rather than at bedtime, can also help.

(Source: Mental Health Foundation UK)

Q: Are sleeping pills a good long-term solution?

A: Sleep medicines can help for short periods, but many can become habit-forming, and long-term changes to lifestyle and sleep habits are generally considered the best approach for ongoing sleep problems.

(Source: MedlinePlus)

Key Takeaways

- Sleep is essential for physical health, mental health, learning, and memory.
- Sleep and mental health have a two-way relationship: each can affect the other.
- Common sleep problems include insomnia, sleep apnea, restless leg syndrome, and circadian rhythm disorders.
- Good sleep hygiene, including a consistent schedule, a comfortable sleep environment, and limiting caffeine, alcohol, and screens before bed, supports better sleep.
- Cognitive behavioral therapy for insomnia is an evidence-based treatment option for chronic sleep problems.
- Sleep problems lasting a month or longer, or occurring alongside mental health symptoms, warrant professional evaluation.

This document provides general educational information based on publicly available guidance from the World Health Organization, MedlinePlus, and the Mental Health Foundation (UK). It is not a substitute for personalized medical advice, diagnosis, or treatment. Always consult a qualified health care provider with questions about a specific medical condition.

References

- Stress — World Health Organization, Questions and Answers (sleep hygiene guidance). <https://www.who.int/news-room/questions-and-answers/item/stress>
- Healthy Sleep — MedlinePlus. <https://medlineplus.gov/healthysleep.html>
- Sleep Disorders — MedlinePlus. <https://medlineplus.gov/sleepdisorders.html>
- Insomnia — MedlinePlus. <https://medlineplus.gov/insomnia.html>
- Insomnia — MedlinePlus Medical Encyclopedia. <https://medlineplus.gov/ency/article/000805.htm>
- Sleep and your health — MedlinePlus Medical Encyclopedia. <https://medlineplus.gov/ency/patientinstructions/000871.htm>

- Sleep and mental health — Mental Health Foundation (UK).
<https://www.mentalhealth.org.uk/explore-mental-health/a-z-topics/sleep-and-mental-health>
- How to sleep better — Mental Health Foundation (UK).
<https://www.mentalhealth.org.uk/explore-mental-health/publications/how-sleep-better>