

Anxiety and Anxiety Disorders

What anxiety is, why it happens, and evidence-based ways to manage it

Introduction

Anxiety is a natural human response to stress or perceived danger, and it is something almost everyone experiences from time to time. This document explains the difference between normal, everyday anxiety and an anxiety disorder, describes the main types of anxiety disorders, and summarizes evidence-based approaches to treatment and self-management. It is written for adults seeking accurate, non-commercial information about anxiety.

What is Anxiety?

Anxiety is a feeling of fear, dread, and uneasiness. It can cause physical reactions such as sweating, restlessness, tension, and a rapid heartbeat. Anxiety can be a normal reaction to stress, for example before a difficult problem at work, a test, or an important decision, and in these situations it can actually help a person cope by providing a boost of energy or improved focus.

(Source: MedlinePlus)

The Mental Health Foundation (UK) describes anxiety as part of the body's natural "fight or flight" response, which helps people react quickly to potential threats. It is normal to feel anxious sometimes, and anxiety is not necessarily a bad thing: it can spur a person on, help them stay alert, and motivate them to solve problems. Anxiety becomes a potential problem when it affects a person's ability to live their life, or when it is ongoing, intense, hard to control, or out of proportion to the situation.

(Source: Mental Health Foundation UK)

For people with an anxiety disorder, the fear is not temporary and can become overwhelming. Anxiety disorders are conditions in which anxiety does not go away and can get worse over time, with symptoms that interfere with daily activities such as job performance, schoolwork, and relationships.

(Source: MedlinePlus)

Common Types of Anxiety Disorders

- Generalized anxiety disorder (GAD): persistent, excessive worry about ordinary issues such as health, money, work, or family, occurring most days for at least six months.
- Panic disorder: recurrent, unexpected panic attacks, which are sudden episodes of intense fear along with physical symptoms such as a racing heart, sweating, or shortness of breath.
- Social anxiety disorder (social phobia): intense, persistent fear of being watched, judged, or embarrassed in social situations.

- Specific phobias: an intense, persistent fear of a specific object or situation, such as heights, flying, or certain animals, that is out of proportion to the actual danger.
- Agoraphobia: intense fear or anxiety about places or situations where escape or help might be difficult if panic-like symptoms occur.
- Separation anxiety disorder: excessive fear or anxiety about being apart from attachment figures.

(Source: NIMH, MedlinePlus)

Causes

The exact cause of most anxiety disorders is not fully understood, but research points to a combination of biological and environmental contributors. Genetics may play a role, since anxiety disorders can run in families, and differences in brain structure and function have been observed between people with and without anxiety disorders. Stressful or traumatic life experiences are also recognized contributors.

(Source: NIMH, MedlinePlus)

The Mental Health Foundation (UK) notes that the causes of anxiety are often complex and can involve a bidirectional relationship with social and environmental factors. Negative life events, such as divorce, abuse, a stressful work or education environment, or a traumatic event like an accident or assault, can contribute to anxiety. Financial strain is also strongly associated with anxiety, as is pressure to meet societal or cultural expectations around appearance, achievement, or success.

(Source: Mental Health Foundation UK)

Risk Factors

- A family history of anxiety disorders or other mental health conditions.
- Exposure to stressful or traumatic events, including in childhood.
- Ongoing stress, such as financial strain, relationship difficulties, or a demanding work or study environment.
- Certain personality traits, such as a tendency toward negative thinking or difficulty tolerating uncertainty.
- Co-occurring physical or mental health conditions, including depression and substance use.
- Younger age: in UK survey data, anxiety levels were highest among adults aged 18 to 34.

(Source: NIMH, MedlinePlus, Mental Health Foundation UK)

Common Symptoms

Anxiety can affect a person physically, mentally, and behaviorally, and symptoms vary from person to person. Cognitive and emotional symptoms can include irritability, difficulty concentrating, and feeling restless, worried, or unable to sit still. Behavioral changes can include avoiding places, situations, or everyday activities that trigger anxiety.

(Source: MedlinePlus)

- Excessive worry that is difficult to control.
- Restlessness, feeling on edge, or being easily fatigued.
- Muscle tension, headaches, or unexplained aches.
- Rapid heartbeat, sweating, trembling, or shortness of breath.
- Difficulty concentrating or a mind that goes blank.
- Sleep problems, including trouble falling or staying asleep.
- Avoidance of situations, places, or activities that trigger anxious feelings.

(Source: MedlinePlus, Mental Health Foundation UK)

A panic attack is a distinct episode of sudden, intense fear accompanied by physical symptoms; it usually peaks within ten to twenty minutes. An isolated panic attack is not itself a mental disorder, but frequent, unexpected panic attacks along with ongoing worry about future attacks may indicate panic disorder.

(Source: NIMH)

Diagnosis

To diagnose an anxiety disorder, a health care provider will ask about symptoms and medical history. A physical exam and laboratory tests may be used to rule out other health problems that could be causing the symptoms. If no other health condition explains the symptoms, the provider may conduct, or refer the person for, a psychological evaluation.

(Source: MedlinePlus)

For generalized anxiety disorder specifically, a provider will typically ask about the duration, frequency, and impact of worry; a diagnosis generally requires frequent worry or tension lasting at least six months, even when there is little or no clear cause, along with associated physical symptoms such as muscle tension or difficulty sleeping.

(Source: MedlinePlus)

Evidence-Based Treatment Options

Anxiety disorders are treatable, and most people who receive appropriate treatment see significant improvement. The main treatments are psychotherapy, medication, or a combination of the two, and the best approach depends on the specific disorder, its severity, and individual preferences.

(Source: NIMH, MedlinePlus)

- Cognitive behavioral therapy (CBT): a structured talk therapy that teaches people to recognize and change distorted thoughts and behaviors, helping them respond differently to situations that trigger fear or anxiety.

- Exposure therapy: a gradual, structured approach in which a person is progressively exposed to a feared object or situation to reduce avoidance and fear over time.
- Medications: including certain antidepressants such as selective serotonin reuptake inhibitors (SSRIs) and serotonin-norepinephrine reuptake inhibitors (SNRIs), as well as other anti-anxiety medicines; benzodiazepines act quickly but can become less effective and habit-forming with long-term use, so they are generally prescribed cautiously and for limited periods.
- Beta blockers: sometimes used to relieve physical symptoms of anxiety, such as a rapid heartbeat or trembling.

(Source: NIMH, MedlinePlus)

Finding the most effective treatment can take some trial and error, and both psychotherapy and medication can take time to produce noticeable benefits. Choosing a treatment plan should be done in consultation with a health care provider or mental health professional, based on individual needs and preferences.

(Source: NIMH)

Healthy Lifestyle Recommendations

A healthy lifestyle can help manage anxiety symptoms and support the effectiveness of formal treatment, although lifestyle changes alone are not usually a substitute for professional care.

(Source: NIMH)

- Getting regular physical activity, which can reduce anxiety symptoms and improve mood.
- Getting enough good-quality sleep on a consistent schedule.
- Reducing caffeine intake, since caffeine can trigger or worsen anxiety symptoms.
- Limiting or avoiding alcohol, since it can worsen anxiety over time despite short-term relief.
- Eating regular, balanced meals to help keep energy levels stable.
- Maintaining a strong social support system through friends, family, or support groups.

(Source: NIMH, MedlinePlus)

Self-Help Techniques

In addition to professional treatment, several self-help strategies can help people manage day-to-day anxiety symptoms.

- Practicing relaxation techniques, such as deep breathing exercises, which can help calm the body's stress response during moments of acute anxiety.
- Practicing mindfulness, which involves paying attention to the present moment; note that some sources indicate mindfulness is not recommended specifically for social anxiety.
- Talking about feelings with a trusted friend, family member, or counselor, since being heard and understood can reduce distress.

- Joining a peer support group to connect with others who have similar experiences.
- Using structured self-help resources, such as guided cognitive behavioral therapy tools, books, or online programs.
- Regular exercise, such as walking, running, swimming, or yoga, which can reduce anxiety and improve sleep.

(Source: Mental Health Foundation UK, NIMH)

When to Seek Professional Help

It is normal to feel anxious at times, and this does not always mean a person has an anxiety disorder. However, professional help should be sought if anxiety is ongoing, intense, difficult to control, out of proportion to the situation, or if it interferes with daily life, work, relationships, or school.

(Source: Mental Health Foundation UK, MedlinePlus)

- Anxiety symptoms last a long time, get worse, or interfere with day-to-day life.
- Physical symptoms, such as chest pain or shortness of breath, are severe or new, since these may also signal another medical condition requiring urgent evaluation.
- A person is avoiding important activities, relationships, or responsibilities because of anxiety.
- A person is using alcohol or drugs to cope with anxiety.
- A person has thoughts of self-harm or suicide, which requires immediate help.

(Source: NIMH, MedlinePlus, Mental Health Foundation UK)

A primary care provider is a good starting point and can refer a person to a mental health specialist, such as a psychologist, psychiatrist, or therapist, for further evaluation and treatment.

(Source: MedlinePlus)

Myths vs Facts

- Myth: Anxiety is always a sign of a disorder. Fact: Anxiety is a normal and often useful emotion; it becomes a disorder only when it is persistent, excessive, and interferes with daily functioning.
- Myth: People with anxiety disorders just need to relax more. Fact: Anxiety disorders are recognized medical conditions with biological and environmental contributors, and they typically require evidence-based treatment such as therapy or medication.
- Myth: Anxiety disorders are rare. Fact: Anxiety disorders are the most common form of mental illness in the United States, and about one third of U.S. adults and adolescents experience an anxiety disorder at some point in their lives.
- Myth: Medication is the only effective treatment. Fact: Psychotherapy, particularly cognitive behavioral therapy, is an effective evidence-based treatment on its own or combined with medication.

(Source: NIMH, MedlinePlus)

Frequently Asked Questions

Q: Is anxiety the same as an anxiety disorder?

A: No. Anxiety is a normal, common emotion that everyone experiences. An anxiety disorder is a diagnosable mental health condition in which anxiety is persistent, excessive, and interferes with daily life.

(Source: Mental Health Foundation UK)

Q: How long does it take for anxiety treatment to work?

A: Both psychotherapy and medication can take time to show benefits, and finding the most effective treatment plan may involve some trial and error in consultation with a health care provider.

(Source: NIMH)

Q: Can lifestyle changes cure an anxiety disorder on their own?

A: Healthy lifestyle choices, such as exercise, good sleep, and reduced caffeine intake, can help reduce anxiety symptoms and support treatment, but they generally cannot replace evidence-based treatment such as psychotherapy or medication for a diagnosed anxiety disorder.

(Source: NIMH)

Q: Can children have anxiety disorders?

A: Yes. Anxiety disorders can affect people of all ages, including children, and can interfere with school performance, family relationships, and social development.

(Source: MedlinePlus)

Key Takeaways

- Anxiety is a normal human response to stress and becomes a disorder only when it is persistent, excessive, and disruptive to daily life.
- There are several distinct anxiety disorders, including generalized anxiety disorder, panic disorder, social anxiety disorder, phobias, and separation anxiety disorder.
- Anxiety disorders are common, affecting about one third of U.S. adults and adolescents at some point in their lives, and are shaped by a mix of genetic, biological, and environmental factors.
- Effective, evidence-based treatments exist, particularly cognitive behavioral therapy and, when appropriate, medication.
- Healthy lifestyle habits and self-help strategies can support treatment but do not usually replace professional care for a diagnosed anxiety disorder.
- Professional help should be sought when anxiety is persistent, severe, or interferes with daily life.

This document provides general educational information based on publicly available guidance from the National Institute of Mental Health, MedlinePlus, and the Mental Health Foundation (UK). It is not a substitute for personalized medical advice, diagnosis, or treatment. Always consult a qualified health care provider with questions about a specific medical condition.

References

- Anxiety Disorders — National Institute of Mental Health (NIMH).
<https://www.nimh.nih.gov/health/topics/anxiety-disorders>
- Generalized Anxiety Disorder: What You Need to Know — NIMH.
<https://www.nimh.nih.gov/health/publications/generalized-anxiety-disorder-gad>
- Panic Disorder: What You Need to Know — NIMH.
<https://www.nimh.nih.gov/health/publications/panic-disorder-when-fear-overwhelms>
- Any Anxiety Disorder — NIMH. <https://www.nimh.nih.gov/health/statistics/any-anxiety-disorder>
- Anxiety — MedlinePlus. <https://medlineplus.gov/anxiety.html>
- Generalized anxiety disorder — MedlinePlus Medical Encyclopedia.
<https://medlineplus.gov/ency/article/000917.htm>
- Panic disorder — MedlinePlus Medical Encyclopedia. <https://medlineplus.gov/ency/article/000924.htm>
- Social anxiety disorder — MedlinePlus Medical Encyclopedia. <https://medlineplus.gov/ency/article/000957.htm>
- Anxiety — Mental Health Foundation (UK).
<https://www.mentalhealth.org.uk/explore-mental-health/a-z-topics/anxiety>
- What is anxiety? — Mental Health Foundation (UK). <https://www.mentalhealth.org.uk/our-work/public-engagement/mental-health-awareness-week/anxiety-report/what-anxiety>
- What causes anxiety? — Mental Health Foundation (UK). <https://www.mentalhealth.org.uk/our-work/public-engagement/mental-health-awareness-week/anxiety-report/what-causes-anxiety>
- What does anxiety feel like? — Mental Health Foundation (UK). <https://www.mentalhealth.org.uk/our-work/public-engagement/mental-health-awareness-week/anxiety-report/what-does-anxiety-feel-like>