

Depression: Symptoms, Causes, and Treatment

An evidence-based overview of depressive disorders and how they are managed

Introduction

Depression is one of the most common mental health conditions worldwide. It is more than ordinary sadness or a temporary low mood; it is a persistent condition that can affect how a person thinks, feels, sleeps, eats, and functions in daily life. This document explains what depression is, its causes and risk factors, common symptoms, how it is diagnosed, evidence-based treatment options, and practical guidance on self-care and when to seek professional help.

What is Depression?

Depression is a mood disorder characterized by a persistent feeling of sadness, hopelessness, or loss of interest in activities that lasts for at least two weeks and interferes with daily life. It is different from the normal emotional experience of sadness or grief, which is typically shorter and less impairing. Depression is extreme and persistent, and it can significantly interfere with a person's ability to function at work, in relationships, and in daily activities.

(Source: NIMH)

Not everyone who is depressed experiences the same symptoms; some people experience only a few symptoms while others experience many, and the severity, frequency, and duration of symptoms vary from person to person. Depression is not a sign of personal weakness or a character flaw, and a person cannot simply “snap out of” clinical depression.

(Source: NIMH)

The Mental Health Foundation (UK) describes depression as ranging from mild forms, which can make everyday tasks feel harder, to severe depression, which can lead to feelings of hopelessness and, in some cases, suicidal thoughts. Depression is common, and recovery is possible even for people who have experienced it for a long time.

(Source: Mental Health Foundation UK)

Types of Depression

- Major depressive disorder: severe symptoms that interfere with the ability to work, sleep, eat, and enjoy life, occurring most of the day, nearly every day, for at least two weeks.
- Persistent depressive disorder (dysthymia): depressive symptoms that last for at least two years, though they may be less severe than major depression.

- Seasonal affective disorder (SAD): a form of depression related to changes in seasons, most often beginning in late fall or early winter and improving in spring and summer.
- Perinatal depression: depression occurring during pregnancy or after childbirth, marked by severe anxiety, sadness, and fatigue.
- Depression as part of bipolar disorder: bipolar disorder involves episodes of depression alternating with episodes of extreme high mood, or mania; it is distinct from depression alone.

(Source: NIMH)

Causes

Depression is caused by a complex combination of biological, psychological, and environmental factors, and researchers continue to study exactly how these factors interact. Differences in brain structure and function, genetics, and imbalances in brain chemistry are all thought to play a role.

(Source: NIMH)

The Mental Health Foundation (UK) notes that different factors can lead to depression, including genetics, physical health problems, difficult childhood experiences, and stressful life events such as unemployment, the end of a relationship, or experiences of bullying or assault. A combination of factors often contributes to depression, and sometimes there is no single, obvious cause.

(Source: Mental Health Foundation UK)

Risk Factors

- A personal or family history of depression or other mental health conditions.
- Major life changes, trauma, or chronic stress, such as unemployment, bereavement, or relationship breakdown.
- Chronic physical health conditions or chronic pain.
- Certain medications and substance use.
- Difficult childhood experiences, including abuse or neglect.
- Being female: after puberty, women are approximately twice as likely as men to develop depression.
- Major hormonal changes, such as during pregnancy, after childbirth, or during perimenopause.

(Source: NIMH)

Common Symptoms

Depression affects people differently, and symptoms can be emotional, cognitive, and physical. To be diagnosed with depression, a person must have symptoms most of the day, nearly every day, for at least two weeks, and these symptoms must interfere with day-to-day functioning and cause significant distress.

(Source: NIMH)

- Persistent sad, anxious, or “empty” mood.
- Loss of interest or pleasure in hobbies and activities once enjoyed.
- Feelings of hopelessness, worthlessness, or excessive guilt.
- Fatigue or decreased energy.
- Difficulty concentrating, remembering details, or making decisions.
- Sleeping too much or too little, or waking up too early.
- Changes in appetite or unplanned weight changes.
- Physical aches or pains, headaches, cramps, or digestive problems without a clear physical cause that do not go away with treatment.
- Thoughts of death or suicide, or suicide attempts.

(Source: NIMH)

Sadness is only a small part of depression, and some people with depression may not feel sadness at all. Symptoms can also present as physical problems, such as a racing heart, chest tightness, or digestive issues, and men in particular are more likely to seek care for physical symptoms than emotional ones, and may be more likely to use alcohol or drugs as a coping strategy.

(Source: NIMH)

Diagnosis

A health care provider diagnoses depression based on a discussion of symptoms, their duration and impact, and family history. A provider may ask when symptoms began, how long they have lasted, how often they occur, and whether they interfere with usual activities. A physical exam and laboratory tests may be used to rule out other medical conditions, such as thyroid disorders, that can cause similar symptoms.

(Source: NIMH)

To be diagnosed with major depression, symptoms must occur most of the day, nearly every day, for at least two weeks, and at least one symptom must be a depressed mood or loss of interest in most activities.

(Source: MedlinePlus)

Evidence-Based Treatment Options

Depression, even in its most severe forms, can be treated, and the earlier treatment begins, the more effective it tends to be. Most adults see an improvement in symptoms with antidepressant medication, psychotherapy, or a combination of both.

(Source: NIMH)

- **Psychotherapy (talk therapy):** approaches such as cognitive behavioral therapy (CBT) and interpersonal therapy help people recognize and change troubling emotions, thoughts, and behaviors; for milder depression, psychotherapy alone is often tried first.
- **Antidepressant medication:** medicines that may help improve how the brain uses certain chemicals that affect mood; for moderate or severe depression, a combination of medication and therapy is often recommended from the start.
- **Brain stimulation therapies:** options such as electroconvulsive therapy (ECT) or repetitive transcranial magnetic stimulation (rTMS) may be considered if psychotherapy and medication do not sufficiently reduce symptoms.
- **Light therapy:** an evidence-based treatment specifically for seasonal affective disorder.

(Source: NIMH, MedlinePlus)

Choosing a treatment plan should be based on a person's individual needs, preferences, and medical situation, in consultation with a health care provider. Finding the most effective treatment can involve some trial and error, and once treatment begins, most people gradually start to feel better.

(Source: NIMH)

Healthy Lifestyle Recommendations

Alongside professional treatment, certain lifestyle habits can support recovery and help manage symptoms of depression.

(Source: NIMH)

- **Getting physical activity:** even about 30 minutes of walking a day can help boost mood.
- **Maintaining a regular bedtime and wake-up time** to support consistent, quality sleep.
- **Eating regular, healthy meals.**
- **Avoiding alcohol, nicotine, and drugs,** including medications not prescribed for the individual.
- **Trying to do things that were previously enjoyable,** even when motivation is low, since engaging in these activities can help improve mood over time.

(Source: NIMH)

Self-Help Techniques

Self-help strategies can complement professional treatment, particularly for milder depression, though they are not a substitute for treatment when depression is moderate to severe.

(Source: Mental Health Foundation UK)

- **Connecting with people:** talking to trusted friends or family about how you are feeling can help reduce isolation.

- Delaying major life decisions until symptoms improve, and discussing important decisions with people who know you well.
- Trying structured self-help resources, such as self-help books or online cognitive behavioral therapy programs.
- Practicing mindfulness, since some evidence indicates that mindfulness-based approaches, including mindfulness-based cognitive therapy, can help manage depression.
- Doing something creative, such as writing, art, or music, which some people find helpful for processing emotions.
- Keeping a mood diary to track changes in mood and notice which activities, places, or people affect how you feel.

(Source: NIMH, Mental Health Foundation UK)

When to Seek Professional Help

Anyone who shows signs or symptoms of depression that persist or do not go away should talk to a health care provider. If a person notices signs of depression in someone they know, they should encourage that person to seek help from a mental health professional.

(Source: NIMH)

- Symptoms have lasted most of the day, nearly every day, for two weeks or more.
- Symptoms are interfering with work, school, relationships, or daily responsibilities.
- There are thoughts of death, self-harm, or suicide.
- Physical symptoms, such as unexplained pain, persist despite other treatment.

(Source: NIMH)

If a person, or someone they know, is struggling or having thoughts of suicide, they should call or text the 988 Suicide and Crisis Lifeline (in the United States) or contact local emergency services immediately. In the UK, Samaritans can be reached at 116 123, and anyone feeling unable to keep themselves safe should call 999 or go to the nearest emergency department.

(Source: NIMH, Mental Health Foundation UK)

Myths vs Facts

- Myth: Depression is just feeling sad and will pass on its own. Fact: Depression is a persistent medical condition, different from ordinary sadness, that usually requires evidence-based treatment to improve.
- Myth: People with depression can simply choose to feel better. Fact: Depression is not a sign of weakness or a character flaw, and a person cannot simply “snap out of” clinical depression.

- Myth: Antidepressants are the only effective treatment. Fact: Psychotherapy alone is often effective, particularly for milder depression, and many people benefit from a combination of therapy and medication.
- Myth: Depression only affects mood. Fact: Depression frequently causes physical symptoms as well, including fatigue, sleep changes, appetite changes, and unexplained aches or pains.

(Source: NIMH, Mental Health Foundation UK)

Frequently Asked Questions

Q: How is depression different from normal sadness or grief?

A: Normal sadness and grief are usually connected to a specific event and tend to ease over time. Depression is extreme and persistent, lasting most of the day for at least two weeks, and it significantly interferes with a person's ability to function.

(Source: NIMH)

Q: Can depression be cured?

A: Depression, even in its most severe forms, can be treated, and most people who receive appropriate care see significant improvement. Some people experience a single episode, while others may have recurring episodes that require ongoing management.

(Source: NIMH)

Q: Is it normal to have physical symptoms with depression?

A: Yes. Depression can cause physical symptoms such as headaches, digestive problems, fatigue, and changes in sleep or appetite, in addition to emotional symptoms.

(Source: NIMH)

Q: How long does depression treatment take to work?

A: It varies by individual and treatment type. Once treatment begins, people should gradually start to feel better, though finding the most effective treatment plan may take some trial and error.

(Source: NIMH)

Key Takeaways

- Depression is a persistent mood disorder, not ordinary sadness, and it significantly affects daily functioning.
- Depression is caused by a complex mix of biological, psychological, and environmental factors, and it is not a sign of personal weakness.
- Common symptoms include persistent low mood, loss of interest in activities, fatigue, sleep and appetite changes, and difficulty concentrating.

- Evidence-based treatments, including psychotherapy and medication, are effective, and most people improve with appropriate care.
- Self-help strategies, such as staying connected with others and maintaining routine, can support recovery alongside professional treatment.
- Any thoughts of self-harm or suicide require immediate professional help.

This document provides general educational information based on publicly available guidance from the National Institute of Mental Health, MedlinePlus, and the Mental Health Foundation (UK). It is not a substitute for personalized medical advice, diagnosis, or treatment. Always consult a qualified health care provider with questions about a specific medical condition.

References

- Depression — National Institute of Mental Health (NIMH). <https://www.nimh.nih.gov/health/publications/depression>
- Depression — National Institute of Mental Health (NIMH), overview page. <https://www.nimh.nih.gov/health/topics/depression>
- Seasonal Affective Disorder — NIMH. <https://www.nimh.nih.gov/health/publications/seasonal-affective-disorder>
- Depression — MedlinePlus. <https://medlineplus.gov/depression.html>
- Depression — MedlinePlus Medical Encyclopedia. <https://medlineplus.gov/ency/article/003213.htm>
- Depression — Mental Health Foundation (UK). <https://www.mentalhealth.org.uk/explore-mental-health/a-z-topics/depression>