

Dataset 2

Objective : Lifestyle and Well-being Impact on Cancer

Cancer is a multifactorial disease influenced by genetic susceptibility, environmental exposures, and lifestyle-related factors. Although inherited genetic variants can predispose individuals to particular cancers, many important cancer risk factors are potentially modifiable. Lifestyle factors such as tobacco use, alcohol consumption, dietary patterns, physical inactivity, excess body weight, and exposure to ultraviolet (UV) radiation can influence cancer development through mechanisms involving DNA damage, inflammation, metabolism, hormonal regulation, immune function, and the gut microbiome.

The relationship between lifestyle and cancer is therefore not limited to a single behavior. Multiple lifestyle factors can interact with one another and with environmental exposures, influencing both cancer development and overall well-being.

1. Tobacco Use and Smoking

Tobacco smoking is one of the most strongly established lifestyle-related causes of cancer. Cigarette smoke contains carcinogenic substances capable of damaging DNA and producing mutations that can contribute to malignant transformation. The strongest association is observed with **lung cancer**, but tobacco exposure is also associated with cancers of the oral cavity, pharynx, larynx, esophagus, stomach, pancreas, bladder, kidney, and several other organs.

The reviewed evidence reports that **more than 80% of lung-cancer cases were attributed to smoking** in one cited analysis. Smoking was also associated with an approximately **10–20-fold increase in lung-cancer risk**, depending on exposure history.

Smoking has also been identified as a causal factor for **laryngeal, oral, pharyngeal, esophageal, bladder, kidney, and pancreatic cancers**, with additional evidence connecting smoking to stomach cancer, cervical cancer, leukemia, and potentially liver cancer.

The effects of tobacco can be amplified when combined with other risk factors. For example, simultaneous tobacco and alcohol exposure can increase the risk of cancers of the upper aerodigestive tract.

Smoking cessation is therefore one of the most important modifiable strategies for reducing cancer risk.

2. Alcohol Consumption

Alcohol consumption is another important lifestyle-related cancer risk factor. The reviewed sources associate alcohol with cancers of the **oral cavity, pharynx, larynx, esophagus, liver, breast, colorectal region, stomach, and pancreas**.

The relationship can be dose-dependent, with increasing alcohol exposure associated with increasing risk for several cancers. One global analysis reported alcohol-attributable fractions in the approximate **25–50% range among men for several cancers**, including oral/pharyngeal, laryngeal, esophageal, and liver cancers.

Alcohol can contribute to carcinogenesis through several mechanisms. Its metabolism produces **acetaldehyde**, which can damage DNA. Alcohol exposure can also contribute to oxidative stress, inflammation, altered hormone regulation, and changes in cellular metabolism.

Alcohol can additionally interact with other risk factors. For example, its combination with tobacco can increase cancer risk, while alcohol exposure combined with chronic hepatitis infections can increase the risk of liver cancer.

The reviewed sources estimate that approximately **4–5% of cancers** are attributable to alcohol consumption in the populations or analyses cited. These figures should be treated as source-specific population estimates rather than as an individual's personal probability of developing cancer.

3. Diet and Nutrition

Diet represents one of the major lifestyle factors associated with cancer. Dietary patterns can influence cancer risk through their effects on **metabolism, inflammation, body weight, hormone regulation, DNA damage, and the gut microbiome**.

The reviewed sources associate dietary factors particularly with cancers of the digestive system, while also discussing relationships with **colorectal, breast, uterine, ovarian, prostate, stomach, pancreatic, and other cancers**.

Certain dietary patterns involving high consumption of processed foods, excessive saturated and trans fats, refined sugars, refined carbohydrates, and high consumption of red or processed meat have been associated with increased cancer risk. However, the evidence is generally stronger for **overall dietary patterns** than for claiming that any single food directly causes or prevents cancer.

Conversely, healthier dietary patterns generally emphasize **fruits, vegetables, whole grains, dietary fiber, healthy fats, and plant-based or lean protein sources**.

4. Red and Processed Meat

Red meat includes **beef, pork, veal, lamb, mutton, and goat**. Processed meat refers to meat treated through smoking, curing, salting, or fermentation and includes foods such as **bacon, ham, deli meats, sausages, and hot dogs**.

The source identifies red meat as a **probable carcinogen** and processed meat as a **known carcinogen**.

Consumption of red and processed meat is particularly associated with an increased risk of **colorectal cancer**. The sources also describe possible associations with **breast, pancreatic, prostate, and stomach cancers**, although additional research is needed for these relationships.

Several mechanisms may contribute to the association. Red meat contains **heme iron**, which can contribute to processes associated with DNA damage. Cooking meat at high temperatures, such as through grilling, frying, or broiling, can produce compounds capable of causing DNA changes in laboratory studies.

Processed meat may contain **nitrates and nitrites**, which can contribute to the formation of potentially DNA-damaging compounds.

Both red and processed meats can also contain substantial amounts of saturated fat. High consumption of saturated fat may contribute to chronic inflammation and excess body weight, which can further influence cancer risk.

The source states that it is **not known whether there is an amount of red or processed meat that can be consumed without increasing cancer risk**. It reports recommendations to avoid processed meat and to

consume no more than **18 ounces (approximately 510 g, or 3–4 servings) of red meat per week** to reduce cancer risk.

Alternative protein sources include **fish, poultry, beans, peas, lentils, nuts, soy, mushrooms, and low-fat dairy products**.

5. Chronic Inflammation and Cancer

Inflammation is normally a protective response to injury or infection. During acute inflammation, immune cells are recruited to damaged areas to help repair tissue.

However, when inflammation continues for a prolonged period, it can become **chronic inflammation**. Persistent inflammation can damage healthy cells and contribute to biological processes associated with cancer development.

The dietary sources identify frequent consumption of certain foods, including **added sugars and processed meat**, as potential contributors to inflammation.

This provides an important connection between lifestyle and cancer:

Unhealthy dietary patterns → chronic inflammation → cellular damage → increased cancer risk

Inflammation is therefore an important biological pathway connecting diet and other lifestyle factors with cancer.

6. Anti-Inflammatory Dietary Patterns

Research described in the source suggests that anti-inflammatory dietary patterns may help reduce chronic inflammation and potentially lower the risk of certain cancers.

The source identifies possible reductions in risk for:

- **Colorectal cancer**
- **Breast cancer**
- **Prostate cancer**
- **Lung cancer**
- **Stomach cancer**
- **Esophageal cancer**
- **Oral cancer**

These dietary patterns may also provide benefits against other chronic diseases, particularly cardiovascular disease. The source further notes research suggesting that an anti-inflammatory dietary pattern may be associated with improved health and survival among people who have already been diagnosed with cancer. This should be presented as an observed research association rather than as a guaranteed therapeutic effect.

Characteristics of Anti-Inflammatory Diets

The sources identify the **Mediterranean, DASH, and Nordic diets** as examples of dietary patterns consistent with an anti-inflammatory approach.

They generally emphasize:

- A variety of **fruits and vegetables**
- **Whole grains** such as brown rice, barley, quinoa, oats, corn, and millet
- **Healthy fats** from nuts, seeds, fatty fish, and plant oils
- **Fish, poultry, and plant-based proteins**
- Beans, peas, lentils, nuts, soy, and mushrooms
- **Low-fat dairy**, such as yogurt and kefir
- Herbs and spices rather than excessive salt

They generally limit:

- Foods high in **added sugar**
- **Ultra-processed foods**
- Processed meats
- Red meat
- Foods high in saturated and trans fats

Thus, the importance of an anti-inflammatory diet lies primarily in its **overall dietary pattern**, rather than in a single “anti-cancer” food.

7. Dietary Patterns, Plant-Based Foods, and Soy

Plant-based dietary patterns are generally characterized by higher consumption of vegetables, fruits, whole grains, fiber, nuts, seeds, legumes, and other plant foods.

Vegetarian diets are generally lower in saturated fat and higher in fiber and beneficial plant compounds. They also exclude red and processed meats. The reviewed evidence indicates that vegetarians may have a **lower overall risk of cancer**, although it is less clear whether vegetarian diets provide additional benefits compared with other healthy diets containing small amounts of animal products.

Overall, the evidence supports a **predominantly plant-based dietary pattern with limited red and processed meat**.

Soy foods such as **tofu, edamame, and soy milk** provide protein, fiber, healthy fats, potassium, magnesium, and other nutrients. Studies in humans indicate that dietary soy either lowers the risk of some cancers or has no significant effect on cancer risk.

The reviewed evidence also does not indicate that normal dietary soy consumption increases breast-cancer risk. However, soy-isoflavone supplements should not be treated as established cancer-prevention interventions.

8. Gut Microbiome and Immune Function

The **gut microbiome** provides another important connection between lifestyle and cancer. It consists of microorganisms living in and on the body, including bacteria, fungi, and viruses.

The gut microbiome interacts closely with the immune system and can influence immune activity. Because diet and other lifestyle behaviors can alter the composition and function of the microbiome, lifestyle may indirectly affect cancer-related immune responses.

The reviewed research discusses **dietary fiber and probiotics** and their potential effects on the gut microbiome and immune responses associated with cancer.

Research involving melanoma has investigated whether changes in the microbiome can influence anti-tumor immune responses.

This remains an evolving field. Therefore, the relationship between:

Diet » microbiome » immune function » cancer

should be described as an emerging biological pathway rather than as proof that specific probiotics or foods prevent cancer.

9. Food Preparation and Cooking

The methods used to prepare food may influence exposure to chemicals associated with cancer risk.

Cooking red meat, poultry, or fish at very high temperatures through **grilling, smoking, or pan-frying** can produce compounds such as **heterocyclic amines**, which may contribute to cancer risk.

Some preservation methods used for red and processed meats can also involve nitrates and contribute to the formation of potentially carcinogenic **N-nitroso compounds**.

Food preparation therefore represents an additional pathway through which dietary behavior may influence cancer risk.

The reviewed source distinguishes microwave cooking from these high-temperature methods. Microwaves use **non-ionizing radiation** and are not identified as a cause of cancer.

10. Other Food-Related Chemical Exposures

Diet-related cancer risk can also involve exposure to environmental chemicals rather than nutrients themselves.

Acrylamide

Acrylamide can form when the amino acid asparagine reacts with sugars during high-temperature cooking, particularly during frying, baking, and roasting.

Common sources include french fries, potato chips, crackers, bread, cookies, breakfast cereals, and coffee.

The source identifies acrylamide as a **probable carcinogen based mainly on animal evidence**, but studies in humans have not established clear evidence that dietary acrylamide increases cancer risk.

Therefore, acrylamide represents a potential exposure of scientific interest, but the human evidence remains uncertain.

Arsenic

Inorganic arsenic is a recognized human carcinogen. Exposure can occur through contaminated drinking water and certain foods.

The reviewed evidence associates arsenic exposure through contaminated drinking water with increased risks of:

- **Lung cancer**
- **Bladder cancer**
- **Skin cancer**

The risk is particularly relevant in areas where groundwater or other drinking-water sources contain elevated arsenic levels.

Pesticides and Herbicides

Some agricultural chemicals can contribute to cancer risk, particularly under occupational or high-level exposure.

The source identifies **glyphosate, malathion, and diazinon** as probable human carcinogens. These have been associated with increased risk of **non-Hodgkin lymphoma**, while malathion has also been associated with prostate cancer and diazinon with lung cancer.

However, fruits and vegetables remain important components of a healthy diet. Washing conventionally grown produce can help remove some pesticide residues and harmful microorganisms.

11. Food Additives, Supplements, and Dietary Claims

Not every food-related chemical or dietary practice represents an established cancer risk.

Antioxidant Supplements

Although antioxidants obtained from whole foods may contribute to healthy dietary patterns, studies of antioxidant supplements have generally shown **little to no benefit in reducing cancer risk**, and some studies have reported increased cancer risk with certain supplements.

Therefore, whole fruits and vegetables should not be replaced with antioxidant supplements as a cancer-prevention strategy.

Glycemic Index and Glycemic Load

The **glycemic index (GI)** describes how rapidly a carbohydrate-containing food raises blood glucose, while **glycemic load (GL)** considers both the quality and quantity of carbohydrate.

The reviewed evidence suggests that diets with a high glycemic load may increase the risk of **endometrial cancer**. Evidence for other cancers remains less established.

Genetically Modified Foods

The source reports that there is currently **no evidence that approved genetically modified foods increase cancer risk**.

Therefore, genetically modified foods should not be classified as an established cancer risk factor based on the reviewed evidence.

Irradiated Foods

Food irradiation is used to kill microorganisms and insects and improve food safety.

The source states that irradiation **does not make food radioactive** and that there is currently no evidence that food irradiation causes cancer.

Juicing and Detox Diets

Juicing can provide nutrients, but whole fruits and vegetables generally provide more dietary fiber.

The source reports **no scientific evidence that juice cleanses or detox diets reduce cancer risk or remove toxins from the body**. The kidneys and liver already perform the body's normal detoxification processes.

Therefore, restrictive detox or juice-only diets should not be presented as established cancer-prevention strategies.

12. Obesity and Metabolic Health

Excess body weight and obesity are important lifestyle-related cancer risk factors. Obesity has been associated with cancers including **colorectal, breast, endometrial, kidney, esophageal, pancreatic, gallbladder, and liver cancers**.

The relationship is particularly notable for certain cancers. One reviewed global analysis estimated that approximately:

- **29% of esophageal adenocarcinoma in men**
- **37% of esophageal adenocarcinoma in women**

were attributable to overweight or obesity.

For **endometrial cancer**, approximately **36%** was estimated to be attributable to overweight or obesity.

For **gallbladder cancer**, the corresponding estimates were approximately **11% in men and 42% in women**.

Obesity can influence cancer development through several interconnected biological pathways. Excess adipose tissue can alter:

- Insulin
- Insulin-like growth factor-1
- Leptin
- Sex hormones
- Insulin resistance
- Inflammatory signaling

Obesity and metabolic disease may also impair immune function, potentially affecting anti-tumor immune responses.

Therefore, metabolic health represents an important intersection between **lifestyle, inflammation, immunity, and cancer development**.

13. Physical Activity and Sedentary Lifestyle

Physical activity is another important lifestyle factor associated with cancer risk. Physical inactivity has been associated with increased risks of several cancers, particularly **colorectal and breast cancers**.

Regular physical activity may influence cancer risk through effects on:

- Insulin regulation
- Hormone levels
- Energy metabolism
- Immune function
- Gastrointestinal activity
- Body-weight regulation

The reviewed evidence reports that high levels of physical activity were associated with reductions in colorectal-cancer risk of **up to approximately 50%** in some studies.

One cited study found that men who maintained moderate physical activity were **48% less likely to develop colon cancer** than men who remained inactive.

Among initially inactive men who subsequently increased their physical activity, the reported risk was approximately **13% lower** than among those who remained inactive.

However, the evidence is not equally strong for every cancer. For example, evidence concerning physical activity and prostate-cancer incidence was inconsistent. Therefore, physical activity should not be presented as definitively preventing every type of cancer.

14. Ultraviolet Radiation and Skin Cancer

UV radiation represents an important environmental exposure that can be influenced by lifestyle and behavior.

The Cancer Research Institute source reports that approximately **90% of skin cancers are caused by exposure to UV radiation**. UV rays are produced naturally by the sun and artificially by tanning beds and sunlamps.

UV radiation can damage DNA within skin cells and contribute to malignant transformation, particularly in areas commonly exposed to sunlight.

Therefore, reducing excessive UV exposure represents an important component of cancer-risk reduction.

15. Environmental Exposures and Air Quality

Not all lifestyle-related cancer risk results from personal choices. Environmental conditions can also contribute to cancer risk.

People may be exposed to carcinogenic substances through the **air they breathe, water they drink, workplaces, food, and their surrounding environment**.

Air quality can therefore contribute to cancer risk independently of individual behavior. Similarly, exposure to contaminated drinking water can contribute to risks such as those associated with inorganic arsenic.

This distinction is important because **cancer risk is influenced by both individual behavior and environmental circumstances**. Cancer prevention should therefore not place responsibility entirely on individuals when harmful exposures may be outside their control.

16. Cancer-Specific Lifestyle Associations

Lung Cancer

Smoking is the dominant lifestyle-related risk factor discussed in the sources. More than **80% of lung-cancer cases** were attributed to smoking in one cited analysis, with smoking associated with approximately **10–20 times greater risk**, depending on exposure history.

Other factors discussed include environmental air quality, obesity-related metabolic changes, and dietary patterns, although the evidence for these factors is less direct than the evidence linking smoking to lung cancer.

Colorectal Cancer

Colorectal cancer has particularly strong relationships with:

- Physical inactivity
- Dietary patterns
- Red and processed meat
- Excess body weight
- Alcohol consumption

High physical activity was associated with up to approximately **50% lower risk** in some studies, while one study reported a **48% lower risk** among men maintaining moderate activity.

Red and processed meat consumption is particularly relevant to colorectal cancer.

Breast Cancer

Breast cancer is influenced by several lifestyle factors, particularly:

- Alcohol consumption
- Excess body weight
- Physical inactivity
- Dietary patterns

Physical activity may influence breast-cancer risk through effects on body weight, metabolism, and hormone exposure, while alcohol consumption is associated with increased risk.

Esophageal Cancer

Tobacco and alcohol are major lifestyle-related risk factors for esophageal cancer.

Obesity is particularly important for **esophageal adenocarcinoma**, with the reviewed analysis estimating attributable fractions of approximately **29% in men and 37% in women** for overweight and obesity.

Very hot beverages have also been identified as a potential exposure associated with esophageal cancer risk in the dietary source.

Endometrial Cancer

Endometrial cancer has a strong association with excess body weight. Approximately **36%** of cases were estimated to be attributable to overweight or obesity in the cited analysis.

The reviewed dietary evidence also suggests that a high glycemic load may increase endometrial-cancer risk.

Physical activity may additionally have a protective association through effects on metabolic and hormonal regulation.

Liver Cancer

Alcohol consumption is an important lifestyle-related risk factor for liver cancer. Alcohol can contribute to chronic inflammation and cellular injury and can interact with chronic hepatitis infections to increase cancer risk.

The reviewed dietary evidence also reports an association between coffee consumption and a **lower risk of liver cancer**.

Kidney Cancer

Kidney cancer is associated with **smoking and excess body weight**. Obesity-related changes in insulin signaling, inflammation, hormones, and metabolism may contribute to this relationship.

Gallbladder Cancer

Overweight and obesity are important risk factors. The cited global analysis estimated that approximately **11% of gallbladder cancer in men and 42% in women** was attributable to overweight or obesity.

Stomach Cancer

Stomach cancer is associated with smoking, while dietary patterns and alcohol exposure may also contribute to risk.

Red and processed meat consumption has been suggested as a possible risk factor, while anti-inflammatory dietary patterns may potentially reduce stomach-cancer risk.

Pancreatic Cancer

Smoking is an established lifestyle-related risk factor for pancreatic cancer. Alcohol consumption, obesity, and dietary factors may also contribute to risk.

Oral and Pharyngeal Cancers

Tobacco and alcohol are major lifestyle-related contributors.

The reviewed global analysis reported alcohol-attributable fractions in the approximate **25–50% range among men** for oral and pharyngeal cancers.

Anti-inflammatory dietary patterns may potentially contribute to lower oral-cancer risk.

Laryngeal Cancer

Laryngeal cancer has strong associations with tobacco and alcohol. Smoking-attributable fractions exceed **50%** in the cited global analysis.

Prostate Cancer

The reviewed evidence discusses possible relationships between prostate cancer and dietary patterns, obesity, alcohol, and physical activity. However, the strength and consistency of these relationships varies.

The dietary source also reports that anti-inflammatory dietary patterns **may decrease prostate-cancer risk**, while the evidence concerning physical activity and prostate-cancer incidence remains inconsistent.

Skin Cancer

UV radiation represents the major lifestyle/environmental exposure discussed for skin cancer. Approximately **90% of skin cancers** were reported in the source as being caused by UV exposure.

UV radiation damages DNA in skin cells and can contribute to cancer development.

17. Overall Impact on Cancer and Well-being

The evidence across the reviewed sources demonstrates that lifestyle affects cancer through **multiple interconnected biological pathways**.

The major relationships can be summarized as:

Tobacco » DNA damage and mutations » cancer

Alcohol » acetaldehyde + oxidative stress + inflammation + hormonal changes » cancer

Unhealthy diet » inflammation + metabolic changes + microbiome alterations » cancer

Red/processed meat » DNA-damaging compounds + inflammation » increased cancer risk

Obesity » hormonal + metabolic + inflammatory + immune changes » cancer

Physical inactivity » metabolic + hormonal changes » increased cancer risk

UV exposure » DNA damage in skin cells » skin cancer

Environmental exposures » carcinogen exposure » cellular/DNA damage » cancer

At the same time, healthier lifestyle patterns may contribute to improved well-being through:

Balanced diet » improved nutrition + metabolic health + microbiome support

Physical activity » improved metabolic + cardiovascular + immune health

Healthy body weight » improved hormonal + metabolic regulation

Reduced tobacco/alcohol exposure » reduced carcinogenic exposure

Reduced excessive UV exposure » reduced DNA damage

The combined effects of lifestyle factors can therefore be substantial. One reviewed estimate attributes approximately **18% of cancer cases and 16% of cancer deaths** to the combined effects of excess body weight, physical inactivity, unhealthy dietary choices, and alcohol consumption.

However, these percentages come from particular analyses and populations. They should therefore be presented as **source-specific population estimates rather than as an individual's personal probability of developing cancer**.

Overall, the evidence indicates that cancer prevention is best understood as a **multifactorial process**. No single food, exercise pattern, supplement, or lifestyle behavior can completely eliminate cancer risk. Instead, cancer risk is influenced by the combined interaction of **genetic susceptibility, lifestyle, environmental exposure, metabolic health, inflammation, immune function, and aging**.

This makes lifestyle and well-being an important area for **early risk identification, prevention, and health education**, while recognizing that many cancers can still occur in individuals who maintain healthy lifestyles due to genetic, environmental, and other factors beyond their control.

Sources Used

1. [National Center for Biotechnology Information — PMC2515569](#)
2. [National Center for Biotechnology Information — Bookshelf, NBK223925](#)
3. [Cancer Research Institute — Why Do We Get Cancer? How Our Lifestyles Impact Our Risk](#)
4. [National Academy of Medical Sciences — Effect of Modifiable Lifestyle Risk Factors on the Incidence and Prevention of Cancer in Modern Society: A Review](#)
5. [American Cancer Society — Anti-inflammatory Diets and Cancer](#)
6. [American Cancer Society — How Diet and Physical Activity Impact Cancer Risk](#)
7. [GEIMS Hospital — How to Reduce Cancer Risk](#)