Let A and its RREF matrix B below as: 
$$A = \begin{bmatrix} 1 & -3 & 0 & 5 & 0 \\ 2 & -6 & 1 & 9 & 0 \\ 2 & -6 & -1 & 11 & 1 \end{bmatrix}$$
,  $B = \begin{bmatrix} 1 & -3 & 0 & 5 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$ 

- a) What is Rank(A) and what is Nullity(A)? c) What is a basis for Row(A)?
- b) What is a basis for Col(A)?

- d) What is a basis for N(A)?

**Problem Solving** - What are the terms/strategies I may need? What do I know?

## Theorems:

- 1) Rank of A is the number of pivots in the RREF matrix.
- 2) Nullity of A is the number of non-pivots in A.
- 3) A basis for the Col Space is the set of pivot columns in A corresponding to the RREF matrix.
- 4) A basis for the Row Space is the set of pivot rows in the RREF matrix of A.
- 5) A basis for the Null Space can be found by solving Ax = 0
- 6) A set of two vectors are linearly independent if the vectors are not scalar multiples of one another.

## Definition of a Basis

A linearly independent set that spans the space.

Definition of 
$$span\{v_1v_2, \dots, v_n\} = \{c_1v_1 + \dots + c_nv_n | c_i \in F\}$$

Let A and its RREF matrix B below as: 
$$A = \begin{bmatrix} 1 & -3 & 0 & 5 & 0 \\ 2 & -6 & 1 & 9 & 0 \\ 2 & -6 & -1 & 11 & 1 \end{bmatrix}$$
,  $B = \begin{bmatrix} 1 & -3 & 0 & 5 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$ 

- a) What is Rank(A) and what is Nullity(A)? c) What is a basis for Row(A)?
- b) What is a basis for Col(A)? d) What is a basis for N(A)?

Steps & Process – Try to answer the question writing in many steps to avoid small errors.

- a) The Rank of A is 3 as there are three pivots in the RREF matrix. The nullity of A is 2 as the number of non-pivots in the RREF matrix is 2.
- b) A basis for Col(A) would be  $\left\{\begin{bmatrix} 1\\2\\1\\-1\end{bmatrix},\begin{bmatrix} 0\\1\\1\end{bmatrix}\right\}$  as these are the corresponding columns in A from the pivots in the B matrix.
- c) A basis for Row(A) would be  $\{[1-3\ 0\ 5\ 0], [0\ 0\ 1-1\ 0], [0\ 0\ 0\ 1]\}$  as these are the corresponding rows in the RREF matrix. (Since row operations preserve the span).
- d) To find the basis for the Null(A), we solve Ax = 0 which reduces to Bx = 0 in this case. Since  $x_2$  and  $x_4$

are free, we get 
$$Null(A) = \left\{ \begin{bmatrix} 3s - 5t \\ s \\ t \\ t \\ 0 \end{bmatrix} \middle| s, t \in \mathbf{R} \right\} = \left\{ s \begin{bmatrix} 3 \\ 1 \\ 0 \\ 0 \end{bmatrix} + t \begin{bmatrix} -5 \\ 0 \\ 1 \\ 1 \\ 0 \end{bmatrix} \middle| s, t \in \mathbf{R} \right\} = span \left\{ \begin{bmatrix} 3 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -5 \\ 0 \\ 1 \\ 1 \\ 0 \end{bmatrix} \right\}$$

Since these two vectors are not scalar multiples of one another, we have that the basis for

Null(A) = 
$$\begin{cases} \begin{bmatrix} 3\\1\\0\\0 \end{bmatrix}, \begin{bmatrix} -5\\0\\1\\1\\0 \end{bmatrix} \end{cases}.$$

Let A and its RREF matrix B below as: 
$$A = \begin{bmatrix} 1 & -3 & 0 & 5 & 0 \\ 2 & -6 & 1 & 9 & 0 \\ 2 & -6 & -1 & 11 & 1 \end{bmatrix}$$
,  $B = \begin{bmatrix} 1 & -3 & 0 & 5 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$ 

- a) What is Rank(A) and what is Nullity(A)? c) What is a basis for Row(A)?
- b) What is a basis for Col(A)?

- d) What is a basis for N(A)?

**Solidify Understanding** – Explain why the steps makes sense by connecting to math you know.

Why do we select the columns of A for col space but the rows of B for the row space? Why does Rank + Nullity = m when a matrix is an  $A_{nm}$ ?

Why could we say that a basis for Col(A) is  $\left\{ \begin{bmatrix} 1\\0\\0 \end{bmatrix}, \begin{bmatrix} 0\\1\\0 \end{bmatrix}, \begin{bmatrix} 0\\0\\1 \end{bmatrix} \right\}$  for this question, but not in general?

For Video Please click the link below:

<u>Video</u>