CronNet-Holo Experiment: Interferometric Detection of Vacuum Resonances Predicted by Quantum Self-Verification Geometry (QSVG)

Luis Morató de Dalmases
morato.lluis@gmail.com
QSVG Research Lead

March 2026

Prepared for: CERN Research Team / Department of Theoretical Physics
Code: https://github.com/biblioteca1960/Interferometric-Detection-Simulation_Vacuum-Resonances_QSVG

Abstract

We propose a high-precision interferometric experiment in the Terahertz (THz) range to detect vacuum resonance frequencies predicted by a discrete model of spacetime—the 3,3,5 polytope network. This model, rooted in Quantum Self-Verification Geometry (QSVG), posits that spacetime emerges from a discrete geometric structure whose resonance frequencies converge to the nontrivial zeros of the Riemann zeta function. We outline the theoretical hypothesis, detailed experimental requirements, and extensive numerical validation via our open-source simulation suite, `CronNet-Holo`. A positive detection would constitute the first empirical evidence of discrete spacetime and establish a direct link between physics and number theory.

> "The most beautiful thing we can experience is the mysterious."
>
> Albert Einstein

1 Scientific Hypothesis

We challenge the paradigm of spacetime as a smooth continuum. Instead, we hypothesize it emerges from a discrete geometric structure: the regular polytope {3,3,5} in 4D, projected into 3D. This network is characterized by:

- A characteristic angular defect of 6.8° per cell.
- An effective coherence radius $R_{\text{eff}} \approx 2 \times 10^{-4}$ m.

The discrete Laplacian operator on this network yields a set of resonance frequencies. Our central hypothesis is that the ratios of these frequencies, f_n/f_1, converge asymptotically to the ratios of the imaginary parts of the nontrivial zeros of the Riemann zeta function, γ_n/γ_1.

2 Predicted Frequencies

The absolute frequency scale is set by the effective radius R_{eff}. For the nominal value $R_{\text{eff}} = 2 \times 10^{-4}$ m, the first four modes are predicted to be: *Note: Frequencies scale with* $1/R_{\text{eff}}$*; values*

Mode n	Riemann Zero γ_n	Ratio γ_n/γ_1	Predicted Frequency f_n (THz)
1	14.1347	1.0000	1.50
2	21.0220	1.4874	2.23
3	25.0108	1.7708	2.66
4	30.4248	2.1533	3.23

Table 1: Predicted resonance frequencies for $R_{\text{eff}} = 2 \times 10^{-4}$ m.

are shown for the nominal radius.

3 Experimental Requirements

To achieve the necessary sensitivity and precision, the experimental setup must meet the following stringent specifications:

Component	Specification
Cavity	Fabry-Pérot, $L = 50$ μm, Finesse $> 10^5$
Source	Photomixing, 1–4 THz tunable
Detector	He-cooled bolometer, NEP $< 10^{-15}$ W/$\sqrt{\text{Hz}}$
Temperature	4 K (liquid helium cryostat)
Shielding	μ-metal magnetic shielding (< 1 nT)

Table 2: Key components and specifications for the experiment.

4 Numerical Validation: CronNet-Holo

We have developed `CronNet-Holo`, an open-source Python simulation suite that provides robust numerical evidence for our hypothesis. The code and all results are available at the repository linked above.

4.1 Convergence Test

A convergence test on a network of 8000 nodes demonstrates the spectrum's stability.

- **Error vs Riemann zeros:** $< 0.01\%$
- **Scaling:** $\epsilon \propto N^{-0.48}$, matching the theoretical prediction $\epsilon \propto N^{-0.5}$.

4.2 Robustness to Noise

We simulated the effect of thermal and electromagnetic noise by adding random perturbations to the network's links. The results confirm topological protection of the signal.

Noise Level	Error vs Riemann	Status
0%	0.04%	✓ STABLE
5%	0.35%	✓ STABLE
10%	0.79%	✓ STABLE
15%	1.12%	△ MARGINAL
20%	1.52%	× DEGRADED

Table 3: Robustness of the frequency ratios against random noise. Signal remains stable up to 10% noise.

4.3 Experimental Simulation

A full experimental scan was simulated by convolving the theoretical spectrum with a realistic detection noise model (2%). The result predicts a signal-to-noise ratio (SNR) > 5 for all four predicted modes, allowing for direct comparison with future measurements.

5 Discovery Criteria

The experiment will be considered successful if it meets the following quantitative criteria:

1. **Spectral Match:** Four or more peaks match the predicted frequencies within $\pm 0.5\%$.
2. **Ratio Match:** The ratios f_n/f_1 match γ_n/γ_1 within an experimental error of $< 1\%$.
3. **Isotropy:** The peak pattern persists under cavity rotation $(0°, 90°, 180°)$.
4. **Control:** No peaks appear in unstructured cavities of the same material.
5. **Significance:** Each peak has a statistical significance $> 5\sigma$ (SNR ¿ 5).

6 Scientific Impact

Confirmation of this prediction would:

- Provide the first empirical evidence of discrete spacetime.
- Establish a direct, empirical link between fundamental physics and pure mathematics (number theory).
- Open a new, energy-accessible experimental window into quantum gravity.
- Offer experimental validation of the Riemann Hypothesis.

7 Why CERN?

CERN's unique and world-leading capabilities in high-precision interferometry, cryogenic systems, low-noise detection, fundamental symmetry tests, and vacuum structure experiments make it the ideal facility to conduct this verification. The expertise and infrastructure required are a perfect match for the demands of this proposal.

A Attachments

The following materials are available upon request or in the digital submission:

- **Full Technical Paper (25 pages):** Complete theoretical framework and derivations.
- **CronNet-Holo Source Code (Python):** The complete simulation suite for independent verification.
- **Simulation Results:** Plots and data for convergence, robustness, and experimental scans.
- **Detailed Experimental Protocol:** A step-by-step measurement procedure.

B Contact Information

Name:	Luis Morató de Dalmases
Role:	QSVG Research Lead
Email:	`morato.lluis@gmail.com`
Phone:	+34 681 346 877
ORCID:	0009-0009-2658-1433
Code:	`https://github.com/biblioteca1960/Interferometric-Detection-Simulation_Vacuum-Resonances_QSVG`

The complete simulation suite, technical paper, and all supplementary materials are available at the GitHub repository above.