

一、今日前沿论文精选

1. Topological Engineering of NIR-II Emissive Conjugated Oligomers With Unexpected Excited-State Manifolds for Efficient Phototheranostics and Cancer Immunotherapy.

- 发表期刊 / 影响因子:Angewandte Chemie (International ed. in English)
- 发表时间:2026-Oct-10
- 第一/通讯作者:Yuliang Yang, Yijian Gao, Ning Li, Yingpeng Wan, Huan Chen
- 核心结论:Phototheranostic agents operating in the second near-infrared (NIR-II) window hold great potential for precision cancer therapy, yet the rational regulation of excited-state processes to simultaneously optimize fluorescence and phototherapeutic performances remains a fundamental challenge. Herein, w...
- 原文DOI / 链接:10.1002/anie.2484646

2. The Effect of Clinicopathological Features and Postoperative Management on Prognosis in Male Breast Cancer.

- 发表期刊 / 影响因子:The breast journal
- 发表时间:2026
- 第一/通讯作者:Shihe Liu, Qiyang Song
- 核心结论:Male breast cancer (MBC) represents a rare type of malignant tumor, comprising less than 1% of all breast cancer cases. Therapies for MBC patients are typically adapted from breast cancer treatments in women. This study systematically performed a retrospective evaluation of the clinicopathological f...
- 原文DOI / 链接:10.1155/tbj/2806453

3. Frequency Schrödinger bridge diffusion model for fast MRI reconstruction in abdominal MR-guided online adaptive radiotherapy.

- 发表期刊 / 影响因子:Journal of applied clinical medical physics
- 发表时间:2026
- 第一/通讯作者:Zhiquan Wang, Shaobin Wang, Nan Liu, Qiu Guan, Qi Chen
- 核心结论:Online adaptive radiotherapy (ART) employs modalities like CBCT, CT, and MRI. For abdominal malignancies, MRI stands out with superior soft-tissue contrast, no ionizing radiation, and functional imaging capabilities. Yet, MRI-guided ART for such cases is challenged by long treatment time. Thus, boos...
- 原文DOI / 链接:10.1002/acm2.70831

4. Microbial cues and T cell immunity: deciphering their interactions in chronic liver disease.

- 发表期刊 / 影响因子:Experimental & molecular medicine
- 发表时间:2026-Oct-09
- 第一/通讯作者:Yinuo Liu, Baichao Yu, Qiaoyan Liu, Xiong Ma, Ruqi Tang
- 核心结论:Chronic liver disease (CLD) poses a major global health challenge, as its progression to cirrhosis and hepatocellular carcinoma significantly contributes to worldwide morbidity and mortality. Mounting evidence has identified the gut microbiota as a critical modulator of CLD progression, primarily vi...
- 原文DOI / 链接:10.1038/s12276-026-01831-y

5. Costs of two-step MASLD screening using ALT and elastography in children with overweight and obesity.

- 发表期刊 / 影响因子:Obesity research & clinical practice
- 发表时间:2026-Oct-09
- 第一/通讯作者:Anne-Sophie R Stroes, Eva Muntjewerff, Laura Draijer, Malika Chegary, Felix Kreier
- 核心结论:Metabolic dysfunction-associated steatotic liver disease (MASLD) is prevalent among children with obesity. Although screening is widely performed in pediatric obesity-related comorbidity assessment, evidence on effectiveness and costs is lacking. This study quantifies costs of MASLD screening using...
- 原文DOI / 链接:10.1016/j.orcp.2026.10.003

6. On the estimation and validity of AI time horizons—a statistical look at the METR plot

- 发表期刊 / 影响因子:arXiv Preprint
- 发表时间:2026-10-08
- 第一/通讯作者:Drew T. Nguyen, William Fithian
- 核心结论:METR' s 50% time horizon measures the human completion time of software tasks that an AI solves with 50% probability, allowing AI capabilities to be expressed in interpretable units. On 228 tasks and 26 AIs, we recompute the time horizons using splines and item-response theory to relax the assumpti...
- 原文DOI / 链接:https://arxiv.org/abs/2610.12466v1

7. From Reactive Containment to Proactive Assurance: Lessons from OpenAI, Anthropic, and Google Agent Security Incidents

- 发表期刊 / 影响因子:arXiv Preprint
- 发表时间:2026-10-08
- 第一/通讯作者:Abbas Raftari
- 核心结论:In 2026, cybersecurity evaluations involving OpenAI, Anthropic, and Google agents reached real systems outside their authorized test scope. The paths were different. OpenAI agents exploited research infrastructure, coordinated across runs, and compromised parts of Hugging Face' s production environme...
- 原文DOI / 链接:https://arxiv.org/abs/2610.12463v1

8. CSF: Contextual Safety Filtering for Motion Generators

- 发表期刊 / 影响因子:arXiv Preprint
- 发表时间:2026-10-08
- 第一/通讯作者:Lizhi Yang, Yiling Hou, Yao Tang, Junheng Li, Daniel Weng

- 核心结论:Text-conditioned motion generators produce trackable whole-body motion, but they have no notion of scene-dependent safety: the same action may target an object or a person. Existing safeguards either inspect the prompt, require labeled motion data, or enforce geometric constraints; therefore, they d…
- 原文DOI / 链接:<https://arxiv.org/abs/2610.12467v1>

二、科研公众号干货

说明:微信公众号文章检索需要微信搜一搜等公开渠道的即时访问能力,当前自动检索模块正在优化中,建议通过以下方式自行获取:- 关注「肿瘤放疗」、「生信技能树」、「医学僧的科研日记」等公众号 - 使用微信搜一搜搜索关键词:肿瘤放疗 生信分析 - 本板块将在后续版本中接入 WeChat 公开内容 API

三、每日高分经典精读

Stereotactic Body Radiation Therapy for Hepatocellular Carcinoma: Current Trends and Future Directions

- 发表期刊 / 影响因子 / 总引用量:International Journal of Radiation Oncology, Biology, Physics / IF=7.0 / 引用量:200+
- 研究背景与里程碑意义:SBRT has emerged as a definitive local treatment for HCC, but optimal dosing, patient selection, and integration with systemic therapies remain active areas of investigation.
- 核心实验设计与方法:Comprehensive review of clinical trials and retrospective studies of SBRT for HCC, including dose-escalation studies, toxicity analyses, and combination strategies.
- 关键结论与行业影响:SBRT achieves local control rates of 80-95% for early-stage HCC with low toxicity. Challenges include BCLC stage-specific outcomes, liver function preservation, and optimal sequencing with immune checkpoint inhibitors.
- 创新点:Systematic framework for SBRT dose selection based on tumor size, location, and liver function; proposed risk-adapted fractionation schedules.
- 原文DOI / 链接:<https://doi.org/10.1016/j.ijrobp.2023.01.045>

四、当日总结

今日共精选8篇最新研究+1篇经典精读。关注方向:Topological Engineering of NIR-II Emissive Conjugated Oligomers With Unexpected Excited-State Manifolds for Efficient Phototheranostics and Cancer Immunotherapy.