

一、今日前沿论文精选

1. Differences of Community-Level Vulnerability Associations With Early- and Later-Onset Nasopharyngeal Cancer Prognosis in the United States.

- 发表期刊 / 影响因子:Otolaryngology–head and neck surgery : official journal of American Academy of Otolaryngology-Head and Neck Surgery
- 发表时间:2026-Sep-11
- 第一/通讯作者:Eric Pan, Hari Patel, David Fei-Zhang, Jill D` Souza, Anthony Sheyn
- 核心结论:To assess age-related differences of community, social vulnerability associations with care and prognosis among those with nasopharyngeal carcinomas. Retrospective Cohort. United States. This retrospective cohort study on 12,480 NPC-patients from 1975 to 2017 linked with 15 county-level vulnerabil...
- 原文DOI / 链接:10.1002/ohn.70430

2. Objective Swallowing Outcomes After Sparing Contralateral Low-Neck Elective Radiotherapy in Oropharyngeal Carcinoma: Single-Institution Clinical Experience.

- 发表期刊 / 影响因子:Otolaryngology–head and neck surgery : official journal of American Academy of Otolaryngology-Head and Neck Surgery
- 发表时间:2026-Sep-11
- 第一/通讯作者:Rex H Lee, Cara Evans, Sarah Honjo, Yue Ma, Alon Witztum
- 核心结论:Dysphagia is a major toxicity of chemoradiation (CRT) for oropharyngeal squamous cell carcinoma (OPSCC). Elective upper-neck irradiation (UNI), omitting the low neck, may reduce esophageal inlet muscle (EIM) dose and improve swallowing outcomes compared to elective whole-neck irradiation (WNI). We...
- 原文DOI / 链接:10.1002/ohn.70429

3. A systematic review of international/national guidelines for the management of nasopharyngeal carcinoma: Convergence and divergence of recommendations.

- 发表期刊 / 影响因子:Cancer
- 发表时间:2026-Sep-15
- 第一/通讯作者:Qin Liu, Handoko, Ji-Shi Li, Qiao-Juan Guo, Fernando Lopez
- 核心结论:Increasing numbers of clinical practice guidelines have been published by international/national groups for nasopharyngeal carcinoma (NPC), providing valuable references for clinicians in making evidence-based decisions on treatment. However, there are substantial discrepancies in various recommenda...
- 原文DOI / 链接:10.1002/cncr.70604

4. Body mass index modifies the association between occupational noise exposure duration and ultrasound-detected hepatic steatosis: A cross-sectional study among male factory workers.

- 发表期刊 / 影响因子:PloS one
- 发表时间:2026
- 第一/通讯作者:Dandan Liu, Huixia Ji, Ye Chen, Tongzhou Zhou
- 核心结论:Metabolic dysfunction-associated steatotic liver disease (MASLD) is a multisystem metabolic disorder. The impact of occupational noise exposure on metabolic processes through non-auditory pathways is also closely associated with hepatic steatosis in addition to obesity. Most studies examine noise ex...
- 原文DOI / 链接:10.1371/journal.pone.0358275

5. Metabolic dysfunction-associated steatotic liver disease predicts graft failure in kidney transplant recipients.

- 发表期刊 / 影响因子:Clinical kidney journal
- 发表时间:2026
- 第一/通讯作者:Kyungho Lee, Yebin Park, Hojin Jeon, Junseok Jeon, Jung Eun Lee
- 核心结论:Metabolic dysfunction-associated steatotic liver disease (MASLD) has been established to classify steatotic liver disease more accurately. We investigated the association between MASLD and long-term kidney transplant outcomes. Kidney transplant recipients who underwent national health screening prog...
- 原文DOI / 链接:10.1093/ckj/sfag294

二、科研公众号干货

说明:微信公众号文章检索需要微信搜一搜等公开渠道的即时访问能力,当前自动检索模块正在优化中,建议通过以下方式自行获取:- 关注「肿瘤放疗」、「生信技能树」、「医学僧的科研日记」等公众号 - 使用微信搜一搜搜索关键词:肿瘤放疗 生信分析 - 本板块将在后续版本中接入 WeChat 公开内容 API

三、每日高分经典精读

Metabolic dysfunction-associated steatotic liver disease (MASLD): a growing global health priority and a real challenge for the internist.

- 发表期刊 / 影响因子 / 总引用量:Internal and emergency medicine / 2026
- 研究背景与里程碑意义:MASLD has emerged as the most prevalent chronic liver disease worldwide, reflecting the global rise in metabolic disorders such as obesity and diabetes.
- 核心实验设计与方法:Narrative review of epidemiology, pathophysiology, diagnostic criteria, and management strategies for MASLD.
- 关键结论与行业影响:Internists play a central role in early detection and multidisciplinary management of MASLD to prevent progression to cirrhosis and hepatocellular carcinoma.
- 创新点:Pragmatic clinical framework for internists to screen, stage, and manage MASLD in routine practice.
- 原文DOI / 链接:https://doi.org/10.1007/s11739-026-04461-4

四、当日总结

