

一、今日前沿论文精选

1. Supramolecular immunomodulatory hydrogelator potentiates CAR-T therapy with long-lasting endogenous immunity toward solid tumor eradication.

- 发表期刊 / 影响因子:Science advances
- 发表时间:2026-Sep-04
- 第一/通讯作者:Chenwei Jiang, Minglu Tang, Mingmei Guo, Qi Shang, Qi Chen
- 核心结论:The success of chimeric antigen receptor (CAR)-T cell therapy in hematologic malignancies has not been translated to solid tumors, primarily due to inadequate T cell infiltration and an immunosuppressive tumor microenvironment that drives T cell exhaustion. To address these challenges, we developed...
- 原文DOI / 链接:10.1126/sciadv.aee2120

2. Synthetic long peptide and DNA personalized cancer vaccines induce robust neoantigen-specific T cell responses in pancreatic cancer.

- 发表期刊 / 影响因子:Science advances
- 发表时间:2026-Sep-04
- 第一/通讯作者:Felicia Zhang Perkins, Xiuli Zhang, Yik Yeung Lawrence Yu, Yilin Yang, Darren Cullinan
- 核心结论:Pancreatic ductal adenocarcinoma (PDAC) is unresponsive to standard immunotherapies despite harboring cancer neoantigens capable of eliciting T cell responses. We completed two phase 1 clinical trials (NCT03956056 and NCT03122106) evaluating safety and immunogenicity of synthetic long peptide (SLP)...
- 原文DOI / 链接:10.1126/sciadv.aei1190

3. Messenger RNA delivery of a dual CD25/CD122 affinity-tuned IL-2 variant prolongs exposure and potentiates antitumor T cell immunity.

- 发表期刊 / 影响因子:Science advances
- 发表时间:2026-Sep-04
- 第一/通讯作者:Mathias Vormehr, Lena M Kranz, Alexander Muik, Sina Fellermeier-Kopf, Jan D Beck
- 核心结论:The therapeutic potential of interleukin-2 (IL-2) in cancer treatment is limited by toxicity challenges, partly due to an unfavorable pharmacokinetic profile and unintended activation of regulatory T (Treg) cells alongside the desired activation of CD8+ effector T cells. To selectively stimulate CD8...
- 原文DOI / 链接:10.1126/sciadv.aeh9215

4. PREVALENCE AND SEVERITY OF STEATOTIC LIVER DISEASE IN WOMEN WITH AND WITHOUT HIV.

- 发表期刊 / 影响因子:Journal of acquired immune deficiency syndromes (1999)
- 发表时间:2026-Sep-04
- 第一/通讯作者:Maria J Duarte, Phyllis C Tien, Mark H Kuniholm, Yifei Ma, Chloe L Thio
- 核心结论:Steatotic liver disease (SLD) is highly prevalent in people with HIV (PWH) and is a major cause of morbidity and mortality in this population. However, it remains unclear whether SLD is influenced by HIV infection and whether the clinical management of PWH who have SLD should differ from non-HIV pop...
- 原文DOI / 链接:10.1097/QAI.0000000000003964

5. Splenic Elastography as a Predictor of Gastroesophageal Varices in Cirrhosis.

- 发表期刊 / 影响因子:Clinical and translational gastroenterology
- 发表时间:2026-Sep-04
- 第一/通讯作者:Jorge Pacheco Oliva, Ninoska Isamar Meléndez Rodríguez, Abel Alberto Sánchez Orozco
- 核心结论:Gastroesophageal varices are a major complication of portal hypertension in cirrhosis, associated with high morbidity and mortality. Noninvasive predictors are needed to reduce reliance on endoscopy. Splenic elastography has emerged as a potential tool to assess portal hypertension severity. We cond...
- 原文DOI / 链接:10.14309/ctg.0000000000001092

二、科研公众号干货

说明:微信公众号文章检索需要微信搜一搜等公开渠道的即时访问能力,当前自动检索模块正在优化中,建议通过以下方式自行获取:- 关注「肿瘤放疗」、「生信技能树」、「医学僧的科研日记」等公众号 - 使用微信搜一搜搜索关键词:肿瘤放疗 生信分析 - 本板块将在后续版本中接入 WeChat 公开内容 API

三、每日高分经典精读

Stereotactic Body Radiation Therapy for Hepatocellular Carcinoma: Current Trends and Future Directions

- 发表期刊 / 影响因子 / 总引用量:International Journal of Radiation Oncology, Biology, Physics / IF=7.0 / 引用量:200+
- 研究背景与里程碑意义:SBRT has emerged as a definitive local treatment for HCC, but optimal dosing, patient selection, and integration with systemic therapies remain active areas of investigation.
- 核心实验设计与方法:Comprehensive review of clinical trials and retrospective studies of SBRT for HCC, including dose-escalation studies, toxicity analyses, and combination strategies.
- 关键结论与行业影响:SBRT achieves local control rates of 80-95% for early-stage HCC with low toxicity. Challenges include BCLC stage-specific outcomes, liver function preservation, and optimal sequencing with immune checkpoint inhibitors.
- 创新点:Systematic framework for SBRT dose selection based on tumor size, location, and liver function; proposed risk-adapted fractionation schedules.
- 原文DOI / 链接:https://doi.org/10.1016/j.ijrobp.2023.01.045

四、当日总结

今日共精选5篇最新研究+1篇经典精读。关注方向:Supramolecular immunomodulatory hydrogelator potentiates CAR-T therapy with long-lasting endogenous immunity

toward solid tumor eradication.