每日医学科研精读日报 | 2026年11月09日

一、今日前沿论文精选

1. Salvage High-Dose-Rate Brachytherapy for Localized Prostate Cancer Recurrence: A Comparative Analysis of 2 Fractionation Schedules-Single-Institution Study of 223 Patients.

- 发表期刊 / 影响因子:Advances in radiation oncology
- 发表时间:2027
- 第一/通讯作者:Magdalena Stankiewicz, Agata Wilk, Marcela Krzempek, Sylwia Kellas-Ślęczka, Marta Szlag
- 核心结论:Salvage high-dose-rate brachytherapy (sHDR-BT) is an effective option for radiorecurrent prostate cancer. This study aimed to compare the efficacy and toxicity of two sHDR-BT fractionation schedules and identify factors associated with treatment outcomes. We retrospectively analyzed 223 patients tre…
- 原文DOI / 链接:10.1016/j.adro.2026.102210

2. Metastasis-directed therapy in addition to definitive locoregional radiotherapy for De novo metastatic nasopharyngeal carcinoma: a retrospective cohort study.

- 发表期刊 / 影响因子:Clinical and translational radiation oncology
- 发表时间:2027
- 第一/通讯作者:Sarbani Ghosh Laskar, Anuj Kumar, Rahat Malhotra, Shwetabh Sinha, Samarpita Mohanty
- 核心结论:De novo metastatic nasopharyngeal carcinoma (mNPC) carries a poor prognosis with systemic therapy alone. Locoregional radiotherapy (LRRT) improves survival in chemotherapy-responsive patients, but the incremental role of metastasis-directed therapy (MDT) remains undefined. This retrospective cohort…
- 原文DOI / 链接:10.1016/j.ctro.2026.101338

3. Osteoradionecrosis in patients with head and neck cancer after reirradiation - a systematic review and meta-analysis.

- 发表期刊 / 影响因子:Clinical and translational radiation oncology
- 发表时间:2027
- 第一/通讯作者:Annemarie Fritz, Nils Gleim, Anita Kloss-Brandstätter, Thomas Kuhnt, Nils H Nicolay
- 核心结论:To determine the observed incidence and risk factors of osteoradionecrosis (ORN) following reirradiation for recurrent or second primary head and neck cancer. A systematic review with meta-analysis was conducted according to the PRISMA guidelines, with the study protocol prospectively registered wit…
- 原文DOI / 链接:10.1016/j.ctro.2026.101342

4. Health-care burden of metabolic dysfunction-associated steatotic liver disease in European tertiary-care outpatient clinics (CONSTANS): a prospective multicentre cohort study.

- 发表期刊 / 影响因子:The Lancet regional health. Europe
- 发表时间:2026
- 第一/通讯作者:Vlad Ratziu, Jean-François Dufour, Maharajah Ponnaiah, Raluca Pais, Maximilian Thormann
- 核心结论:Despite modeling-based epidemiological studies of MASLD, precise estimates of the prevalence and health-care resource utilization in outpatient clinical practice are unknown. We assessed, in a real-life setting, the prevalence of suspected/confirmed MASLD and described health-care resource utilizati…
- 原文DOI / 链接:10.1016/j.lanepe.2026.101893

5. Author Reply to Letter to the Editor Regarding ‘’ Beyond Awareness: A Qualitative Assessment of Barriers and Facilitators to Implementing Metabolic dysfunction-Associated Steatotic Liver Disease (MASLD) Care Pathways in Primary Care’’.

- 发表期刊 / 影响因子:Gastro hep advances
- 发表时间:2026
- 第一/通讯作者:Leonard Niels D Broekman, Stan Driessen, Adriaan Onno G Holleboom
- 核心结论:参见原文摘要
- 原文DOI / 链接:10.1016/j.gastha.2026.101120

6. Phantom Consensus: Measuring Social Reasoning in Multi-Agent LLM Decision-Making

- 发表期刊 / 影响因子:arXiv Preprint
- 发表时间:2026-11-06
- 第一/通讯作者:Disha Sheshanarayana, Rajat Subhra Pal, Manjira Sinha, Tirthankar Dasgupta
- 核心结论:Multi-agent LLM systems increasingly aggregate judgments across model instances, but reasoning about other agents can itself introduce failure. We study a setting in which LLMs shift their public decisions after observing ambiguous social signals while reporting inaccurate estimates of what other ag…
- 原文DOI / 链接:https://arxiv.org/abs/2611.08971v1

7. PhysioWave-T: Token-Efficient Wavelet Foundation Models for Biosignals

- 发表期刊 / 影响因子:arXiv Preprint
- 发表时间:2026-11-06
- 第一/通讯作者:Yanlong Chen, Simone Benatti, Luca Benini, Cuntai Guan, Yawei Li
- 核心结论:We present PhysioWave-T, a token-efficient wavelet foundation model family for EEG, ECG, and sEMG. Dynamic scale folding recombines J+1 wavelet subbands prior to tokenization, compressing sequence length from (J+1)CS to CS tokens for J levels, C channels, and S temporal patches. Route-specific front…
- 原文DOI / 链接:https://arxiv.org/abs/2611.08969v1

8. Energy-Based Reasoners: Your Recurrent Reasoner Should Be an Energy-Based Model

- 发表期刊 / 影响因子:arXiv Preprint

- 发表时间:2026-11-06
- 第一/通讯作者:William Xingxu Chen, Alexi Gladstone, Zeyu Michael Li, Heng Ji, Yilun Du
- 核心结论:Looping is a powerful inductive bias for reasoning. By repeatedly iterating a recurrent block, looped models can scale their effective depth and enable multi-step reasoning in latent space. However, existing looped models are not inherently guided by an explicit measure of progress, making it diffic…
- 原文DOI / 链接:https://arxiv.org/abs/2611.08977v1

二、科研公众号干货

说明:微信公众号文章检索需要微信搜一搜等公开渠道的即时访问能力,当前自动检索模块正在优化中,建议通过以下方式自行获取: - 关注「肿瘤放疗」、「生信技能树」、「医学僧的科研日记」等公众号 - 使用微信搜一搜搜索关键词:肿瘤放疗 生信分析 - 本板块将在后续版本中接入 WeChat 公开内容 API

三、每日高分经典精读

Metabolic dysfunction-associated steatotic liver disease (MASLD): a growing global health priority and a real challenge for the internist.

- 发表期刊 / 影响因子 / 总引用量:Internal and emergency medicine / 2026
- 研究背景与里程碑意义:MASLD has emerged as the most prevalent chronic liver disease worldwide, reflecting the global rise in metabolic disorders such as obesity and diabetes.
- 核心实验设计与方法:Narrative review of epidemiology, pathophysiology, diagnostic criteria, and management strategies for MASLD.
- 关键结论与行业影响:Internists play a central role in early detection and multidisciplinary management of MASLD to prevent progression to cirrhosis and hepatocellular carcinoma.
- 创新点:Pragmatic clinical framework for internists to screen, stage, and manage MASLD in routine practice.
- 原文DOI / 链接:https://doi.org/10.1007/s11739-026-04461-4

四、当日总结

今日共精选8篇最新研究+1篇经典精读。关注方向:Salvage High-Dose-Rate Brachytherapy for Localized Prostate Cancer Recurrence: A Comparative Analysis of 2 Fractionation Schedules-Single-Institution Study of 223 Patients.

日报自动生成于 2026年11月09日 | 数据来源:PubMed、arXiv | 下一期:明日7:00