每日医学科研精读日报 | 2026年10月30日

一、今日前沿论文精选

1. Anti-PD-L1 glycoengineering enhances tumor killing through coordinated interactions between PD-L1⁺ DCs and CD16A⁺ cytotoxic lymphocytes.

- 发表期刊 / 影响因子:Cancer research communications
- 发表时间:2026-Oct-29
- 第一/通讯作者:Anne Louise Blomberg, Weihua Tian, Judit Pina Agullet, Natasha Helleberg Madsen, Katrine Grove Linde
- 核心结论:Blockade of the PD-1/PD-L1 axis has transformed cancer immunotherapy, yet many patients fail to achieve durable responses. Although Fc glycoengineering by defucosylation is well established to enhance NK cell-mediated antibody-dependent cellular cytotoxicity through increased FcγRIIIa binding, its e…
- 原文DOI / 链接:10.1158/2767-9764.CRC-26-0696

2. Indazole Derivatives as Emerging Scaffolds for Anti-Breast Cancer Therapy: Synthesis, Structure-Activity Relationship, and Therapeutic Potential.

- 发表期刊 / 影响因子:Archiv der Pharmazie
- 发表时间:2026
- 第一/通讯作者:Soubhik Tanti, Akhilesh Gangwar, Vikas Jaitak
- 核心结论:Breast cancer (BC) is the second most common cause of death in women. Current therapies like hormonal therapy, surgery, radiotherapy, and chemotherapy become ineffective due to severe side effects and multidrug resistance. Therefore, there is an urgent need for the development of more effective and…
- 原文DOI / 链接:10.1002/ardp.70374

3. Phase 1 study of OX40 agonist ABBV-368 as monotherapy or in combination with budigalimab in advanced solid tumors.

- 发表期刊 / 影响因子:Future oncology (London, England)
- 发表时间:2026-Oct-29
- 第一/通讯作者:Marc Oliva, John Powderly, Christophe Le Tourneau, Amani Makkouk, Philippe A Cassier
- 核心结论:This first-in-human phase 1 trial evaluated the OX40 agonist antibody ABBV-368 as monotherapy and with the anti-programmed cell death protein (PD)-1 budigalimab in advanced solid tumors. In dose escalation, patients with solid tumors received ABBV-368 monotherapy. In dose expansion, patients with re…
- 原文DOI / 链接:10.1080/14796694.2026.2736946

4. Nanoplastics as Hepatic Toxicant Vectors: Interactions with Environmental Chemicals, Organelle Dysfunction, and Regulated Cell Death.

- 发表期刊 / 影响因子:Toxicology mechanisms and methods
- 发表时间:2026-Oct-29
- 第一/通讯作者:Amr Ali Mohamed Abdelgawwad El-Sehrawy, Safa Al-Kayyat, Djaloliddin Mansurov, Dhara N Patel, Priya Priyadarshini Nayak
- 核心结论:The environmental accumulation of nanoplastics (NPs) raises concerns regarding their potential effects on hepatic function. Experimental studies suggest that NP exposure can be related with altered mitochondrial electron transport chain activity, reduced ATP availability, changes in Keap1/Nrf2-media…
- 原文DOI / 链接:10.1080/15376516.2026.2755752

5. Olvanil Reduces Subcutaneous Fat Accumulation and Lipogenic Gene Expression, Without Alleviating Hepatic Oxidative Stress or Histopathological Alterations in Mice with Metabolic Dysfunction-Associated Steatotic Liver Disease.

- 发表期刊 / 影响因子:Journal of medicinal food
- 发表时间:2026-Oct-29
- 第一/通讯作者:Elda Cristina Villaseñor-Tapia, David Alejandro Curiel-Pedraza, Edgar Rubén Mendieta-Condado, Jorge Alberto Rodríguez-González, Alejandro Arturo Canales-Aguirre
- 核心结论:Metabolic dysfunction-associated steatotic liver disease (MASLD) is characterized by excess intrahepatic lipid accumulation due to lipid overflow and synthesis, as well as impaired lipid oxidation. Olvanil (OLV) is a nonpungent capsaicin analog that has previously been shown to have anti-obesity eff…
- 原文DOI / 链接:10.1177/1096620X261499871

6. Autonomous Quantum Measurement with Agentic AI

- 发表期刊 / 影响因子:arXiv Preprint
- 发表时间:2026-10-28
- 第一/通讯作者:Eunbi Rha, Abhijatmedhi Chotrattanapituk, Mouyang Cheng, Chung-tao Chou, Shenyong Ye
- 核心结论:Quantum materials exhibit diverse electronic and functional phenomena, but their characterization requires extensive experimental expertise, iterative decision-making and accumulated laboratory knowledge that is difficult to document and transfer. Here we introduce Quantum Agent (QUANTA), an agentic…
- 原文DOI / 链接:https://arxiv.org/abs/2610.38850v1

7. Discovering Jailbreaks via Curriculum of Models

- 发表期刊 / 影响因子:arXiv Preprint
- 发表时间:2026-10-28
- 第一/通讯作者:Michał Burzyński, Noah T. Rossi, Shomit Basu, Andy Arditi
- 核心结论:Automated jailbreak discovery aims to find prompts that elicit prohibited behaviors from safety-trained language models, helping developers identify vulnerabilities before deployment. One class of approaches uses reinforcement learning to train an investigator model to generate prompts that elicit a…
- 原文DOI / 链接:https://arxiv.org/abs/2610.38833v1

8. Discovery of Diverse and Realistic Financial Tail-Risk Using Generative Flow Networks

- 发表期刊 / 影响因子:arXiv Preprint
- 发表时间:2026-10-28
- 第一/通讯作者:Adir Serruya, Gilad Katz
- 核心结论:Accurate modeling of rare, high-impact financial events is essential for robust risk management. Sampling-based techniques such as Sequential Markov Chain Monte Carlo (SMCMC) generate diverse scenarios but inefficiently explore high-risk regions, while Deep Reinforcement Learning (DRL) converges on…
- 原文DOI / 链接:https://arxiv.org/abs/2610.38838v1

二、科研公众号干货

说明:微信公众号文章检索需要微信搜一搜等公开渠道的即时访问能力,当前自动检索模块正在优化中,建议通过以下方式自行获取: - 关注「肿瘤放疗」、「生信技能树」、「医学僧的科研日记」等公众号 - 使用微信搜一搜搜索关键词:肿瘤放疗 生信分析 - 本板块将在后续版本中接入 WeChat 公开内容 API

三、每日高分经典精读

Metabolic dysfunction-associated steatotic liver disease (MASLD): a growing global health priority and a real challenge for the internist.

- 发表期刊 / 影响因子 / 总引用量:Internal and emergency medicine / 2026
- 研究背景与里程碑意义:MASLD has emerged as the most prevalent chronic liver disease worldwide, reflecting the global rise in metabolic disorders such as obesity and diabetes.
- 核心实验设计与方法:Narrative review of epidemiology, pathophysiology, diagnostic criteria, and management strategies for MASLD.
- 关键结论与行业影响:Internists play a central role in early detection and multidisciplinary management of MASLD to prevent progression to cirrhosis and hepatocellular carcinoma.
- 创新点:Pragmatic clinical framework for internists to screen, stage, and manage MASLD in routine practice.
- 原文DOI / 链接:https://doi.org/10.1007/s11739-026-04461-4

四、当日总结

今日共精选8篇最新研究+1篇经典精读。关注方向:Anti-PD-L1 glycoengineering enhances tumor killing through coordinated interactions between PD-L1^{+} DCs and CD16A^{+} cytotoxic lymphocytes.

日报自动生成于 2026年10月30日 | 数据来源:PubMed、arXiv | 下一期:明日7:00