

一、今日前沿论文精选

1. Berberine induces PD-L1 degradation via the autophagy-lysosome pathway through the PI3K-Akt pathway and enhances immunogenic cell death in triple-negative breast cancer.

- 发表期刊 / 影响因子:Frontiers in immunology
- 发表时间:2026
- 第一/通讯作者:Yuqin Wei, Qiudan Huang, Xingwen Wei, Ruilan Mo, Wei Zhao
- 核心结论:This study aimed to investigate the regulatory effect and molecular mechanism of Berberine (BBR) on programmed death-ligand 1 (PD-L1) expression in Triple-Negative Breast Cancer (TNBC) cells. In vitro cellular experiments (using MDA-MB-231 and BT-549 cell lines) and mouse models (Balb/c-nu and Balb/...
- 原文DOI / 链接:10.3389/fimmu.2026.1797840

2. Neoadjuvant therapy combined with immunotherapy for anal sphincter preservation rate, clinical efficacy, and safety in rectal cancer patients: a meta-analysis.

- 发表期刊 / 影响因子:Frontiers in oncology
- 发表时间:2026
- 第一/通讯作者:Sijie Chen, Yuansha Ge, Xue He, Wenkang Zhang, Jie Li
- 核心结论:Neoadjuvant therapy combined with immunotherapy has emerged as a promising strategy for rectal cancer (RC), but its efficacy in anal sphincter preservation, clinical outcomes, and safety profile remains to be systematically validated. This meta-analysis aims to comprehensively evaluate the anal sphi...
- 原文DOI / 链接:10.3389/fonc.2026.1866026

3. Overcoming Resistance to Immune Checkpoint Blockade: Mechanistic Insights and Emerging Therapeutic Directions.

- 发表期刊 / 影响因子:Iranian journal of immunology : IJI
- 发表时间:2026-Aug-22
- 第一/通讯作者:Yuyu Xu, Huiling Shen
- 核心结论:参见原文摘要
- 原文DOI / 链接:10.22034/iji.2026.111387.3205

4. Non-lean versus lean MASLD: associations with advanced cardiovascular-kidney-metabolic syndrome in adults with type 2 diabetes mellitus, a cross-sectional study.

- 发表期刊 / 影响因子:Frontiers in endocrinology
- 发表时间:2026
- 第一/通讯作者:Jun-Wei Wang, Le Jiang, Lin-Lin Gao, Jie Ren, Shi-Yu Zhao
- 核心结论:This study seeks to explore the impact of lean and non-lean phenotypes of metabolic dysfunction-associated steatotic liver disease (MASLD) on the prevalence of advanced cardiovascular-kidney-metabolic (CKM) syndrome stage in individuals with type 2 diabetes mellitus (T2DM). Based on a BMI cutoff of...
- 原文DOI / 链接:10.3389/fendo.2026.1910398

5. Partial depletion of plasminogen activator inhibitor-1 decreases subcutaneous fat cell hypertrophy and liver cholesterol in high-fat-fed female mice.

- 发表期刊 / 影响因子:FEBS letters
- 发表时间:2026-Aug-21
- 第一/通讯作者:Claudia E Ramirez Bustamante, Robert Sharp, Pradip K Saha, Elizabeth S Anaya, Elizabeth L Lieu
- 核心结论:Higher levels of plasminogen activator inhibitor-1 (PAI-1) circulate in the blood of people with obesity and predict hyperlipidemia and metabolic dysfunction-associated steatotic liver disease in studies predominantly conducted in women. To understand the roles of PAI-1 in women with obesity, we gen...
- 原文DOI / 链接:10.1002/1873-3468.70439

6. G-CARL: Grounded Checklist-Aligned Reward Learning for Patient-Oriented Medical Report Interpretation

- 发表期刊 / 影响因子:arXiv Preprint
- 发表时间:2026-08-20
- 第一/通讯作者:Shiao Xie, Siyu Chen, Jianwei Lv, Bo Yuan, Yujin Wang
- 核心结论:Personalized interpretation of medical reports has emerged as an increasingly important need among patients. Addressing this need requires both evidence-grounded medical factuality and context-dependent patient communication, yet existing medical vision-language tasks do not adequately capture these...
- 原文DOI / 链接:https://arxiv.org/abs/2608.20331v1

7. An Agentic Approach for Active Data Collection, Travel Behavior Modeling, and Weather-Sensitive Demand Prediction

- 发表期刊 / 影响因子:arXiv Preprint
- 发表时间:2026-08-20
- 第一/通讯作者:Narges Ahmadi, Yubo Jiao, Jônatas Augusto Manzolli, Jiangbo Yu, Luis Miranda-Moreno
- 核心结论:Travel behavior research increasingly combines digital data collection with predictive modeling, yet these stages are often developed and evaluated separately. This study proposes a three-agent workflow integrating conversational data collection, structured data processing, and behavioral prediction...
- 原文DOI / 链接:https://arxiv.org/abs/2608.20320v1

8. Information on trajectories: martingales and random times

- 发表期刊 / 影响因子:arXiv Preprint
- 发表时间:2026-08-20
- 第一/通讯作者:Akshay Balsubramani
- 核心结论:Accounting for information flow on the path space of trajectories of a nonnegative martingale yields exact variational identities for it, even at arbitrary random times. This recovers the widely used classical concentration inequalities, from Ville to PAC-Bayes, and measures what each one discards...
- 原文DOI / 链接:<https://arxiv.org/abs/2608.20337v1>

二、科研公众号干货

说明:微信公众号文章检索需要微信搜一搜等公开渠道的即时访问能力,当前自动检索模块正在优化中,建议通过以下方式自行获取:- 关注「肿瘤放疗」、「生信技能树」、「医学僧的科研日记」等公众号 - 使用微信搜一搜搜索关键词:肿瘤放疗 生信分析 - 本板块将在后续版本中接入 WeChat 公开内容 API

三、每日高分经典精读

Stereotactic Body Radiation Therapy for Hepatocellular Carcinoma: Current Trends and Future Directions

- 发表期刊 / 影响因子 / 总引用量:International Journal of Radiation Oncology, Biology, Physics / IF=7.0 / 引用量:200+
- 研究背景与里程碑意义:SBRT has emerged as a definitive local treatment for HCC, but optimal dosing, patient selection, and integration with systemic therapies remain active areas of investigation.
- 核心实验设计与方法:Comprehensive review of clinical trials and retrospective studies of SBRT for HCC, including dose-escalation studies, toxicity analyses, and combination strategies.
- 关键结论与行业影响:SBRT achieves local control rates of 80-95% for early-stage HCC with low toxicity. Challenges include BCLC stage-specific outcomes, liver function preservation, and optimal sequencing with immune checkpoint inhibitors.
- 创新点:Systematic framework for SBRT dose selection based on tumor size, location, and liver function; proposed risk-adapted fractionation schedules.
- 原文DOI / 链接:<https://doi.org/10.1016/j.ijrobp.2023.01.045>

四、当日总结

今日共精选8篇最新研究+1篇经典精读。关注方向:Berberine induces PD-L1 degradation via the autophagy-lysosome pathway through the PI3K-Akt pathway and enhances immunogenic cell death in triple-negative breast cancer.