每日医学科研精读日报 | 2026年11月06日

一、今日前沿论文精选

1. Evaluation of the Clinical Utility of Circulating Tumor DNA in the Management of Patients With Nonmetastatic Head and Neck Cancer: Protocol for a Prospective Bicentric Study (CIRVAD).

- 发表期刊 / 影响因子:JMIR research protocols
- 发表时间:2026-Nov-05
- 第一/通讯作者:Romina Mastronicola, Maëlys Bresson, Giacomo Gravante, Sophie Cortese, Emilie Beulque
- 核心结论:Human papillomavirus (HPV)-negative head and neck cancers, often linked to tobacco and alcohol exposure, have a poor prognosis and high recurrence rates. Circulating tumor DNA (ctDNA) is a promising biomarker for detecting minimal residual disease and monitoring treatment response. Recent studies sh…
- 原文DOI / 链接:10.2196/89973

2. Practical Recommendations for Implementing Remote Patient Monitoring for Patients Receiving Bispecific Antibodies for Multiple Myeloma.

- 发表期刊 / 影响因子:JMIR cancer
- 发表时间:2026-Nov-05
- 第一/通讯作者:Cindy Varga, Eric Schaefer
- 核心结论:Bispecific antibodies are novel immunotherapies approved for the treatment of patients with relapsed or refractory multiple myeloma (MM). While these therapies induce deep and durable responses, they are associated with adverse events, such as cytokine release syndrome and immune effector cell-assoc…
- 原文DOI / 链接:10.2196/95949

3. HPV Vaccination Among Medicaid-Insured 9-12-year-Olds in Texas.

- 发表期刊 / 影响因子:Cancer control : journal of the Moffitt Cancer Center
- 发表时间:2026
- 第一/通讯作者:Erika L Thompson, Yong Shan, Jacqueline M Hirth, Miranda Cano, Yong-Fang Kuo
- 核心结论:IntroductionNational organizations support starting HPV vaccination at age 9 to increase initiation and completion of the vaccine series. This study aimed to assess the patterns of HPV vaccination initiation for Texas children who are enrolled in Medicaid.MethodsUsing a retrospective cohort design,…
- 原文DOI / 链接:10.1177/10732748261505753

4. Comparing attenuation parameter and liver stiffness measurement using Hepatus and Fibroscan for the evaluation of steatosis and fibrosis in patients with chronic liver disease: A prospective study.

- 发表期刊 / 影响因子:PloS one
- 发表时间:2026
- 第一/通讯作者:Cain-En Lim, Ram Prasad Sinnanaidu, Xin Tong Ng, Jin Ying Teh, Farahani Khamis
- 核心结论:There are limited studies comparing Hepatus (HP), a novel visual transient elastography device with Fibroscan (FS), one of the most studied and validated non-invasive tests in the field of hepatology. HP and FS examinations were performed on patients with chronic liver disease by two operators, heal…
- 原文DOI / 链接:10.1371/journal.pone.0359831

5. Lean MAFLD as an independent risk factor for carotid plaque: The value of FIB-4 index in subgroup analysis.

- 发表期刊 / 影响因子:PloS one
- 发表时间:2026
- 第一/通讯作者:Guanqun Chao, Ailing Lin, Tianxiang Mai
- 核心结论:This study aimed to investigate the associations between different subtypes of Metabolic dysfunction-associated fatty liver disease (MAFLD) and carotid plaque, and to evaluate the independent association of the Fibrosis-4 (FIB-4) index with carotid plaque risk. Multivariable logistic regression was…
- 原文DOI / 链接:10.1371/journal.pone.0343781

6. IMUDINO: Self-distilled Foundation Model for Inertial Sensing

- 发表期刊 / 影响因子:arXiv Preprint
- 发表时间:2026-11-05
- 第一/通讯作者:Zhoutong Ye, Chengwen Zhang, Zixuan Wang, Xufeng Jian, Zhe He
- 核心结论:We present IMUDINO, the first foundation model (FM) for inertial sensing to achieve competitive performance across diverse sensing tasks including human activity recognition (HAR), health, motion capture, and odometry. Prior inertial FMs often focus on one subfield of inertial sensing and sometimes…
- 原文DOI / 链接:https://arxiv.org/abs/2611.07437v1

7. Typed Knowledge over Constructive Institutions

- 发表期刊 / 影响因子:arXiv Preprint
- 发表时间:2026-11-05
- 第一/通讯作者:Hans-Martin Will
- 核心结论:Type theories designed for mathematics or program verification need only that a proposition holds; proof irrelevance in conversion erases, at the kernel level, how a claim was established. A persistent knowledge store, by contrast, must retain that record: who established a claim, by which derivatio…
- 原文DOI / 链接:https://arxiv.org/abs/2611.07432v1

8. Two-stage residual learning for calibrated uncertainty in solar irradiance forecasts

- 发表期刊 / 影响因子:arXiv Preprint
- 发表时间:2026-11-05

- 第一/通讯作者:RR Flanagan, E Kasha, PW Talbot, AS Epiny, MR Deinert
- 核心结论:Accurate forecasting of renewable generation requires not only point predictions but also reliable estimates of the uncertainty attached to them, particularly at extended horizons. Machine learning models are increasingly used to predict solar output, but they typically report a single value and cap…
- 原文DOI / 链接:https://arxiv.org/abs/2611.07444v1

二、科研公众号干货

说明:微信公众号文章检索需要微信搜一搜等公开渠道的即时访问能力,当前自动检索模块正在优化中,建议通过以下方式自行获取: - 关注「肿瘤放疗」、「生信技能树」、「医学僧的科研日记」等公众号 - 使用微信搜一搜搜索关键词:肿瘤放疗 生信分析 - 本板块将在后续版本中接入 WeChat 公开内容 API

三、每日高分经典精读

Stereotactic Body Radiation Therapy for Hepatocellular Carcinoma: Current Trends and Future Directions

- 发表期刊 / 影响因子 / 总引用量:International Journal of Radiation Oncology, Biology, Physics / IF=7.0 / 引用量:200+
- 研究背景与里程碑意义:SBRT has emerged as a definitive local treatment for HCC, but optimal dosing, patient selection, and integration with systemic therapies remain active areas of investigation.
- 核心实验设计与方法:Comprehensive review of clinical trials and retrospective studies of SBRT for HCC, including dose-escalation studies, toxicity analyses, and combination strategies.
- 关键结论与行业影响:SBRT achieves local control rates of 80-95% for early-stage HCC with low toxicity. Challenges include BCLC stage-specific outcomes, liver function preservation, and optimal sequencing with immune checkpoint inhibitors.
- 创新点:Systematic framework for SBRT dose selection based on tumor size, location, and liver function; proposed risk-adapted fractionation schedules.
- 原文DOI / 链接:https://doi.org/10.1016/j.ijrobp.2023.01.045

四、当日总结

今日共精选8篇最新研究+1篇经典精读。关注方向:Evaluation of the Clinical Utility of Circulating Tumor DNA in the Management of Patients With Nonmetastatic Head and Neck Cancer: Protocol for a Prospective Bicentric Study (CIRVAD).
