

Basic Statistics

Sudipta Das

Assistant Professor,
Department of Computer Science,
Ramakrishna Mission Vivekananda Educational & Research Institute

1 Statistical Inference

- Introduction
- Point Estimation

Chapter 8: Statistical Inference

Sudipta Das

- The main objective in any statistical enquiry is the properties of one or more population.
 - However, the population(s) is (are) usually unknown to us, and we simply have a sample from the population (or, a sample from each of the given populations)

- Statistical Inference:-

Given the properties of the sample (or, of the samples), to infer about those of the population(s) is the problem of statistical inference

- It is analogous to the inductive logic, the only difference being that the induction is achieved under probabilistic framework
 - Probability comes due to random sampling
- It is a process of going over from the known sample to unknown population.

Statistical Inference III

Statistical set-up of the problem of inference

- Let (X_1, X_2, \dots, X_n) be a random sample of size n drawn from a population (discrete/continuous) with p.m.f/p.d.f $f(\underline{x}; \underline{\theta}) = f_{\underline{\theta}}(\underline{x})$, where $\underline{\theta}$ is the unknown parameter(s) of interest.
 - Our problem is to infer about $\underline{\theta}$
- Let Θ be the set of all possible values of θ
 - Θ is called the parameter space
- Note:
 - In the problem of statistical inference, Θ is known, although θ is unknown.
 - Example 1: $X_1, X_2, \dots, X_n \sim \text{Bernoulli}(p)$
 - $\theta = p$ is unknown, $\Theta = [0, 1]$ is known
 - Example 2: $X_1, X_2, \dots, X_n \sim \text{Normal}(\mu, \sigma)$
 - $\underline{\theta} = [\mu, \sigma]'$ is unknown, $\Theta = (-\infty, \infty) \times (0, \infty)$ is known
 - $\theta = \mu$ is unknown, $\Theta = (-\infty, \infty)$ is known
 - $\theta = \sigma$ is unknown, $\Theta = (0, \infty)$ is known

- Statistical Inference
 - ① Estimation
 - i Point Estimation
 - ii Interval Estimation
 - ② Hypothesis-testing

1 Estimation:-

Here, we have **no idea** about the true value of θ and the problem is **to estimate** the likely value of θ on the basis of the random sample (X_1, X_2, \dots, X_n) drawn from the population

i Point Estimation:-

Here, we estimate θ by a **single** value (i.e., by a point)

- Let $T = T(X_1, X_2, \dots, X_n)$ be a statistic which is used to estimate the parameter θ , is called an **estimator** of θ
- For the observed sample $(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n)$, the observed value of the estimator, namely,

$$t = T(x_1, x_2, \dots, x_n)$$

is called an estimate of θ

ii Interval Estimation:-

Here, we estimate θ by an **interval** of values

- Let $T_1 = T_1(X_1, X_2, \dots, X_n)$ and $T_2 = T_2(X_1, X_2, \dots, X_n)$ be two statistics such that

$$P[T_1 \leq \theta \leq T_2] = 1 - \alpha,$$

where α is a pre-assigned small quantity. Usually, we take $\alpha = 0.05$ or 0.01 etc.

- If $\alpha = 0.05$, then $P[T_1 \leq \theta \leq T_2] = 0.95$. Hence, the observed values of $[T_1, T_2] = [t_1, t_2]$, say, is called a 95% confidence interval of θ

2 Hypothesis-testing:-

Here we have some idea about the true value of θ , in the form of a hypothesis, say, $\theta = \theta_0$,

- Our problem is **to judge or test** the validity/ feasibility/ tenability of the given hypothesis $\theta = \theta_0$ on the basis of random sample of the population

Chapter 8a: Point Estimation

Sudipta Das

- In this case, we estimate θ by a **single** value (i.e., by a point)
 - Let $T = T(X_1, X_2, \dots, X_n)$ be a statistic which is used to estimate the parameter θ , is called an **estimator** of θ
 - For the observed sample $(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n)$, the observed value of the estimator, namely,

$$t = T(x_1, x_2, \dots, x_n)$$

is called an **estimate** of θ .

- Methods of calculating point estimates
 - Method of moments
 - Method of maximum likelihood

- Method of moments:

It consists in equating the first few moments of the population ($\mu'_k = E[X^k]$) with the corresponding moments of the sample

$$\left(m'_k = \frac{1}{n} \sum_{i=1}^n x_i^k \right), \text{ i.e.,}$$

$$\mu'_k = m'_k.$$

- The method of moments procedure:
Suppose there are l parameters to be estimated, say $\theta = (\theta_1, \dots, \theta_l)'$.
 - Find l population moments, μ'_k , for $k = 1, 2, \dots, l$.
 - μ'_k will contain one or more parameters $\theta_1, \dots, \theta_l$.
 - Find the corresponding l sample moments, m'_k , for $k = 1, 2, \dots, l$.
 - The number of sample moments should equal the number of parameters to be estimated.
 - From the system of equations, $\mu'_k = m'_k$, for $k = 1, 2, \dots, l$, solve for the parameter $\theta = (\theta_1, \dots, \theta_l)'$.
 - This will be a moment estimator of $\hat{\theta}$

Point Estimation (method of moment)

To Estimate	Notation	Point
Mean	\bar{X}	$\frac{1}{n} \sum_{i=1}^n X_i$
Proportion	\hat{p}	$\frac{1}{n} \sum_{i=1}^n I_{(X_i=1)}$
Variance	S_n^2	$\frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$

- Method of maximum likelihood:

It consists in choosing as estimator of $\underline{\theta}$ that statistic, which when substituted for $\underline{\theta}$, maximizes the likelihood function

$$L = f_{\underline{X}}(\underline{x}, \underline{\theta}) = f_{X_1, \dots, X_n}(x_1, \dots, x_n, \theta_1, \dots, \theta_l).$$

- Procedure to find maximum likelihood estimate (mle):

- Define the likelihood function, $L(\theta)$.
- Often it is easier to take the natural logarithm (\ln) of $L(\theta)$.
- When applicable, differentiate $l(\theta) = \ln L(\theta)$ with respect to θ , and then equate the derivative to zero.
- Solve for the parameter θ , and we will obtain $\hat{\theta}$.
- Check whether it is a maximizer or global maximizer.

Some Desirable Properties of Point Estimators

- Unbiased
- Sufficiency
- Consistency
- Efficiency

Unbiased Estimators

- A point estimator $\hat{\theta}$ is called an unbiased estimator of the parameter θ if

$$E(\hat{\theta}) = \theta$$

for all possible values of θ .

- Otherwise $\hat{\theta}$ is said to be biased.
 - Furthermore, the bias of $\hat{\theta}$ is given by

$$B(\hat{\theta}) = E[\hat{\theta}] - \theta.$$

- *Theorems:*

- The sample mean, $\bar{X} \left(= \frac{1}{n} \sum_{i=1}^n X_i \right)$ is an unbiased estimator of the population mean μ .

- Sketch of proof: $E[\bar{X}] = E \left(\frac{1}{n} \sum_{i=1}^n X_i \right) = \mu$

Unbiased Estimators III

- The statistic, $s_n^2 \left(= \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2 \right)$ is not an unbiased estimator of the population variance σ^2 .

- Sketch of proof:

$$\begin{aligned} E[S_n^2] &= E\left[\frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2\right] = \frac{1}{n} E\left[\sum_{i=1}^n ((X_i - \mu) - (\bar{X} - \mu))^2\right] \\ &= \frac{1}{n} \left(\sum_{i=1}^n E(X_i - \mu)^2 - nE(\bar{X} - \mu)^2 \right) = \frac{1}{n} \left(n\sigma^2 - n\frac{\sigma^2}{n} \right) = \left(\frac{n-1}{n} \right) \sigma^2 \end{aligned}$$

- The sample variance, $s^2 \left(= \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2 \right)$ is an unbiased estimator of the population variance σ^2 .

- Sketch of proof: $E[S^2] = E\left[\frac{n}{n-1} S_n^2\right] = \frac{n}{n-1} E[S_n^2] = \sigma^2$

Mean Square Error of an Estimator

- The mean square error of the estimator $\hat{\theta}$, denoted by $MSE(\hat{\theta})$, is defined as

$$MSE(\hat{\theta}) = E(\hat{\theta} - \theta)^2.$$

- Theorem:** $MSE(\hat{\theta}) = Var(\hat{\theta}) + B^2(\hat{\theta})$.

- Sketch of proof:

$$\begin{aligned} MSE(\hat{\theta}) &= E(\hat{\theta} - \theta)^2 = E\left((\hat{\theta} - E(\hat{\theta})) + (E(\hat{\theta}) - \theta)\right)^2 \\ &= E\left(\hat{\theta} - E(\hat{\theta})\right)^2 + \left(E(\hat{\theta}) - \theta\right)^2 + 2\left(E(\hat{\theta}) - \theta\right)\left(E(\hat{\theta}) - E(\hat{\theta})\right) \\ &= Var(\hat{\theta}) + B^2(\hat{\theta}). \end{aligned}$$

- If $\hat{\theta}$ is an unbiased estimator of θ , then

$$B^2(\hat{\theta}) = 0 \text{ and } MSE(\hat{\theta}) = Var(\hat{\theta}).$$

- Minimum variance unbiased estimator (MVUE) of θ :
 - The unbiased estimator $\hat{\theta}$ that minimizes the mean square error is called the *MVUE* of θ .

Sufficient Estimators

- A statistic U is a sufficient statistic for a parameter θ if U contains all the information available in the data about the value of θ .
- If $U(\mathbf{X})$ is a sufficient statistic for a parameter θ , then any inference about θ should depend on the sample \mathbf{X} only through the value $U(\mathbf{X})$.
 - That is, if \mathbf{x} and \mathbf{y} are two sample points such that $U(\mathbf{x}) = U(\mathbf{y})$, then the inference about θ should be the same whether $\mathbf{X} = \mathbf{x}$ or $\mathbf{X} = \mathbf{y}$ is observed.
- Example
 - the sample mean (\bar{X}) may contain all the relevant information about the parameter μ , and in that case $U(\mathbf{X}) = \bar{X}$ is called a sufficient statistic for μ .

Formal definition

- Let X_1, \dots, X_n be a random sample from a probability distribution with unknown parameter θ .
 - Then, the statistic $U = g(X_1, \dots, X_n)$ is said to be **sufficient statistic** for θ if $f_{X_1, \dots, X_n}(x_1, \dots, x_n | U = u)$ does not depend on θ for any value of u .
 - An estimator of θ that is a function of a sufficient statistic for θ is said to be a **sufficient estimator** of θ .

Sufficient Estimators III

- Example: Let X_1, \dots, X_n be i.i.d. Bernoulli random variables with parameter θ . Then $U = \sum_{i=1}^n X_i$ is sufficient for θ .

- Sketch of proof:

$$\begin{aligned} f(X_1, \dots, X_n; \theta) &= \theta^{\sum_{i=1}^n X_i} (1 - \theta)^{n - \sum_{i=1}^n X_i} \\ &= \theta^U (1 - \theta)^{n-U}. \end{aligned}$$

Since, $U \sim \text{Bin}(n, \theta)$,

$$f(U; \theta) = {}^n C_U \theta^U (1 - \theta)^{n-U}.$$

Thus,

$$f(x_1, \dots, x_n | U = u) = \frac{f(x_1, \dots, x_n, u)}{f_U(u)} = \begin{cases} \frac{1}{{}^n C_U} & ; \text{ if } u = \sum x_i \\ 0 & ; \text{ otherwise.} \end{cases}$$

Neyman–Fisher factorization theorem to *spot* a sufficient statistic.

- *Theorem:* -

Let U be a statistic based on the random sample X_1, \dots, X_n . Then, U is a sufficient statistic for θ if and only if the joint p.d.f or p.m.f., $f(x_1, \dots, x_n; \theta)$ can be factored into two non-negative functions, i.e.,

$$f(x_1, \dots, x_n; \theta) = g(u, \theta)h(x_1, \dots, x_n), \text{ for all } x_1, \dots, x_n,$$

where

- $g(u, \theta)$ is a function only of u and θ
- and $h(x_1, \dots, x_n)$ is a function of only x_1, \dots, x_n and not of θ .

Sufficient Estimators V

- Sketch of proof: discrete case
 - Sufficient \Rightarrow Factorization

$$\begin{aligned}f(x_1, \dots, x_n; \theta) &= P_\theta(X_1 = x_1, \dots, X_n = x_n, U = u) \\&= P_\theta(X_1 = x_1, \dots, X_n = x_n | U = u) P_\theta(U = u) \\&\stackrel{\text{uff.}}{=} h(x_1, \dots, x_n) g(u, \theta).\end{aligned}$$

- Factorization \Rightarrow Sufficient

$$\begin{aligned}P_\theta(X_1 = x_1, \dots, X_n = x_n | U = u) &= \frac{P_\theta(X_1 = x_1, \dots, X_n = x_n, U = u)}{P_\theta(U = u)} \\&= \begin{cases} \frac{P_\theta(X_1 = x_1, \dots, X_n = x_n, U = u)}{P_\theta(U = u)} & \text{if } (x_1, \dots, x_n) \in A_u \\ 0 & \text{if } (x_1, \dots, x_n) \notin A_u, \end{cases}\end{aligned}$$

where A_u is the set of all (x_1, \dots, x_n) such that U maps it into u .

Sufficient Estimators VI

- When $(x_1, x_2, \dots, x_n) \in A_u$,

$$\begin{aligned} P_\theta(X_1 = x_1, \dots, X_n = x_n | U = u) &= \frac{P_\theta(X_1 = x_1, \dots, X_n = x_n, U = u)}{P_\theta(U = u)} \\ &= \frac{P_\theta(X_1 = x_1, \dots, X_n = x_n)}{P_\theta(U = u)} \\ &= \frac{f(x_1, \dots, x_n, \theta)}{\sum_{(x_1, \dots, x_n) \in A_u} f(x_1, \dots, x_n, \theta)} \\ &\stackrel{\text{fact}}{=} \frac{g(u, \theta)h(x_1, \dots, x_n)}{\sum_{(x_1, \dots, x_n) \in A_u} g(u, \theta)h(x_1, \dots, x_n)} \\ &= \frac{h(x_1, \dots, x_n)}{\sum_{(x_1, \dots, x_n) \in A_u} h(x_1, \dots, x_n)} \perp \theta \end{aligned}$$

- When $(x_1, x_2, \dots, x_n) \notin A_u$,

$$\begin{aligned} P_\theta(X_1 = x_1, \dots, X_n = x_n | U = u) &= \frac{P_\theta(X_1 = x_1, \dots, X_n = x_n, U = u)}{P_\theta(U = u)} \\ &= 0 \perp \theta \end{aligned}$$

- Procedure to verify Sufficiency
 - 1 Obtain the joint pdf or pmf $f_{\theta}(x_1, \dots, x_n)$.
 - 2 If necessary, rewrite the joint pdf or pmf in terms of the given statistic and parameter so that one can use the factorization theorem.
 - 3 Define the functions g and h , in such a way that g is a function of the statistic and parameter only and h is a function of the observations only.
 - 4 If step 3 is possible, then the statistic is sufficient. Otherwise, it is not sufficient.

Sufficient Estimators VIII

- Example: Let X_1, \dots, X_n denote a random sample from a geometric population with parameter p . Show that \bar{X} is sufficient for p .

- Sketch of proof:

$$\begin{aligned} f(x_1, \dots, x_n) &= \prod_{i=1}^n p(1-p)^{x_i-1} \\ &= p^n (1-p)^{-n + \sum_{i=1}^n x_i} \\ &= p^n (1-p)^{-n + n\bar{x}} \\ &= g(\bar{x}, p) \underbrace{h(x_1, \dots, x_n)}_1 \end{aligned}$$

- *Joint Sufficiency:*

Two statistics U_1 and U_2 are said to be jointly sufficient for the parameters θ_1 and θ_2 if the conditional distribution of X_1, \dots, X_n given U_1 and U_2 does not depend on θ_1 or θ_2 .

In general, the statistic $U = (U_1, \dots, U_n)$ is jointly sufficient for $\theta = (\theta_1, \dots, \theta_n)$ if the conditional distribution of X_1, \dots, X_n given U is free of θ .

- Factorization criteria for Joint Sufficiency

Theorem: -

The two statistics U_1 and U_2 are jointly sufficient for θ_1 and θ_2 if and only if the likelihood function can be factored into two non-negative functions,

$$f(x_1, \dots, x_n; \theta_1, \theta_2) = g(u_1, u_2; \theta_1, \theta_2)h(x_1, \dots, x_n)$$

where $g(u_1, u_2; \theta_1, \theta_2)$ is only a function of $u_1, u_2; \theta_1$ and θ_2 , and $h(x_1, \dots, x_n)$ is free of θ_1 or θ_2

- Example: Let X_1, \dots, X_n be a random sample from $N(\mu, \sigma^2)$.
 - ① If μ is unknown and $\sigma^2 = \sigma_0^2$ is known, then $U_1 = \bar{X}$ is a sufficient statistic for μ .

$$L = \prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma_0} e^{-\frac{1}{2}\left(\frac{x_i - \mu}{\sigma_0}\right)^2} = \underbrace{(2\pi\sigma_0^2)^{-n/2} e^{-\frac{1}{2\sigma_0^2} \left[\sum_{i=1}^n x_i^2 \right]}}_{h(x_1, \dots, x_n)} \times \underbrace{e^{-\frac{1}{2\sigma_0^2} [-2n\mu\bar{x} + n\mu^2]}}_{g(u_1, \theta_1)}$$

- ② If $\mu = \mu_0$ is known and σ^2 is unknown, then $U_1 = \sum_{i=1}^n (X_i - \mu_0)^2$ is a sufficient statistic for σ^2 .

$$L = \prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}\left(\frac{x_i - \mu_0}{\sigma}\right)^2} = \underbrace{(2\pi)^{-n/2}}_{h(x_1, \dots, x_n)} \times \underbrace{\sigma^{-n} e^{-\frac{\sum_{i=1}^n (x_i - \mu_0)^2}{2\sigma^2}}}_{g(u_1, \theta_1)}$$

- 3 If μ and σ^2 are both unknown, then $U_1 = \sum_{i=1}^n X_i$ and $U_2 = \sum_{i=1}^n X_i^2$ are jointly sufficient for μ and σ^2 .

$$L = \prod_{i=1}^n \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{1}{2}\left(\frac{x_i - \mu}{\sigma}\right)^2} = \underbrace{(2\pi)^{-n/2}}_{h(x_1, \dots, x_n)} \times \underbrace{\sigma^{-n} e^{-\frac{1}{2\sigma^2} \left[\sum_{i=1}^n x_i^2 - 2\mu \sum_{i=1}^n x_i + n\mu^2 \right]}}_{g(u_1, \theta_1)}$$

- *Theorem:* -

If U is a sufficient statistic for θ , then the maximum likelihood estimator of θ , if unique, is a function of U .

- Sketch of proof.

$$f(x_1, \dots, x_n; \theta) \stackrel{\text{suff}}{=} g(u, \theta)h(x_1, \dots, x_n).$$

Thus, the joint pdf/pmf depends on θ only through the statistic U . To maximize L we need to maximize $g(U, \theta)$.

Sufficient Estimators XV

- *Theorem:* -

Let X_1, \dots, X_n be a random sample from a population with pdf or pmf of the exponential form

$$f(x_1, \dots, x_n; \theta) = \begin{cases} \exp[k(x)c(\theta) + S(x) + d(\theta)] & , x \in B \\ 0 & , x \notin B, \end{cases}$$

where B does not depend on the parameter θ . The statistic

$U = \sum_{i=1}^n k(X_i)$ is sufficient for θ .

- Sketch of proof.

$$\begin{aligned} f(x_1, \dots, x_n; \theta) &= \exp \left[c(\theta) \sum_{i=1}^n k(x_i) + \sum_{i=1}^n S(x_i) + nd(\theta) \right] \\ &= \underbrace{\exp \left[c(\theta) \sum_{i=1}^n k(x_i) + nd(\theta) \right]}_{g(u, \theta)} \underbrace{\exp \left[\sum_{i=1}^n S(x_i) \right]}_{h(x_1, \dots, x_n)}. \end{aligned}$$

Some Observations on Sufficiency

- 1 All the statistics need not be sufficient, in addition
- 2 Any function of a sufficient statistic needs not to be sufficient, however
- 3 Any one-to-one function of a sufficient statistic is also sufficient statistic

- *RAO-BLACKWELL Theorem:* -

Let X_1, \dots, X_n be a random sample with joint pmf or pdf $f(x_1, \dots, x_n; \theta)$ and let $U = (U_1, \dots, U_n)$ be jointly sufficient for $\theta = (\theta_1, \dots, \theta_n)$. If T is any unbiased estimator of $k(\theta)$, and if $T^* = E(T|U)$, then:

- T^* is an unbiased estimator of $k(\theta)$.

Sketch of proof: - $ET^* = E(E(T|U)) = E(T) = k(\theta)$.

Hence, T^* is an unbiased estimator of $k(\theta)$.

- T^* is a function of U , and does not depend on θ .

Sketch of proof: - Because U is sufficient for θ , the conditional distribution of any statistic (hence, for T), given U , does not depend on θ .

- $Var(T^*) \leq Var(T)$ for every θ , and $Var(T^*) < Var(T)$ for some θ unless $T^* = T$ with probability 1.

Sketch of proof: - $Var(T) = E(Var(T|U)) + Var(E(T|U)) = E(Var(T|U)) + Var(T^*)$.

Because $Var(T|U) \geq 0$ for all u , it follows that $E(Var(T|U)) \geq 0$. Hence, $Var(T^*) \leq Var(T)$.

Also $Var(T^*) = Var(T)$ iff $Var(T|U) = 0$ or T is a function of U , in which case $T^* = E(T|U) = T$.

More Observations

- If one is searching for an unbiased estimator with minimal variance, it has to be restricted to functions of a sufficient statistics.
- If $k(\theta) = \theta$, and T is an unbiased estimator of θ , then $T^* = E(T|U)$ will typically give the MVUE of θ .

- Minimal sufficient statistic:

A sufficient statistic $T(X)$ is called a minimal sufficient statistic if for any other sufficient statistic $T'(X)$,

$$T(X) = g(T'(X)),$$

i.e., $T(X)$ is a function of $T'(X)$.

- Intuitively, a minimal sufficient statistic most efficiently captures all possible information about the parameter θ .

Sufficient Estimators XX

- Lehmann and Scheffe method to find a minimal sufficient statistic
 - Let X_1, \dots, X_n be a random sample with pdf or pmf $f(x)$ that depends on a parameter θ .
 - Let (x_1, \dots, x_n) and (y_1, \dots, y_n) be two different sets of values of (X_1, \dots, X_n) .
 - Let $\frac{L(\theta; x_1, \dots, x_n)}{L(\theta; y_1, \dots, y_n)}$ be the ratio of the likelihoods evaluated at these two points.
 - Suppose it is possible to find a function $g(x_1, \dots, x_n)$ such that this ratio will be free of the unknown parameter θ if and only if $g(x_1, \dots, x_n) = g(y_1, \dots, y_n)$,
 - in other words

$$\frac{L(\theta; x_1, \dots, x_n)}{L(\theta; y_1, \dots, y_n)} \text{ is independent of } \theta \Leftrightarrow g(x_1, \dots, x_n) = g(y_1, \dots, y_n).$$

- If such a function g can be found, then $g(X_1, \dots, X_n)$ is a minimal sufficient statistic for θ .

- Example: -

Let (X_1, \dots, X_n) be a random sample from the *Bernoulli*(p), where p is unknown. Find a minimal sufficient statistic for p .

- Solution: - The ratio of the likelihoods is

$$\frac{L(x_1, \dots, x_n)}{L(y_1, \dots, y_n)} = \frac{p(x_1, \dots, x_n)}{p(y_1, \dots, y_n)} = \frac{p^{\sum x_i} (1-p)^{n-\sum x_i}}{p^{\sum y_i} (1-p)^{n-\sum y_i}} = \left(\frac{p}{1-p} \right)^{\sum x_i - \sum y_i}$$

This ratio is to be independent of p , if and only if

$$g(x_1, \dots, x_n) = \sum_{i=1}^n x_i = \sum_{i=1}^n y_i = g(y_1, \dots, y_n).$$

Therefore, $g(X_1, \dots, X_n) = \sum_{i=1}^n X_i$ is a minimal sufficient statistic for p .

Consistent Estimators

- A statistic is a consistent estimator if its value becomes closer to the value of the true parameter which is being estimated, as the sample size becomes larger.

Consistent Estimators II

Formal definition

- The estimator $\hat{\theta}_n$ is said to be a consistent estimator of θ if, for any $\epsilon > 0$,

$$\lim_{n \rightarrow \infty} P \left[|\hat{\theta}_n - \theta| \leq \epsilon \right] = 1$$

or equivalently,

$$\lim_{n \rightarrow \infty} P \left[|\hat{\theta}_n - \theta| > \epsilon \right] = 0,$$

i.e. $\hat{\theta}_n$ converges in probability to θ .

Consistent Estimators III

- Example: - Let X_1, \dots, X_n be a random sample with true mean μ and finite variance, σ^2 . Then, the sample mean \bar{X} is a consistent estimator of the population mean μ .

- Sketch of proof: -
Note that, for any positive r.v. X (Markov Inequality)

$$E[X] = \int_0^\epsilon x f_X(x) dx + \int_\epsilon^\infty x f_X(x) dx \geq \int_0^\epsilon x f_X(x) dx + \epsilon \int_\epsilon^\infty f_X(x) dx \geq \epsilon P(X \geq \epsilon).$$

Hence, (Chebyshev's Inequality)

$$P[|\bar{X} - \mu| \geq \epsilon] = P[(\bar{X} - E[\bar{X}])^2 \geq \epsilon^2] \leq \frac{E(\bar{X} - E[\bar{X}])^2}{\epsilon^2} = \frac{\text{Var}(\bar{X})}{\epsilon^2},$$

Thus,

$$\lim_{n \rightarrow \infty} P[|\bar{X} - \mu| \geq \epsilon] \leq \lim_{n \rightarrow \infty} \frac{\text{Var}(\bar{X})}{\epsilon^2} = \lim_{n \rightarrow \infty} \frac{\sigma^2}{n\epsilon^2} = 0$$

Consistent Estimators IV

Test for Consistency (Sufficient conditions)

- *Theorem:* - An unbiased estimator $\hat{\theta}_n$ of θ is a consistent estimator for θ if

$$\lim_{n \rightarrow \infty} \text{Var}(\hat{\theta}_n) = 0.$$

- Sketch of proof: - $\lim_{n \rightarrow \infty} P[|\hat{\theta}_n - \theta| \geq \epsilon] \stackrel{U.E.}{=} \lim_{n \rightarrow \infty} P[|\hat{\theta}_n - E(\hat{\theta}_n)| \geq \epsilon] \stackrel{C.I.}{\leq} \lim_{n \rightarrow \infty} \frac{\text{Var}(\hat{\theta}_n)}{\epsilon^2} = 0$
- An unbiased estimator $\hat{\theta}_n$ is consistent if $\text{Var}(\hat{\theta}_n) \rightarrow 0$ as $n \rightarrow \infty$.
- *Theorem:* - An estimator $\hat{\theta}_n$, with finite variance, is a consistent estimator for θ if

$$\lim_{n \rightarrow \infty} E(\hat{\theta}_n - \theta)^2 = 0.$$

- Sketch of proof: - $\lim_{n \rightarrow \infty} P[|\hat{\theta}_n - \theta| \geq \epsilon] \stackrel{C.I.}{\leq} \lim_{n \rightarrow \infty} \frac{E(\hat{\theta}_n - \theta)^2}{\epsilon^2} = 0$
- A biased estimator $\hat{\theta}_n$ is consistent if both $\text{Var}(\hat{\theta}_n) \rightarrow 0$ and $B(\hat{\theta}_n) \rightarrow 0$ as $n \rightarrow \infty$.

Consistent Estimators V

- Example: -

Let X_1, \dots, X_n be a random sample from $N(\mu, \sigma^2)$ population.

- Then the sample variance S^2 is a consistent estimator for σ^2 .

- Sketch of proof: -

$$(n-1) \frac{S^2}{\sigma^2} \sim \chi^2(n-1) \Rightarrow \text{Var} \left[(n-1) \frac{S^2}{\sigma^2} \right] = 2(n-1) \Rightarrow \text{Var}(S^2) = \frac{2\sigma^4}{n-1}.$$

S^2 is u.e. of σ^2 and $\lim_{n \rightarrow \infty} \text{Var}(S^2) = 0$.

- The maximum likelihood estimators \bar{X} and S_n^2 for μ and σ^2 , respectively, are consistent estimators for μ and σ^2 .

- Sketch of proof: -

\bar{X} is u.e. of μ and $\lim_{n \rightarrow \infty} \text{Var}(\bar{X}) = 0$, thus \bar{X} is consistent estimator of μ .

$$\text{Var}(S_n^2) = \text{Var} \left(\frac{n-1}{n} S^2 \right) = \frac{2(n-1)\sigma^4}{n^2}, \text{ thus } \lim_{n \rightarrow \infty} \text{Var}(S_n^2) = 0 \text{ and}$$

$$B(S_n^2) = \frac{n-1}{n} \sigma^2 - \sigma^2 = -\frac{1}{n} \sigma^2, \text{ thus } \lim_{n \rightarrow \infty} B(S_n^2) = 0$$

Efficiency

- It is a relative comparison between variances of two unbiased/biased estimators

Efficiency II

Formal Definitions

- If $\hat{\theta}_1$ and $\hat{\theta}_2$ are two unbiased estimators for θ , the efficiency of $\hat{\theta}_1$ relative to $\hat{\theta}_2$ is the ratio

$$e(\hat{\theta}_1, \hat{\theta}_2) = \frac{\text{Var}(\hat{\theta}_2)}{\text{Var}(\hat{\theta}_1)}.$$

- If $\text{Var}(\hat{\theta}_1) < \text{Var}(\hat{\theta}_2)$, or equivalently, $e(\hat{\theta}_1, \hat{\theta}_2) > 1$, then, $\hat{\theta}_1$ is relatively more efficient than $\hat{\theta}_2$.
- If $\hat{\theta}_1$ and $\hat{\theta}_2$ are two biased estimators for θ , the efficiency of $\hat{\theta}_1$ relative to $\hat{\theta}_2$ is the ratio

$$e(\hat{\theta}_1, \hat{\theta}_2) = \frac{E(\hat{\theta}_2 - \theta)^2}{E(\hat{\theta}_1 - \theta)^2} = \frac{\text{MSE}(\hat{\theta}_2)}{\text{MSE}(\hat{\theta}_1)}.$$

- If $\text{MSE}(\hat{\theta}_1) < \text{MSE}(\hat{\theta}_2)$, or equivalently, $e(\hat{\theta}_1, \hat{\theta}_2) > 1$, then, $\hat{\theta}_1$ is relatively more efficient than $\hat{\theta}_2$.

- Example:-

Let X_1, \dots, X_n , $n \geq 2$ be a random sample from a normal population with a true mean μ and variance σ^2 . Consider the following two estimators of σ^2 : $\theta_1 = S^2$, and $\theta_2 = S_n^2$. Find $e(\theta_1, \theta_2)$.

- Sketch of proof: -

$$MSE(\hat{\theta}_1) = 0 + \frac{2\sigma^4}{n-1} = \frac{2\sigma^4}{n-1}$$

$$MSE(\hat{\theta}_2) = \left(-\frac{\sigma^2}{n}\right)^2 + \frac{2(n-1)\sigma^4}{n^2} = \frac{(2n-1)\sigma^4}{n^2}$$

$$e(\hat{\theta}_1, \hat{\theta}_2) = \frac{(n-1)(2n-1)}{2n^2} < 1, \text{ for } n \geq 2.$$

Hence, S_n^2 is relatively more efficient than S^2 .

Uniformly Minimum Variance Unbiased Estimator

- An unbiased estimator $\hat{\theta}_0$, is said to be a uniformly minimum variance unbiased estimator (UMVUE) for the parameter θ if, for any other unbiased estimator $\hat{\theta}$,

$$\text{Var}(\hat{\theta}_0) \leq \text{Var}(\hat{\theta}),$$

for all possible values of θ .

Cramer–Rao Inequality:

(Lower bound for the variance of any estimator)

- *Theorem:* - Let X_1, \dots, X_n be a sample with pdf $f(\mathbf{x}|\theta)$, and let $W(\mathbf{X}) = W(X_1, \dots, X_n)$ be any estimator satisfying

$$\frac{d}{d\theta} E_{\theta}[W(\mathbf{X})] = \int_{\mathcal{X}} \frac{\delta}{\delta\theta} [W(\mathbf{x})f(\mathbf{x}|\theta)] d\mathbf{x}$$

and

$$\text{Var}_{\theta}[W(\mathbf{X})] < \infty,$$

then

$$\text{Var}_{\theta}[W(\mathbf{X})] \geq \frac{\left(\frac{d}{d\theta} E_{\theta}[W(\mathbf{X})]\right)^2}{E_{\theta}\left[\left(\frac{\delta}{\delta\theta} \ln f(\mathbf{x}|\theta)\right)^2\right]}.$$

Efficiency VI

- Sketch of proof: -
Note that

$$\begin{aligned}\frac{d}{d\theta} E_{\theta}[W(\mathbf{X})] &= \int_{\mathcal{X}} \frac{\delta}{\delta\theta} [W(\mathbf{x})f(\mathbf{x}|\theta)] d\mathbf{x} \\ &= \int_{\mathcal{X}} W(\mathbf{x}) \left[\frac{\delta}{\delta\theta} f(\mathbf{x}|\theta) \right] d\mathbf{x} \\ &= \int_{\mathcal{X}} W(\mathbf{x}) \left[\frac{\delta f(\mathbf{x}|\theta)}{\delta\theta} \frac{1}{f(\mathbf{x}|\theta)} \right] f(\mathbf{x}|\theta) d\mathbf{x} \\ &= E_{\theta} \left[W(\mathbf{X}) \frac{\delta}{\delta\theta} \ln f(\mathbf{x}|\theta) \right] \\ &= \text{Cov}_{\theta} \left(W(\mathbf{X}), \frac{\delta}{\delta\theta} \ln f(\mathbf{x}|\theta) \right),\end{aligned}$$

$$\text{since } E_{\theta} \left[\frac{\delta}{\delta\theta} \ln f(\mathbf{x}|\theta) \right] = \int_{\mathcal{X}} \left[\frac{\delta}{\delta\theta} \ln f(\mathbf{x}|\theta) \right] f(\mathbf{x}|\theta) d\mathbf{x} = \int_{\mathcal{X}} \left[\frac{\delta}{\delta\theta} f(\mathbf{x}|\theta) \right] d\mathbf{x} = \frac{d}{d\theta} \left[\int_{\mathcal{X}} f(\mathbf{x}|\theta) d\mathbf{x} \right] = 0.$$

Also

$$\text{Var}_{\theta} \left(\frac{\delta}{\delta\theta} \ln f(\mathbf{x}|\theta) \right) = E_{\theta} \left[\left(\frac{\delta}{\delta\theta} \ln f(\mathbf{x}|\theta) \right)^2 \right].$$

Now,

$$\frac{\left[\text{Cov}_{\theta} \left(W(\mathbf{X}), \frac{\delta}{\delta\theta} \ln f(\mathbf{x}|\theta) \right) \right]^2}{\text{Var}_{\theta} [W(\mathbf{X})] \text{Var}_{\theta} \left[\frac{\delta}{\delta\theta} \ln f(\mathbf{x}|\theta) \right]} \leq 1 \Rightarrow \frac{\left[\frac{d}{d\theta} E_{\theta}[W(\mathbf{X})] \right]^2}{E_{\theta} \left[\left(\frac{\delta}{\delta\theta} \ln f(\mathbf{x}|\theta) \right)^2 \right]} \leq \text{Var}_{\theta} [W(\mathbf{X})]$$

- *Corollary:* -

Let X_1, \dots, X_n be an iid random sample from a population with pdf or pmf $f_\theta(x)$ that depends on a parameter θ . If $\hat{\theta} = W(\mathbf{x})$ is an unbiased estimator of $\psi(\theta)$, then

$$\text{Var}(\hat{\theta}) \geq \frac{\left[\frac{d}{d\theta}\psi(\theta)\right]^2}{nE\left[\left(\frac{\delta}{\delta\theta}\ln f_\theta(x)\right)^2\right]}.$$

Efficiency VIII

- Sketch of proof: - Note that

$$\begin{aligned} E \left[\left(\frac{\delta}{\delta\theta} \ln f(\mathbf{x}|\theta) \right)^2 \right] &= E \left[\left(\frac{\delta}{\delta\theta} \ln \prod_{i=1}^n f_{\theta}(x_i) \right)^2 \right] = E \left[\left(\sum_{i=1}^n \frac{\delta}{\delta\theta} \ln f_{\theta}(x_i) \right)^2 \right] \\ &= \sum_{i=1}^n E \left[\left(\frac{\delta}{\delta\theta} \ln f_{\theta}(x_i) \right)^2 \right] + \sum_{i \neq j} E \left[\left(\frac{\delta}{\delta\theta} \ln f_{\theta}(x_i) \right) \left(\frac{\delta}{\delta\theta} \ln f_{\theta}(x_j) \right) \right] \\ &= nE \left[\left(\frac{\delta}{\delta\theta} \ln f_{\theta}(x) \right)^2 \right] + \sum_{i \neq j} E \left[\left(\frac{\delta}{\delta\theta} \ln f_{\theta}(x_i) \right) \right] E \left[\left(\frac{\delta}{\delta\theta} \ln f_{\theta}(x_j) \right) \right] \\ &= nE \left[\left(\frac{\delta}{\delta\theta} \ln f_{\theta}(x) \right)^2 \right] \end{aligned}$$

and

$$\left[\frac{d}{d\theta} E_{\theta}[W(\mathbf{x})] \right]^2 = \left[\frac{d}{d\theta} E_{\theta}[\hat{\theta}] \right]^2 = \left[\frac{d}{d\theta} \psi(\theta) \right]^2.$$

Hence,

$$\text{Var}(\hat{\theta}) \geq \frac{\left[\frac{d}{d\theta} E_{\theta}[\hat{\theta}] \right]^2}{E_{\theta} \left[\left(\frac{\delta}{\delta\theta} \ln f(\mathbf{x}|\theta) \right)^2 \right]} = \frac{\left[\frac{d}{d\theta} \psi(\theta) \right]^2}{nE \left[\left(\frac{\delta}{\delta\theta} \ln f_{\theta}(x) \right)^2 \right]}.$$

- *Corollary:* -

Let X_1, \dots, X_n be an iid random sample from a population with pdf or pmf $f_\theta(x)$ that depends on a parameter θ . If $\hat{\theta}$ is an unbiased estimator of θ , then

$$\text{Var}(\hat{\theta}) \geq \frac{1}{nE \left[\left(\frac{\delta}{\delta\theta} \ln f_\theta(x) \right)^2 \right]}.$$

Efficient Estimator

- If $\hat{\theta}$ is an unbiased estimator of θ and if

$$\text{Var}(\hat{\theta}) = \frac{1}{nE \left[\left(\frac{\delta}{\delta\theta} \ln f_{\theta}(x) \right)^2 \right]}$$

then $\hat{\theta}$ is a uniformly minimum variance unbiased estimator (UMVUE) of θ .

Sometimes $\hat{\theta}$ is also referred to as an efficient estimator.

- **Result:-** If the function $f(\cdot)$ is sufficiently smooth, specifically if $\frac{d}{d\theta} E_{\theta} \left(\frac{\delta}{\delta\theta} \ln f_{\theta}(x) \right) = \int \frac{\delta}{\delta\theta} \left[\left(\frac{\delta}{\delta\theta} \ln f_{\theta}(x) \right) f_{\theta}(x) \right] dx$, then

$$E_{\theta} \left[\left(\frac{\delta}{\delta\theta} \ln f_{\theta}(x) \right)^2 \right] = -E_{\theta} \left(\frac{\delta^2}{\delta\theta^2} \ln f_{\theta}(x) \right) = \text{Var} \left[\frac{\delta}{\delta\theta} \ln f_{\theta}(x) \right]$$

and for an unbiased estimator $\hat{\theta}$ for θ the Cramer-Rao inequality can be rewritten as

$$\text{Var}(\hat{\theta}) \geq \frac{1}{-nE \left(\frac{\delta^2}{\delta\theta^2} \ln f_{\theta}(x) \right)} = \frac{1}{n\text{Var} \left[\frac{\delta}{\delta\theta} \ln f_{\theta}(x) \right]}.$$

Efficiency XII

- Sketch of proof: -

$$\begin{aligned} E_{\theta} \left(\frac{\delta^2}{\delta\theta^2} \ln f_{\theta}(x) \right) &= E_{\theta} \left[\frac{\delta}{\delta\theta} \left(\frac{\delta}{\delta\theta} \ln f_{\theta}(x) \right) \right] = E_{\theta} \left[\frac{\delta}{\delta\theta} \left(\frac{\frac{\delta}{\delta\theta} f_{\theta}(x)}{f_{\theta}(x)} \right) \right] \\ &= E_{\theta} \left[\left(\frac{\frac{\delta^2}{\delta\theta^2} f_{\theta}(x)}{f_{\theta}(x)} \right) - \left(\frac{\frac{\delta}{\delta\theta} f_{\theta}(x)}{f_{\theta}(x)} \right)^2 \right] \\ &= \int \frac{\delta^2}{\delta\theta^2} f_{\theta}(x) dx - E_{\theta} \left[\left(\frac{\delta}{\delta\theta} \ln f_{\theta}(x) \right)^2 \right] \\ &\stackrel{\text{assumption}}{=} \frac{\delta}{\delta\theta} \int \frac{\delta}{\delta\theta} f_{\theta}(x) dx - E_{\theta} \left[\left(\frac{\delta}{\delta\theta} \ln f_{\theta}(x) \right)^2 \right] \\ &= \frac{d}{d\theta} E_{\theta} \left[\frac{\delta}{\delta\theta} \ln f_{\theta}(x) \right] - E_{\theta} \left[\left(\frac{\delta}{\delta\theta} \ln f_{\theta}(x) \right)^2 \right] \\ &= -E_{\theta} \left[\left(\frac{\delta}{\delta\theta} \ln f_{\theta}(x) \right)^2 \right] \end{aligned}$$

Hence,

$$\text{Var}(\hat{\theta}) \geq \frac{1}{n E_{\theta} \left[\left(\frac{\delta}{\delta\theta} \ln f_{\theta}(x) \right)^2 \right]} = \frac{1}{-n E_{\theta} \left(\frac{\delta^2}{\delta\theta^2} \ln f_{\theta}(x) \right)} = \frac{1}{n \text{Var}_{\theta} \left[\frac{\delta}{\delta\theta} \ln f_{\theta}(x) \right]}.$$

- Example: - Let X_1, \dots, X_n be a random sample from an $N(\mu, \sigma^2)$ population. Then \hat{X} is an efficient estimator for μ .

- Sketch of proof: - $l(x, \mu) = \ln f(x, \mu) = c - \frac{(x-\mu)^2}{2\sigma^2}$.

Thus,

$$\frac{\delta}{\delta \mu} l(x, \mu) = \frac{x - \mu}{\sigma^2} \text{ and } \frac{\delta^2}{\delta \mu^2} l(x, \mu) = -\frac{1}{\sigma^2}.$$

Hence,

$$\frac{1}{nE \left[\left(\frac{\delta}{\delta \theta} \ln f_{\theta}(x) \right)^2 \right]} = \frac{1}{nE \left[\left(\frac{\delta}{\delta \mu} l(x, \mu) \right)^2 \right]} = \frac{1}{nE \left[-\frac{\delta^2}{\delta^2 \mu} l(x, \mu) \right]} = \frac{\sigma^2}{n} = \text{Var}(\bar{X})$$

Note: -

- For a given problem UMVUE may not exist.
- Even when an UMVUE exists, it is not necessary that it have a variance equal to the Cramer–Rao lower bound.
- The term $I(\theta) = E \left[\left(\frac{\delta}{\delta \theta} \ln f_{\theta}(x) \right)^2 \right]$ is called the Fisher information.
- It can be shown that the Fisher information in a sample of size n , denoted by $I_n(\theta)$, is n times the Fisher information in one observation. That is,

$$I_n(\theta) = nI(\theta).$$