

# CREATE A GDP BASED ENTITY

A climada additional module

Version December 14

## Contents

Part 1: All-in-one functions.....	2
1.    All-in-one functions.....	2
1a)    Create centroids and base entity for a country.....	2
1b)    GDP based entity for a certain region within a country.....	3
Part 2: Background information and individual functions .....	4
2.    Night time lights for spatial distribution of values .....	4
3.    Gross Domestic Product (GDP) data .....	7
3a)    GDP today.....	7
3b)    Future projection of GDP .....	9
4.    Countries: Use polygons and rasters as helper files.....	11
5.    Night light intensities as a proxy for spatial distribution of values within a country: Linear or nonlinear relationship between light intensity and asset value.....	12
6.    Create the centroids.....	15
6a)    Define the resolution .....	15
6b)    Cut out coastal areas (hollow out) and expand with a buffer zone .....	15
6c)    Create the centroids.....	16
7.    Create the entity .....	18
7a)    Base entity: assets sum up to 100 for a given country.....	18
7b)    Entity with assets based on the country's GDP today .....	21
7c)    Future entity with assets based on the country's future GDP (e.g. 2030) .....	22
7d)    Select a specific region within a country .....	22
8.    Appendix .....	24
8a)    Downscale resolution of original night light tif in ArcGIS.....	24
8b)    Night light and GDP relationship .....	25

## Part 1: All-in-one functions

### 1. All-in-one functions

#### 1a) Create centroids and base entity for a country

The following function creates a portfolio for a specific country, consisting of the base entity and centroids.

MATLAB call:

```
[centroids,      climada_create_centroids_entity_base(country_name,
entity_base]=    asset_resolution_km, hollowout, check_for_groups, night_light,
                    pp,borders,border_mask,check_figure, save_on_entity_centroids)

for coastal areas only
climada_create_centroids_entity_base('Mexico',10,1)

for the entire country
climada_create_centroids_entity_base('Mexico',10,1)
```

All input parameters are optional. If not given as an input, the country can be selected through a graphical user interface. The required resolution can be specified as a value in km; however default value is roughly 10 km. Per default, the base entity and centroids will be saved in the folders `..\climada\data\entities` and `..\climada\data\system\centroids`, respectively.




Figure 1: Graphical user interface to select country.

#### Output

```
>> climada_create_centroids_entity_base('Mexico',10,1);
0)      a) Load world borders including regions
0)      b) Load border masks... done
*** Mexico on roughly 10 km ***
Transform night lights to values (nonlinearly or linearly)
y =  0.0000*x^0 + -0.0817*x^1 + 0.0172*x^2
Distribute values according to night lights within Mexico on
a 10 km resolution
1) Downscale distributed values to ~10km ... Requested resolution
(9 km) corresponds already to input matrix (9.26 km)
(exactly ~9km) done
2) Create centroids for Mexico on a ~9 km resolution
   a) Create buffer of ~50km
   b) Select entire country (hollowout is set to 0)
```

```
Requested resolution (9 km) corresponds already to input
matrix (9.26 km)
--> 36681 centroids (8692 on land and within buffer, 2891
outside)
d) Save centroids in

\\CHRB1065.CORP.GWPNET.COM\homes\X\S3BXXW\Documents\lea\climada_test_environment\climada\data\system\centroids_Mexico_9km_
3) Create base entity
a) Load wildcard entity without assets (damagefunctions, measures,
discount)
b) Take assets from distributed values matrix
c) Encode assets to centroids
d) Save entity_base in
\\CHRB1065.CORP.GWPNET.COM\homes\X\S3BXXW\Documents\lea\climada_test_environment\climada\data\entities\entity_Mexico_base_-0817x1_0172x2_9km_
e) Save entity as excel-file
- Assets sheet
- Damagefunctions sheet
- Measures sheet
- Discount sheet
Save entity as xls file
\\CHRB1065.CORP.GWPNET.COM\homes\X\S3BXXW\Documents\lea\climada_test_environment\climada\data\entities\entity_Mexico_base_-0817x1_0172x2_9km_.xls
>>
```

## 1b) GDP based entity for a certain region within a country

The function `climada_create_GDP_entity` wraps it all in one function. It creates a portfolio with centroids, as an intermediary step creates the base entity, and scales the entity assets up to GDP today and future GDP projection.

MATLAB call:

```
[centroids, entity, climada_create_GDP_entity(country_name, polygon)
entity_forecast] =
climada_create_GDP_entity('Mexico')
```

## Part 2: Background information and individual functions

### 2. Night time lights for spatial distribution of values

Stable night time lights can be downloaded from NOAA, e.g. the data for 2010, Version 4 DMSP-OLS Night time Lights Time Series

<http://ngdc.noaa.gov/eog/dmsp/downloadV4composites.html#AVSLCFC><sup>1</sup>

The files are cloud-free composites made using all the available archived DMSP-OLS smooth resolution data for calendar years. In cases where two satellites were collecting data - two composites were produced. The products are **30 arc second grids**, spanning -180 to 180 degrees longitude and -65 to 75 degrees latitude.

Credit: Image and data processing by NOAA's National Geophysical Data Center. DMSP data collected by US Air Force Weather Agency

A number of constraints are used to select the highest quality data for entry into the composites:

- Data are from the center half of the 3000 km wide OLS swaths. Lights in the center half have better geolocation, are smaller, and have more consistent radiometry.
- Sunlit data are excluded based on the solar elevation angle.
- Glare is excluded based on solar elevation angle.
- Moonlit data are excluded based on a calculation of lunar illuminance.
- Observations with clouds are excluded based on clouds identified with the OLS thermal band data and NCEP surface temperature grids.
- Lighting features from the aurora have been excluded in the northern hemisphere on an orbit-by-orbit manner using visual inspection.

Each composite set is named with the satellite and the year (F121995 is from DMSP satellite number F12 for the year 1995). Three image types are available as geotiffs for download from the version 4 composites:

- **F1?YYYY\_v4b\_stable\_lights.avg\_vis.tif:** The cleaned up avg\_vis contains the lights from cities, towns, and other sites with persistent lighting, including gas flares. Ephemeral events, such as fires have been discarded. Then the background noise was identified and replaced with values of zero. **Data values range from 1-63.** Areas with zero cloud-free observations are represented by the value 255.



Figure 2: Night time lights( "Nighttime Lights.bmp") in 2010. Credit: Image and data processing by NOAA's National Geophysical Data Center. DMSP data collected by US Air Force Weather Agency.

<sup>1</sup> In the past, also at [http://www.ngdc.noaa.gov/dmsp/data/web\\_data/v4composites/F182010.v4.tar](http://www.ngdc.noaa.gov/dmsp/data/web_data/v4composites/F182010.v4.tar) and <http://www.ngdc.noaa.gov/dmsp/downloadV4composites.html>

Create a GDP based entity, a climada additional module

The downloaded tif file is too big to be read directly in matlab, therefore resample (downscale resolution) in ArcGIS. See Appendix for further information on resampling in ArcGIS. Save in png-file format ...\\climada\_additional\\2\_globalGDP\\data\\night\_light\_2010\_10km.png

Read stable night light in matlab and save variable as "night\_light\_2010\_10km.mat" (1680 x 4320 cells, 10km resolution) in ...\\climada\_additional\\2\_globalGDP\\data\\night\_light\_2010\_10km.mat

MATLAB call:

```
night_light = climada_night_light_read(png_filename, check_figure,
                                         check_printplot, save_on)
climada_night_light_read('night_light_2010_10km.png')

.values      1680x4320 sparse double indicating night light intensities in
               the range 180 to 180 degrees longitude and -65 to 75 degrees
               latitude
.lon_range   Longitudinal range of masks, e.g. [-180 180]
.lat_range   Latitudinal range of masks, e.g. [-65 75]
.resolution_x Resolution in x direction in degree, e.g. 0.0833°, corresponds
               roughly to 10km
.resolution_y Resolution in y direction in degree, e.g. 0.0833°, corresponds
               roughly to 10km
.comment     Night time lights, 2010
```




Figure 3: Stable night lights in matlab (1680 x 4320 cells, 10km resolution), zoomed to Europe.




Figure 4: Stable night lights in matlab (1680 x 4320 cells, 10km resolution) on a global scale.

### 3. Gross Domestic Product (GDP) data

#### 3a) GDP today

GDP Data is available from the World Bank

<http://data.worldbank.org/indicator/NY.GDP.MKTP.CD>

##### GDP (current US\$)

 DATABANK


GDP at purchaser's prices is the sum of gross value added by all resident producers in the economy plus any product taxes and minus any subsidies not included in the value of the products. It is calculated without making deductions for depreciation of fabricated assets or for depletion and degradation of natural resources. Data are in current U.S. dollars. Dollar figures for GDP are converted from domestic currencies using single year official exchange rates. For a few countries where the official exchange rate does not reflect the rate effectively applied to actual foreign exchange transactions, an alternative conversion factor is used.

World Bank national accounts data, and OECD National Accounts data files.

Catalog Sources World Development Indicators



Figure 5: GDP Data from Worldbank Website, July 2012.



<http://data.worldbank.org/indicator/NY.GDP.MKTP.CD/countries>

Figure 6: GDP 2010 per country.

##### GDP per capita (current US\$)

 DATABANK


GDP per capita is gross domestic product divided by midyear population. GDP is the sum of gross value added by all resident producers in the economy plus any product taxes and minus any subsidies not included in the value of the products. It is calculated without making deductions for depreciation of fabricated assets or for depletion and degradation of natural resources. Data are in current U.S. dollars.

World Bank national accounts data, and OECD National Accounts data files.

Catalog Sources World Development Indicators



Figure 7: GDP per capita data from Worldbank Website, July 2012.



<http://data.worldbank.org/indicator/NY.GDP.PCAP.CD/countries>

Figure 8: GDP 2010 per capita.

Use the excel file as a key to match country names of the World Bank with climada world map  
[..\\climada\\_additional\\GDP\\_entity\\data\\country\\_names\\_key\\_between\\_worldbank\\_and\\_worldmap.xlsx](#)

MATLAB call:

```
GDP = climada_GDP_read(xlsfilename,special_cases_on,
check_names, save_on, silent_mode)
climada_GDP_read('World_GDP_current_1960_2010.xls')

.country_names      216x1 cell, list with all country names, e.g. Afghanistan,
Albania etc.
.year              1x51 double, with years from 1960 to 2010
.value             216x51 double, GDP value per year and country
.description        GDP in USD
.comment            World Bank/IMF GDP data, from year 1960 to 2010
.country_borders_index  216x1 cell, index that links GDP with border mask
structure (see next chapter), e.g.
GDP.country_borders_index(1) = 52; country number 1
in GDP list is Afghanistan and is on position 51 in the
borders structure. One country in the GDP list can index
to multiple countries in the border mask structure, for
instance GDP India (position 88) refers to border masks
position 55 (Kashmir and Jammu (claimed by India and
Pakistan) and 100 (India).
```

Output

```
***  

GDP in USD  

Worldbank/IMF GDP data, from year 1960 to 2010  

read successfully and loaded into workspace  

GDP available for 216 countries  

***  

***  

Check country names...  

Climada world border file loaded to compare country names.  

- GDP Australia is applied to  

    Christmas Island, Cocos (Keeling) Islands,  

Australia, Norfolk Island, Heard Island and Mcdonald Islands  

39: no match found for GDP country name Channel Islands  

- GDP China is applied to  

    Aksai Chin (administered by China. Claimed by  

India), China, Taiwan  

- GDP France is applied to  

    Saint Pierre and Miquelon, France, Martinique,  

French Guyana, Wallis and Futuna Islands, Reunion, Europa Islands, French  

Southern Territories  

74: no match found for GDP country name Gibraltar  

- GDP India is applied to  

    Kashmir and Jammu (claimed by India and Pakistan),  

India  

93: no match found for GDP country name Isle of Man  

- GDP Israel is applied to  

    Golan Heights, Israel, Gaza Strip  

104: no match found for GDP country name Kosovo  

116: no match found for GDP country name Macao SAR, China  

121: no match found for GDP country name Maldives  

131: no match found for GDP country name Monaco  

133: no match found for GDP country name Montenegro  

- GDP Morocco is applied to  

    Morocco, Western Sahara  

- GDP New Zealand is applied to  

    Cook Islands, New Zealand  

- GDP Norway is applied to  

    Jan Mayen, Svalbard, Norway  

- GDP Russia is applied to
```

```
Russia, Kuril Islands (administered by Russia.
Claimed by Japan)
171: no match found for GDP country name Sint Maarten (Dutch part)
177: no match found for GDP country name South Sudan
- GDP Turkey is applied to
    Turkey, Cyprus Turkish part
- GDP United Kingdom is applied to
    United Kingdom, Virgin Islands (British), Anguilla,
    Montserrat, British Indian Ocean Territory, Saint Helena, Pitcairn, Falkland
    Islands (Malvinas), South Georgia and South Sandwich Islands
- GDP United States (USA) is applied to
    United States (USA), Wake Island, Palmyra Atoll,
    Jarvis
```

MATLAB call:

```
GDP = climada_GDP_check_countrynamess(GDP, borders, silent_mode)
```

Output

```
GDP country name matched for 239 countries within climada world map.
No GDP information available for following 4 countries (within climada
worldmap):
- Antarctica
- Bouvet Island
- Nauru
- Niue
```

### 3b) Future projection of GDP

GDP development data is available from the International Monetary Fund, World Economic Outlook Database 2012 and from the European Central Bank

<http://www.imf.org/external/ns/cs.aspx?id=28>

<http://www.imf.org/external/pubs/ft/weo/2012/01/weodata/index.aspx>

<http://www.ecb.int/home/html/index.en.html>

Gross domestic product, current prices, U.S. dollars, GDP growth until 2010, 2011

<http://data.worldbank.org/indicator/NY.GDP.MKTP.KD.ZG>



Figure 9: GDP growth data from world bank.

GDP projection data from 2010 to 2017 can be read from the excel file (GDP evolution from International Monetary Fund (IMF)), that is saved in

[climada\\_additional\GDP\\_entity\data\World\\_GDP\\_2010\\_2017.xlsx](http://climada_additional/GDP_entity/data/World_GDP_2010_2017.xlsx)

MATLAB call:

```
GDP_forecast = climada_GDP_read(xlsfilename, special_cases_on,
check_names, save_on, silent_mode)
climada_GDP_read('World_GDP_constant_2000_2017.xls')
```

.country_names	<i>216x1 cell, list with all country names, e.g. Afghanistan, Albania etc.</i>
.year	<i>1x51 double, with years from 2000 to 2017</i>
.value	<i>216x51 double, GDP value per year and country</i>
.description	<i>constant GDP in national currency</i>
.comment	<i>IMF GDP data, from year 2010 to 2017</i>
.country_borders_index	<i>216x1 cell, index that links GDP with border mask structure (see next chapter), e.g. GDP.country_borders_index(1) = 52; country number 1 in GDP list is Afghanistan and is on position 51 in the borders structure. One country in the GDP list can index to multiple countries in the border mask structure, for instance GDP India (position 88) refers to border masks position 55 (Kashmir and Jammu (claimed by India and Pakistan) and 100 (India).</i>

We extrapolate a future GDP based on the GDP forecast data by IMF (current price, national currency) with a second order polynomial. A scaleup factor is calculated on the basis of a linear extrapolation between today and the specified year in the future.




Figure 10: GDP development forecast from 2012 to 2030 for countries in North America. Forecast data until 2017 is available from IMF (GDP current prices, national currency).

## 4. Countries: Use polygons and rasters as helper files

The climada world map is based on the text file `world_50m.gen` that is stored in the system folder `..\climada\data\system\world_50m.gen`

A matlab variable is generated the first time from the gen-file while running the code `climada_plot_world_borders`. The mat-file is a structure variable and consists a list of country names, and according polygons. `..\climada\data\system\world_50m.mat`

**MATLAB call:**

```
[varargout]= climada_plot_world_borders(linewidth,check_country,
map_border_file, keep_boundary, country_color)
```

We create a raster for every country polygon on given lat-long-range and cellsize, size 1680 x 4320, filled with zeros and ones (one for within country, zero for out of country). This resolution corresponds roughly to 10 km. This takes a bit of time. For ~50km resolution (raster size 336 x 864) the function is a lot faster (roughly 2min). All country rasters are saved as `border_mask.mat` structure in `..\climada_additional\GDP_entity\data\border_mask_10km.mat` or `..\climada_additional\GDP_entity\data\border_mask_50km.mat`

**MATLAB call:**

```
border_mask= climada_polygon2raster(borders,raster_size,save_on)
climada_polygon2raster('',[336 864])

.mask      243 matrices (for each country) masking 1 for within country
           and zero for out of the country
.name      Name of each country
.world_mask All countries within one mask (1 for land, 0 for sea)
.lon_range Longitudinal range of masks, e.g. [-180 180]
.lat_range Latitudinal range of masks, e.g. [-65 75]
.resolution_x Resolution in x direction in degree, e.g. 0.0833°, corresponds
           to ~10km
.resolution_y Resolution in y direction in degree, e.g. 0.0833°, corresponds
           to ~10km
```




Figure 11: Raster of Canada polygons.  
Saved in

`border_mask.mask{country_Canada}`




Figure 12: Mask (zeros and ones) of the whole world.  
Resolution roughly 50km, 336\*864 pixels.

## 5. Night light intensities as a proxy for spatial distribution of values within a country: Linear or nonlinear relationship between light intensity and asset value

We use the following function to identify the night light intensities within a country (given through the country mask) and to assign the values according to night light intensities.

The relationship between light intensities and asset values can be linear or nonlinear and in the latter case can be inputted through the factor pp. The factor pp describes the variables of a second order polynomial function without y-indent, such as  $y = pp(1)*x^2 + pp(2)*x$ , where y is the asset value and x is the night light intensities (a value between 1 and 63). To use a linear relationship set pp to  $pp = [1 0]$ . pp is optional. If pp is not given, a standard nonlinear relationship is used, which is  $y = -0.0817*x + 0.0172*x^2$ . The sum of all values distributed within a country is set to 100, i.e.

```
>> sum(full(values_distributed.values(:)))
ans =
100.0000
```

Like this, the values can easily be up scaled to any required value, such as GDP for a given year or any other value.

MATLAB call:

```
[values_distr    climada_night_light_to_country(country_name,           pp,
ibuted, pp]=      night_light,   borders,   border_mask,   check_figure,
                  check_printplot, save_on, silent_mode)
                  climada_night_light_to_country('Switzerland',[1 0])
                  climada_night_light_to_country('Switzerland')
```


values\_distributed is a structure with the following fields

.values	<i>1680x4320 sparse matrix, distribution of values within the range indicated in lon_range and lat_range, ~10km resolution</i>
.lon_range	<i>Longitudinal range of masks, e.g. [-180 180]</i>
.lat_range	<i>Latitudinal range of masks, e.g. [-65 75]</i>
.resolution_x	<i>Resolution in x direction in degree, e.g. 0.0833°, corresponds to ~10km</i>
.resolution_y	<i>Resolution in y direction in degree, 0.0833°, corresponds to ~10km</i>
.comment	<i>e.g. nonlinear function, <math>y = 0.0000*x^0 + -0.0817*x^1 + 0.0172*x^2</math></i>

And in there the function transforms the night lights to asset values.


MATLAB call:

```
[values_out, pp]= climada_nightlight_nonlinear_transformation
                    (values_in, pp, check_figure, check_printplot)
```




**Figure 13:** Distributed values within Switzerland on the basis of a linear relationship between night light intensities and asset values.

The sum of all values corresponds to 100. This value distribution forms the basis for a given scaling, e.g. based on a country's GDP.



**Figure 14:** Distributed values within Switzerland on the basis of a nonlinear relationship between night light intensities and asset values.

The sum of all values corresponds to 100. This value distribution forms the basis for a given scaling, e.g. based on a country's GDP.



**Figure 15:** Second order polynomial transformation function (without y-indent) to define the relationship between night light intensities and asset values.




Figure 16: GDP distributed according to night lights within one specific country, zoom to USA and Mexico. Bright spots are redder in the US than in Mexico, because Mexico has in total a lower GDP.




Figure 17: GDP distributed according to night lights within one specific country, zoom to Europe. See bright spot differences in Portugal versus Spain, or Germany versus Poland.




Figure 18: Night time lights in Europe.




Figure 19: GDP per country in 2010.

## 6. Create the centroids

### 6a) Define the resolution

In order to create centroids at a requested resolution the value distribution matrix needs to be downscaled accordingly.

To downscale the matrix to a required resolution we use the following function. The `high_resolution_matrix` is a matlab structure as described in `values_distributed`, namely with the fields `.values`, `.lon_range`, `.lat_range`, `.resolution_x`, `.resolution_y`. Specification is a string, to determine the value for the downsampled resolution matrix, with either summing up all values with the keyword '`sum`', taking the average with keyword '`average`' or taking the most counted value with keyword '`unique`'.

MATLAB call:

```
[low_resolution    climada_resolution_downscale      (high_resolution_matrix,
_matrix, X, Y,   resolution_km, specification)
resolution_km]=
climada_resolution_downscale(values_distributed, 50, 'sum')
```

`low_resolution_matrix` is a structure with the following fields

<code>.values</code>	<i>336x864 sparse matrix, distribution of values within the range indicated in lon_range and lat_range, ~50km resolution</i>
<code>.lon_range</code>	<i>Longitudinal range of masks, e.g. [-180 180]</i>
<code>.lat_range</code>	<i>Latitudinal range of masks, e.g. [-65 75]</i>
<code>.resolution_x</code>	<i>Resolution in x direction in degree, e.g. 0.4167°, corresponds to ~50km</i>
<code>.resolution_y</code>	<i>Resolution in y direction in degree, 0.4167°, corresponds to ~50km</i>
<code>.comment</code>	<i>e.g. nonlinear function, <math>y = 0.0000*x^0 + -0.0817*x^1 + 0.0172*x^2</math>, 46km, sum</i>

`X` and `Y` are helper matrices containing the longitude and latitude information, respectively for plotting the `low_resolution_matrix`.

### 6b) Cut out coastal areas (hollow out) and expand with a buffer zone

We create a buffer zone around the specified country in order to have centroids also in the oceans (e.g. to create nice wind field plots). Additionally the country area can be hollowed out to keep only the coastal areas. The buffer zone is characterized by the number of pixels into the sea. The coastal region to be kept is defined through the number of pixels as well. If no cut out is needed `no_pixel_hollow` can be set to 0.

MATLAB call:

```
matrix_buffer = climada_mask_buffer_hollow (matrix, no_pixel_buffer,
no_pixel_hollow, border_mask, check_figure,
check_printplot, printname, cbar_label)
climada_mask_buffer_hollow(border_mask.mask{45},2,5)
```

`matrix_buffer` is a matrix masking 1 for on land, zero for sea, 2 for buffer.




Figure 20: Mexico (`no_pixel_hollow = 0`), and buffer zone of about 80 km (8 pixels) around Mexico (grey).

Figure 21: Coastal Mexican areas (within 150 km from the coast, 15 pixels, turquoise) and buffer zone of about 80 km (8 pixels) around Mexico (grey).




Figure 22: Entire world map with buffer of about 200 km (20 pixels).

### 6c) Create the centroids

This function creates the centroids on coastal land areas, within the buffer zone at the coast and on a coarser level further away, such as on sea and more distant from the coast. The resolution for coastal areas and buffer zone corresponds to the required input resolution. The regular grid has a five times coarser resolution, e.g. 50 km for resolution of coastal areas on 10 km.

MATLAB call:

```
centroids = climada_matrix2centroid(matrix_buffer, lon_range, lat_range,
                                     country_name)
climada_resolution_downscale(values_distributed, 50, 'sum')
```

`low_resolution_matrix` is a structure with the following fields

<code>.values</code>	336x864 sparse matrix, distribution of values within the range indicated in <code>lon_range</code> and <code>lat_range</code> , ~50km resolution
<code>.lon_range</code>	Longitudinal range of masks, e.g. [-180 180]
<code>.lat_range</code>	Latitudinal range of masks, e.g. [-65 75]
<code>.resolution_x</code>	Resolution in x direction in degree, e.g. 0.4167°, corresponds to ~50km
<code>.resolution_y</code>	Resolution in y direction in degree, 0.4167°, corresponds to ~50km
<code>.comment</code>	e.g. nonlinear function, $y = 0.0000*x^0 + -0.0817*x^1 + 0.0172*x^2$ , 46km, sum

`X` and `Y` are a helper matrices containing the longitude and latitude information, respectively for plotting the `low_resolution_matrix`.

The centroids can be visualized on a map.

MATLAB call:

```
fig = climada_plot_centroids(centroids, country_name, check_printplot,  
printname)
```




Figure 23: Centroids in entire Mexico (turquoise, 10 km resolution), buffer zone (grey, 10 km resolution) and further away (on sea or in another country, orange, 50 km resolution).




Figure 24: Centroids in coastal Mexico (turquoise, 10 km resolution), buffer zone (grey, 10 km resolution) and further away (on sea or too far from the coast, orange, 50 km resolution).

## 7. Create the entity

### 7a) Base entity: assets sum up to 100 for a given country

The entity file contains all asset, vulnerability and adaptation measure data for a certain country or region. This is essentially an excel file with four tabs. As described in chapter 0, in the base entity, the sum of all assets within the entire country corresponds to 100. If only coastal areas are taken (hollowed out matrix), the sum of coastal assets is smaller than 100. This presents the basis for scaling up assets to any given value, such as the country's GDP for any given year or future year.

- Tab "assets": List of assets with longitude, latitude, asset value, damage function ID
- Tab "damagefunctions": A damage function describes how the asset responds to a given wind intensity. The damage function consists of two curves which put wind intensity in relation to mean damage degree and percentage of affected assets for a certain class of assets. There can be different damage functions depending on type and age of building.
- Tab "measures": Adaptation measures are available to avert todays and future risks. These can range from mangroves conservation, beach nourishment, to wind-resistant roofs and cross-bracing of houses, sandbags, construction of a seawall and risk transfer. This tab features a list of possible measures and its associated cost and the impacts on damage reduction. The impact of a measures is parameterized through the component of hazard reduction (e.g. mangroves reduce wind speed and wave heights), vulnerability reduction (e.g. wind-resistant roofs can withstand all damage up to a certain wind speed).
- Tab "discount": This tab contains the discount information, such as discount rate per year from today until the future (e.g. 2030 or 2050). This is used for discounting future loss reductions achieved through specific adaptation measures.

The base entity is saved as an excel file and is used in climada as a matlab structure. As in this module we create the assets only (longitude, latitude, asset values) of a given country or region, we use a wildcard entity with dummy values for the tabs damagefunctions, measures and discount. Based on this wildcard entity, we add the assets (location and values) to the entity matlab structure. In order to map a single asset to the closest calculation centroid, the assets are encoded to centroids. In core climada there is the function `climada_assets_encode`, that encodes the assets on the basis of the hazard, however here we encode the assets on the basis of the centroids.

MATLAB call:

```
entity_base = climada_entity_base_assets_add(values_distributed,centroids,  
country_name, matrix_hollowout, X, Y, hollow_name)  
  
.assets .excel_file_name  
.Latitude  
.Longitude  
.Value  
.Deductible  
.Cover  
.DamageFunID  
.centroid_index  
.hazard  
.filename  
.DamageFunID  
.Intensity  
.MDD  
.PAA  
.MDR  
.name  
.color  
.cost  
.hazard_intensity_impact  
.hazard_high_frequency_cutoff
```

```
.MDD_impact_a  
.MDD_impact_b  
.PAA_impact_a  
.PAA_impact_b  
.damagefunctions_map  
.risk_transfer_attachement  
.risk_transfer_cover  
.color_RGB  
.damagefunctions_mapping  
.discount           .filename  
                      .yield_ID  
                      .year  
                      .discount_rate
```

This function combines the following functions

To read the wildcard entity from excel file

MATLAB call:

```
entity_base = climada_entity_read_wo_assets(entity_filename)
```

To encode assets from the distributed value matrix

MATLAB call:

```
[entity_base.assets, climada_assets_encode_centroids(assets, centroids)  
centroids]=
```

Finally the assets within the entity can be visualized on a map.

MATLAB call:

```
fig = climada_plot_entity_assets(entity,centroids,  
country_name, check_printplot, printname)
```




Figure 25: Distribution of assets in Mexico (coastal areas and inland areas) on a 10 km resolution. The sum of all values corresponds to 100.




Figure 26: Distribution of assets in Mexican coastal areas only, on a 10 km resolution. The sum of all values corresponds to 39.9 as the inland areas omitted.


## 7b) Entity with assets based on the country's GDP today

The assets need to be scale up from the base value of 100 (or less in the case of coastal areas only) to sum up to a country's GDP. This is done with the following function:


MATLAB call:

```
entity = climada_entity_GDP(entity_base,GDP,year_start,centroids,
borders, check_figure, check_printplot)
climada_entity_GDP(entity_100,GDP,2010,centroids)
```

Note that the distribution of assets (look at the color scheme) is the same as in Figure 25 and Figure 26. Only the sum of all assets has changed.



*Figure 27: Distribution of assets in Mexico, on a 10 km resolution. The sum of all asset values corresponds to the Mexican GDP in 2014.*



*Figure 28: Distribution of assets in Mexican coastal areas only, on a 10 km resolution. The sum of all values corresponds to 39.9% of the Mexican GDP in 2014 (inland areas omitted).*

### 7c) Future entity with assets based on the country's future GDP (e.g. 2030)

We use the GDP projection data from IMF for the years 2010 to 2017, from where we derive linear extrapolation formulas (see GDP chapter). Based on the linear extrapolation we project GDP for any given year in the future, e.g. 2030 or 2050. This is also used if the start year is after 2010, as the GDP data from the World Bank is only available up to 2010 in the excel sheet.

This following function scales the assets up based on GDP growth between two given periods. It includes the reading of GDP projection data, the extrapolation and the up scaling of the assets in the entity structure.

MATLAB call:

```
entity = climada_entity_scaleup_GDP(entity,GDP_forecast,year_start,  
centroids, borders, check_figure, check_printplot)
```

And in there the function `climada_entity_scaleup_factor` is used to scale up all assets within a given country with a given multiplicator.

MATLAB call:

```
entity = climada_entity_scaleup_GDP(entity,GDP_forecast,year_start,  
centroids, borders, check_figure, check_printplot)
```

### 7d) Select a specific region within a country

We can select a specific region within a country, and carve out assets and centroids within that region. The region can be defined through a polygon (vector), or is defined interactively by using the mouse.

MATLAB call:

```
[centroids, climada_cut_out_GDP_entity(entity, centroids, polygon)  
entity, polygon]=
```

`polygon` is a vector that characterizes the polygon, ie, `polygon(:,1)` are the longitudes, `polygon(:,2)` are the latitudes.




Figure 29: Define the region that is required with the mouse (see red lines).




Figure 30: Assets of within selected regions are still visible. All assets outside of the polygon are in grey.

## 8. Appendix

### 8a) Downscale resolution of original night light tif in ArcGIS




Figure 31: Night time lights in ArcGIS, 2010.

- Open georeferenced image in ArcGIS and save ArcGIS file in  
[\\CHRB1065.CORP.GWPNET.COM\homes\X\S3BXXW\Documents\lea\cc\\_SwissRe\\_modeling\climada\\_cc\\_i mpact\data\globalGDP\night\\_light\20120713\\_night\\_time\\_light.mxd](\\CHRB1065.CORP.GWPNET.COM\homes\X\S3BXXW\Documents\lea\cc_SwissRe_modeling\climada_cc_i mpact\data\globalGDP\night_light\20120713_night_time_light.mxd)
- Resample in ArcToolbox (search in ArcToolbox) from original resolution ~1km to ~10km cellsize




Figure 32: Resample details in ArcToolbox ArcGIS.




Figure 33: Resample cell resolution from ~1km (left) to ~10km (right).

- Save stable nights lights as image in
- [\\CHRB1065.CORP.GWPNET.COM\homes\X\S3BXXW\Documents\lea\cc\\_SwissRe\\_modeling\climada\\_cc\\_i mpact\data\system\night\\_light\\_2010\\_10km.png](\\CHRB1065.CORP.GWPNET.COM\homes\X\S3BXXW\Documents\lea\cc_SwissRe_modeling\climada_cc_i mpact\data\system\night_light_2010_10km.png)

## 8b) Night light and GDP relationship

### 8.b.1. Literature

- U.S. Department of Commerce, Bureau of Economic Analysis  
<http://www.bea.gov/regional/index.htm#gsp>
- Global estimates of market and non-market values derived from nighttime satellite imagery, land cover, and ecosystem service valuation, Sutton and Costanza, 2002, Ecological Economics.  
<http://www.sciencedirect.com/science/article/pii/S0921800902000976>

$$\ln(\text{GDP}) = \alpha + \beta \ln(\text{night light})$$

$$\alpha = -4.25$$

$$\beta = 1.05$$




Figure 34: First order polynomial fit to natural logarithmic relation between night light and GDP.

- Using luminosity data as a proxy for economic statistics, Chen and Nordhaus, 2011, PNAS.  
<http://www.pnas.org/content/108/21/8589.full.pdf+html>  
[http://www.nsf.gov/discoveries/disc\\_summ.jsp?cntn\\_id=119737](http://www.nsf.gov/discoveries/disc_summ.jsp?cntn_id=119737)
- Geographically based Economic data (G-Econ)  
<http://gecon.yale.edu/usa>
- Night Lights and Economic Activity in India: A study using DMSP-OLS night time images, Bhandari and Roychowdhury, 2011  
[http://www.google.ch/url?sa=t&rct=j&q=&esrc=s&frm=1&source=web&cd=4&cad=rja&ved=0CF0QFjAD&url=http%3A%2F%2Fusymposia.upm.my%2Findex.php%2FAPAN\\_Proceedings%2F32nd\\_APAN%2Fpaper%2Fdownload%2F137%2F96&ei=HUUzUNfeCcrStAaf6lCoAg&usg=AFQjCNG5fRnfSoK1j0SgnzcUIO1u1m-84w&sig2=3RiTp-r650dRW0t7V58Cpg](http://www.google.ch/url?sa=t&rct=j&q=&esrc=s&frm=1&source=web&cd=4&cad=rja&ved=0CF0QFjAD&url=http%3A%2F%2Fusymposia.upm.my%2Findex.php%2FAPAN_Proceedings%2F32nd_APAN%2Fpaper%2Fdownload%2F137%2F96&ei=HUUzUNfeCcrStAaf6lCoAg&usg=AFQjCNG5fRnfSoK1j0SgnzcUIO1u1m-84w&sig2=3RiTp-r650dRW0t7V58Cpg)
- Night time lights averaged by admin level 2 areas – indicator of GDP  
<http://www.edenextdata.com/?q=content/night-time-lights-averaged-admin-level-2-areas-indicator-gdp-0>
- Shedding Light on the Global Distribution of Economic Activity, Gosh et al., 2010, The Open Geography Journal.  
[http://www.ngdc.noaa.gov/dmsp/pubs/Ghosh\\_TOGEOGJ.pdf](http://www.ngdc.noaa.gov/dmsp/pubs/Ghosh_TOGEOGJ.pdf)