Difference Equations

2.1.1 Compound Interest

2.1.2 Loan Repayment

2.1.3 Gambler's Ruin

2.2.2 Exponential Population Growth

2.2.3 Average Lifespan

2.2.★ Rabbit Populations

# 2.2.4 Nonlinear Population Models

**Goal.** Compare nonlinear Differential Equations vs Difference Equations.

**Goal.** Compare nonlinear Differential Equations vs Difference Equations.

Consider the continuous Logistic Population Model

$$ho'(t) = R
ho(t)\left(1 - rac{
ho(t)}{K}
ight)$$

**Goal.** Compare nonlinear Differential Equations vs Difference Equations.

Consider the continuous Logistic Population Model

$$ho'(t) = R
ho(t)\left(1 - rac{
ho(t)}{K}
ight)$$

and the discrete Logistic Population Mode

$$p_{n+1} - p_n = Rp_n \left(1 - \frac{p_n}{K}\right)$$

**Goal.** Compare nonlinear Differential Equations vs Difference Equations.

Consider the continuous Logistic Population Model

$$ho'(t) = R
ho(t)\left(1 - rac{
ho(t)}{K}
ight)$$

and the discrete Logistic Population Mode

$$p_{n+1} - p_n = Rp_n \left(1 - \frac{p_n}{K}\right)$$

Using 
$$x_n = \frac{R}{1+R} \frac{p_n}{K}$$
 and  $\mu = 1+R$ , obtain

$$x_{n+1} = \mu x_n (1 - x_n).$$

**Goal.** Compare nonlinear Differential Equations vs Difference Equations.

Consider the continuous Logistic Population Model

$$ho'(t) = R
ho(t)\left(1 - rac{
ho(t)}{K}
ight)$$

and the discrete Logistic Population Mode

$$p_{n+1} - p_n = Rp_n \left(1 - \frac{p_n}{K}\right)$$


Using 
$$x_n = \frac{R}{1+R} \frac{p_n}{K}$$
 and  $\mu = 1+R$ , obtain

$$x_{n+1} = \mu x_n (1 - x_n).$$

**2** Consider the influenza virus (R = 3). Then  $\mu = 4$ .


We can run the difference equation for  $x_n$  with different initial conditions:

We can run the difference equation for  $x_n$  with different initial conditions:




 $x_0 = 0.1$ 

We can run the difference equation for  $x_n$  with different initial conditions:




$$x_0 = 0.1$$



$$x_0 = 0.101$$

compare

We can run the difference equation for  $x_n$  with different initial conditions:



What happens to the continuous model with the same initial conditions?

$$x_{n+1} = \mu x_n (1 - x_n)$$

$$\mu = 1$$
 , 2 , 4 , 8 ?

$$x_{n+1} = \mu x_n (1 - x_n)$$


$$\mu = 1 , 2 , 4 , 8 ?$$

- $\circ$  Allow  $\mu \in \mathbb{C}$ .
- O Which values of  $\mu \in \mathbb{C}$ , for which  $x_n$  doesn't go to infinity?

$$x_{n+1} = \mu x_n (1 - x_n)$$

$$\mu = 1 , 2 , 4 , 8 ?$$


- $\circ$  Allow  $\mu \in \mathbb{C}$ .
- O Which values of  $\mu \in \mathbb{C}$ , for which  $x_n$  doesn't go to infinity?



$$x_{n+1} = \mu x_n (1 - x_n)$$

$$\mu = 1$$
 , 2 , 4 , 8 ?

- $\circ$  Allow  $\mu \in \mathbb{C}$ .
- O Which values of  $\mu \in \mathbb{C}$ , for which  $x_n$  doesn't go to infinity?



$$x_{n+1} = \mu x_n (1 - x_n)$$

$$\mu = 1$$
 , 2 , 4 , 8 ?

- $\circ$  Allow  $\mu \in \mathbb{C}$ .
- O Which values of  $\mu \in \mathbb{C}$ , for which  $x_n$  doesn't go to infinity?

