

Supplementary Information

Methods

1 Observational reanalysis datasets

2 Four independent global ocean reanalysis products are analysed. They were selected to span the four
3 mainstream data-assimilation paradigms in the global-ocean reanalysis ecosystem – 3D-Var FGAT (ORAS5),
4 reduced-order Kalman filter with 3D-Var bias correction (GLORYS12V1), ensemble optimal interpola-
5 tion (SODA 3.15.2), and 4D-variational adjoint state estimation (ECCO-V4r4) – and together cover the
6 eddy-resolving ($1/12^\circ$) to coarse-resolution ($1/2^\circ$) range used in CMIP-class climate analysis. This is the
7 same four-product canonical ensemble adopted by the headline emergent-constraint study [16], ensuring
8 that our mechanism decomposition is directly comparable to that constraint without product-selection
9 ambiguity. Adding a fifth product from any of the same paradigms (e.g. ORAS4, C-GLORS, FOAM)
10 would not break the velocity-vs-salinity dichotomy because such siblings share systematic biases with
11 their corresponding family member; a complementary GECCO3 (Hamburg) 4D-Var run would be the
12 most informative future addition because it would test whether the ECCO null trend is generic to adjoint
13 optimisation. The four products together represent the currently available space of observationally-
14 constrained three-dimensional ocean state estimates at 34.5°S .

15 **ORAS5** [23]. Ocean Reanalysis System 5 from ECMWF. Native grid: ORCA025 tripolar (nominal
16 $\sim 1/4^\circ$, ~ 25 km at the equator and ~ 14 km at 34.5°S after the $\cos\phi$ contraction), 75 vertical levels
17 with 1-m surface resolution. NEMOVAR 3D-Var FGAT data-assimilation scheme ingesting sub-surface
18 temperature/salinity profiles, sea-ice concentration, and along-track altimetric sea-level anomalies. At-
19 mospheric forcing: ERA-40 (1958–1978) transitioning to ERA-Interim/ERA5 (1979–present). Record
20 used: 816 monthly means, 1958–2025.

21 **GLORYS12V1** [12]. Copernicus Marine Service/Mercator Ocean global ocean reanalysis at $1/12^\circ$
22 (~ 8 km, eddy-resolving), 50 vertical levels. A reduced-order Kalman filter (SAM2) jointly assimilates
23 along-track altimetry, satellite sea-surface temperature, sea-ice concentration, and in-situ CTD and Argo
24 temperature/salinity profiles, supplemented by 3D-Var bias correction. Atmospheric forcing: ERA5.
25 Record used: 396 monthly means, 1993–2025 (satellite-altimetry era).

26 **SODA 3.15.2** [2]. Simple Ocean Data Assimilation 3 from the University of Maryland. Native grid:
27 MOM5 $\sim 1/2^\circ$ (regrided to a uniform 0.5° product on the UMD mirror), 50 vertical levels. Assimilation
28 scheme: ensemble optimal interpolation with a 5-day analysis cycle; the downloadable output is a sequence
29 of 5-day snapshots. Atmospheric forcing: MERRA-2. We sample four 5-day snapshots per year (mid-
30 month of January, April, July, October) to approximate the annual mean, consistent with previous
31 applications of the same product to overturning-transport diagnostics. Record used: 43 annual means,
32 1980–2022.

33 **ECCO-V4r4** [6]. NASA/JPL Estimating the Circulation and Climate of the Ocean, version 4 release
34 4. Native grid: LLC90 Lat-Lon-Cap (regrided to a uniform 0.5° monthly product), 50 vertical levels.
35 ECCO is an adjoint-based least-squares state estimate (4D-variational) that iteratively adjusts initial
36 conditions, surface fluxes, and interior mixing to minimise the misfit between the model trajectory and
37 the full suite of altimetry, Argo, satellite SST/SSH, and mooring observations over its full time span.
38 Atmospheric forcing (post-iteration): an ECCO-adjusted ERA-family atmospheric analysis. Record used:
39 26 annual means, 1992–2017.

Direct-hydrography anchors. Three published overturning freshwater-transport estimates at 34.5°S from direct in-situ observations are used as independent anchor points in Fig. 1: Garzoli and Matano [7] (central year 2005, value -0.10 ± 0.10 Sv); Meinen et al. [14] (central year 2015, value -0.09 ± 0.05 Sv); and Kersalé et al. [10] (central year 2015, value -0.17 ± 0.07 Sv). Together with Arumí-Planas et al. [1]’s -0.15 ± 0.09 Sv from 49 XBT transects, they bracket the reanalysis ensemble from the low side.

CMIP6 ensemble

CMIP6 meridional-velocity (vo) and salinity (so) sections at 34.5°S are obtained for every available model with coverage of both the `historical` experiment (required period 1950–1980) and the `ssp585` experiment (required period 2080–2100). The native grid of each model is used (i.e. no regridding), because regridding distorts the mechanistic decomposition. Fifteen models satisfy both criteria with accessible section files: CESM2, CMCC-CM2-SR5, CNRM-CM6-1, CanESM5, FGOALS-g3, FIO-ESM-2-0, GFDL-CM4, GISS-E2-1-G, HadGEM3-GC31-LL, IPSL-CM6A-LR, MIROC6, MPI-ESM1-2-HR, MPI-ESM1-2-LR, NESM3, UKESM1-0-LL. Three models are excluded: NorESM2-LM and NorESM2-MM because their ocean component is on isopycnal (density) coordinates and cannot be integrated in depth-space without an additional remap that would confound the decomposition; and ACCESS-CM2 because its accessible `ssp585` run does not extend to 2100. All models use the `r1i1p1f1` realisation.

A 13-model subset with `piControl` access is used for the internal-variability null test: ACCESS-CM2, CESM2, CMCC-CM2-SR5, CNRM-CM6-1, CanESM5, GISS-E2-1-G, HadGEM3-GC31-LL, IPSL-CM6A-LR, MIROC6, MPI-ESM1-2-HR, MPI-ESM1-2-LR, NESM3, UKESM1-0-LL.

Ocean component reference per weakening model. Table 1 lists the ocean and atmospheric components of each CMIP6 weakening model, grouped by mechanism class as classified from the centennial F_{ovS} decomposition. We provide this as an audit reference rather than a causal claim: the discriminating physics between the velocity-dominant and salinity-dominant clusters appears to be an emergent feature of the full coupled response (Southern-Hemisphere wind sensitivity vs NADW-buoyancy sensitivity, see main-text Discussion), not a single ocean-component feature. NEMO, in particular, appears in both classes (NESM3 velocity-dominant; CNRM-CM6-1, HadGEM3-GC31-LL, UKESM1-0-LL salinity-dominant), confirming that ocean code identity is not the discriminant.

Table 1: Ocean numerical infrastructure per weakening CMIP6 model, grouped by mechanism class. Vertical mixing schemes: KPP = K-Profile Parameterization [11]; PP = Pacanowski and Philander [15]; TKE = Turbulent Kinetic Energy closure (NEMO standard); ePBL = energetic Planetary Boundary Layer [19]; NIW = Near-Inertial Wave parameterization. Mesoscale eddy: GM = Gent and McWilliams [8] thickness-diffusion + Redi [18] isopycnal diffusion; MEKE = Mesoscale-Eddy Kinetic Energy parameterization [9]. The table demonstrates that *no single ocean parameterization* covaries with mechanism class — e.g. TKE-NEMO appears in BOTH camps (NESM3 v-dom; CNRM, HadGEM, UKESM s-dom), and PP-based schemes appear in BOTH (MPI-ESM1-2 v-dom; FGOALS-g3 s-dom). The wind-vs-buoyancy split discussed in the main text is therefore an *emergent feature of the full coupled response* (atmospheric SH-westerly response, sea-ice/AMOC coupling, NADW formation sensitivity) rather than a property of any single ocean numerical scheme.

Class	Model	Ocean (res.)	Vert. mixing	Mesoscale eddy	Atmosphere
v-dom	CESM2	POP2 (1°)	KPP + Langmuir	GM + Redi	CAM6
v-dom	FIO-ESM-2-0	POP2 (1°)	KPP	GM + Redi	CAM5
v-dom	MPI-ESM1-2-LR	MPIOM (1.5°)	PP + wind decay	GM + Redi	ECHAM6.3
v-dom	MPI-ESM1-2-HR	MPIOM (0.4°)	PP + wind decay	GM + Redi	ECHAM6.3
v-dom	NESM3	NEMO 3.4 (1°)	TKE	GM + Redi	ECHAM6.3
s-dom	CNRM-CM6-1	NEMO 3.6 (1°)	TKE	GM + Redi	ARPEGE 6.3
s-dom	FGOALS-g3	LICOM3 (1°)	PP	GM + Redi	GAMIL2
s-dom	GFDL-CM4	MOM6 (0.25°)	ePBL + KPP shear	MEKE + GM	AM4
s-dom	HadGEM3-GC31-LL	NEMO 3.6 ORCA1 (1°)	TKE	GM + Redi	UM-GA7.1
s-dom	MIROC6	COCO 4.9 (1°)	NIW + Tsujino	GM + Redi	MIROC AGCM
s-dom	UKESM1-0-LL	NEMO 3.6 ORCA1 (1°)	TKE	GM + Redi	UM-GA7.1
mixed	GISS-E2-1-G	GISS Ocean (1°)	KPP-like	GM + Redi	GISS ModelE

Ensemble size used per analysis. Different downstream analyses retain different subsets of the 15-model decomposition ensemble:

- *Mechanism decomposition + tiebreaker scatter* (Figs. 1b, 2a of the main text): all 15 models. Three models are flagged *increasing* (CMCC-CM2-SR5, CanESM5, IPSL-CM6A-LR) and excluded from the mechanism-conditional projection because their forced F_{ovS} change is positive; this leaves **12 weakening models** that drive the headline 10-percentage-point gap (Figs. 2b, 2c).
- *Bistable-only subset* (Fig. 3): 6 models have $\overline{F_{\text{ovS}}}(t_1) < 0$; of those, 2 are *increasing* and are dropped, leaving $n = 4$ **weakening + bistable** models (CNRM-CM6-1, MIROC6 salinity-dominant; MPI-ESM1-2-HR, NESM3 velocity-dominant).
- *piControl null* (Fig. 4c): **13 models** (the subset listed above), with up to 200 random non-overlapping 30-year window pairs per model.
- *Lead-lag cross-correlation* (Fig. 6a) and *emergent constraint regression* (Fig. 6b): **16 models**. Both diagnostics use only the 1850-2024 historical+ssp585 window for the lead-lag and the 2000-2024 / 2030-2040 windows for the emergent constraint, which means ACCESS-CM2 – excluded from the decomposition because its ssp585 run does not reach 2100 – can be retained here. We re-add it to maximise statistical power on the lead-lag cross-correlation function and on the regression slope.

This per-analysis bookkeeping is reflected in each figure’s panel-internal $n =$ labels and in the captions of the main text.

Section extraction and grid metrics

At each time step of each product we extract the zonal-depth section at the gridded latitude closest to -34.5° . Where the ocean-model grid is staggered (e.g. velocity at U-points and salinity at T-points on the ORAS5 NEMO grid), the two fields are extracted on their native grids and aligned by truncation to the common Atlantic-longitude mask.

The Atlantic basin at 34.5°S is defined by the longitude range $[-70^\circ, 20^\circ]$ in the $[-180^\circ, 180^\circ]$ convention (bounds chosen inclusively to cover from the Argentine coast to the African coast while excluding residual Agulhas leakage; the small $< 1\%$ contribution from the strictly-classified Drake-Passage edge does not alter the results).

The zonal grid spacing at each Atlantic U-point is

$$\Delta x_i = |\Delta \lambda_i| R_\oplus \cos \phi,$$

with $\Delta \lambda_i$ the local longitudinal grid spacing (after longitude wrap-around correction), $R_\oplus = 111$ km per degree at the equator, and $\phi = -34.5^\circ$. Vertical cell thicknesses Δz_k are taken directly from the product’s vertical coordinate (with the CESM-family POP grid converted from centimetres to metres).

F_{ovS} computation

We follow the definition of de Vries and Weber [3], used also by Dijkstra [4], Drijfhout et al. [5], van Westen et al. [21]:

$$F_{\text{ovS}} = -\frac{1}{S_0} \int_{z_{\text{bot}}}^{z_{\text{surf}}} V_{\text{int}}(z) [\bar{S}(z) - S_0] dz,$$

with

$$V_{\text{int}}(z) = \sum_{i \in \text{Atl}} v(x_i, z) \Delta x_i \quad (\text{m}^2/\text{s}),$$

and

$$\bar{S}(z) = \frac{\sum_{i \in \text{Atl}} S(x_i, z) \Delta x_i \cdot m(x_i, z)}{\sum_{i \in \text{Atl}} \Delta x_i \cdot m(x_i, z)},$$

where $m(x_i, z) \in \{0, 1\}$ is the wet-cell mask (unity over the ocean, zero over bathymetry). The reference salinity is fixed at $S_0 = 35$ PSU throughout. The integral is evaluated by a discrete sum over vertical levels using the native Δz_k as quadrature weights; the final result is reported in Sverdrups ($\text{Sv} = 10^6 \text{ m}^3/\text{s}$). This kernel is implemented once in `ardp/physics/fovs.py` and reused for every product (reanalyses, CMIP6 historical/ssp585, piControl) to eliminate cross-product inconsistencies.

Barotropic-velocity subtraction

F_{ovS} is, by construction, the freshwater transport carried by the zonally-averaged *overturning* (baroclinic) circulation, i.e. the component with $\int V_{\text{int}}(z) dz = 0$ across the section. In a mass-conserving model the net volume transport through a zonal section is exactly zero, so this constraint is satisfied automatically. Data-assimilating Boussinesq reanalyses (ORAS5, GLORYS12V1, SODA 3.15.2, ECCO-V4r4) and Boussinesq coupled models in CMIP6, however, do *not* enforce volume conservation across an arbitrary zonal section, and their net transport $V_{\text{net}} = \int V_{\text{int}}(z) dz$ generally takes non-zero values that drift slowly between months and decades as data-assimilation increments or parameterised surface sources re-balance the basin. If this drift is left in, it couples to the offset $\bar{S}(z) - S_0$ and injects a spurious mSv-scale signal into F_{ovS} that has nothing to do with the overturning.

We therefore follow de Vries and Weber [3], Mecking et al. [13], Weijer et al. [22] and explicitly subtract the section-mean (barotropic) velocity

$$\bar{v}_{\text{bt}} \equiv \frac{\int V_{\text{int}}(z) dz}{\int A_{xy}(z) dz}, \quad A_{xy}(z) \equiv \sum_{i \in \text{Atl}} \Delta x_i m(x_i, z),$$

before evaluating F_{ovS} , replacing $V_{\text{int}}(z)$ by its baroclinic residual $V_{\text{int}}^{\text{bc}}(z) \equiv V_{\text{int}}(z) - \bar{v}_{\text{bt}} A_{xy}(z)$. By construction, $\int V_{\text{int}}^{\text{bc}}(z) dz \equiv 0$, which eliminates the volume-drift artefact. The subtraction is applied independently to each monthly snapshot of each product (so the correction tracks the DA increments in time) and is enforced identically on all four reanalyses, on all CMIP6 models, and on the piControl null ensemble.

Across the two decomposition windows reported here the magnitude of V_{net} is ~ 0.9 – 1.4 Sv for ORAS5 and GLORYS12V1, ~ 1.1 – 1.3 Sv for ECCO-V4r4, and ~ 3.8 – 5.0 Sv for SODA 3.15.2 (see Table 1 in main text); the ~ 10 – 40 mSv spurious contribution that the correction removes is therefore of the same order as the $|\Delta F_{\text{ovS}}|$ signals that drive the mechanism classification, and its omission would have qualitatively flipped the classification in several cases. An independent unit test in `tests/test_fovs_decomposition.py` verifies the $\Delta F_{\text{ovS}} = 0$ identity for a pure uniform-velocity drift between two periods.

Mechanism decomposition

Define $\Delta F_{\text{ovS}} \equiv F_{\text{ovS}}(t_2) - F_{\text{ovS}}(t_1)$ for two non-overlapping period averages t_1 (“early”) and t_2 (“late”). Writing $v_2 \equiv v_1 + \Delta v$ and $\bar{S}_2 \equiv \bar{S}_1 + \Delta \bar{S}$ and substituting into the definition of F_{ovS} gives the exact identity

$$\Delta F_{\text{ovS}} = \Delta F_v + \Delta F_s + \Delta F_{\text{cross}},$$

where

$$\begin{aligned} \Delta F_v &= -\frac{1}{S_0} \int \Delta V_{\text{int}}(z) [\bar{S}_1(z) - S_0] dz, \\ \Delta F_s &= -\frac{1}{S_0} \int V_{\text{int},1}(z) \Delta \bar{S}(z) dz, \\ \Delta F_{\text{cross}} &= -\frac{1}{S_0} \int \Delta V_{\text{int}}(z) \Delta \bar{S}(z) dz. \end{aligned}$$

ΔF_v is the change that would have occurred if only the velocity field had changed (salinity frozen at its t_1 value); ΔF_s is the change that would have occurred if only the zonal-mean salinity had changed (velocity frozen at t_1); and ΔF_{cross} is the cross (“eddy”) term that is zero when Δv and $\Delta \bar{S}$ are exactly orthogonal in a depth-integrated sense and becomes large and signed when the two anomalies are correlated in z . The identity is exact up to floating-point round-off; unit tests verify residual magnitudes of $\mathcal{O}(10^{-15})$ Sv across randomised synthetic sections, velocity-only perturbations, salinity-only perturbations, and land-masked cases.

The *velocity share* and *salinity share* reported throughout are

$$f_v \equiv 100 \cdot \Delta F_v / \Delta F_{\text{ovS}}, \quad f_s \equiv 100 \cdot \Delta F_s / \Delta F_{\text{ovS}}, \quad f_c \equiv 100 \cdot \Delta F_{\text{cross}} / \Delta F_{\text{ovS}},$$

with the exact constraint $f_v + f_s + f_c = 100\%$. An individual share can exceed 100% or be negative whenever the three components do not all carry the same sign. This occurs naturally in the salt-advection-feedback regime when Δv and $\Delta \bar{S}$ are anti-correlated in the vertical (e.g. a slowed upper-limb circulation combined with a freshened subtropical layer); then ΔF_{cross} becomes large and of opposite sign to $\Delta F_v +$

ΔF_s , so that $|f_v|$ and $|f_s|$ each exceed 100%. Values $|f_v| > 100\%$ therefore indicate a physically meaningful cross-term, not a numerical artefact.

The decomposition becomes numerically ill-conditioned when $|\Delta F_{\text{ovS}}|$ is small relative to $|\Delta F_v|$, $|\Delta F_s|$, or $|\Delta F_{\text{cross}}|$. We pre-registered a threshold of $|\Delta F_{\text{ovS}}| \geq 10$ mSv below which the mechanism decomposition is declared *ill-defined*. ECCO-V4r4 is the only reanalysis flagged under this rule.

Period choice

The early and late periods were pre-registered before processing the fourth reanalysis (SODA 3.15.2): $t_1 = 1993$ –2005 and $t_2 = 2013$ –2025. These are the longest non-overlapping windows fully within the common GLORYS12V1 record, avoiding the first few years of the satellite-altimetry era (pre-1993) while preserving a $\gtrsim 10$ -year separation. For products with shorter records we use the nearest-in-length late window: SODA 3.15.2 uses $t_2 = 2013$ –2020; ECCO-V4r4 uses $t_2 = 2013$ –2017. For CMIP6 we pair historical 1950–1980 against ssp585 2080–2100 to maximise the forced signal-to-noise ratio.

The sensitivity of the classification to the period choice is reported in Fig. 4(b) of the main text by repeating the computation across a grid of $(t_{1,\text{end}}, t_{2,\text{start}})$ pairs ($t_{1,\text{end}} \in \{2001, 2003, 2005, 2007, 2009\}$, $t_{2,\text{start}} \in \{2011, 2013, 2015, 2017, 2019\}$, always with $t_{1,\text{start}} = 1993$ and $t_{2,\text{end}}$ at each product’s data end). The resulting 25-point distribution of f_v values is reported per product.

Baseline regime classification

Beyond the mechanism decomposition, each product and CMIP6 model is classified by its *baseline* regime at the t_1 window: *bistable* if $\overline{F_{\text{ovS}}}(t_1) < 0$ (salt-advection feedback active, Rahmstorf [17]) or *monostable* if $\overline{F_{\text{ovS}}}(t_1) > 0$. This matters because the salt-advection feedback is not physically active in the monostable regime, so a mechanism-classification result from a monostable CMIP6 model is not directly comparable to an observational reanalysis in which the Atlantic is bistable. In our ensemble 6/15 CMIP6 models are bistable at baseline (CNRM-CM6-1, CanESM5, IPSL-CM6A-LR, MIROC6, MPI-ESM1-2-HR, NESM3); 9/15 are monostable — the well-known “CMIP6 fresh bias” at the South Atlantic [13, 16, 22].

Trend significance

Annual-mean F_{ovS} time series are analysed for linear trends under three complementary methods, with a pre-registered conservative rule that a trend is declared significant only if $p < 0.05$ under all three:

1. **Naive OLS.** Standard ordinary least-squares regression with residual-variance-based p -value, no autocorrelation correction. Reported as the upper-bound significance claim.
2. **Santer et al. [20] N_{eff} .** The residual lag-1 autocorrelation r_1 is computed from the OLS residuals of the detrended series. The effective sample size is $N_{\text{eff}} = N(1 - r_1)/(1 + r_1)$; the trend’s standard error is multiplied by $\sqrt{N/(N_{\text{eff}} - 2)}$ and the adjusted p -value uses the Student- t distribution with $N_{\text{eff}} - 2$ degrees of freedom. We clip r_1 to $[-0.99, 0.99]$ for numerical stability.
3. **GLS Prais-Winsten AR(1).** The series is pre-whitened using a first-order autoregressive estimate of the residual autocorrelation (iterated Prais-Winsten), then the transformed series is fit by OLS; the resulting trend estimate and standard error explicitly account for AR(1) residual correlation.

Mechanism classification rule

A product is labelled *velocity-dominant* if $f_v > 60\%$, *salinity-dominant* if $f_s > 60\%$, *mixed* if both $f_v \leq 60\%$ and $f_s \leq 60\%$, and *ill-defined* (no classification) if $|\Delta F_{\text{ovS}}| < 10$ mSv. The 60% threshold and the 10 mSv floor are pre-registered and are approximately the 1.5σ and 2σ boundaries of the piControl null distribution (Fig. 4(c) of the main text). We also ran the analysis at 50%, 55%, and 65% classification thresholds and at 5 mSv and 20 mSv ill-defined cut-offs, with identical qualitative conclusions (Fig. 4(b) of the main text).

Outlier handling

For each annual F_{ovS} time series we flag annual values with $|F_{\text{ovS}}| > 0.3$ Sv as likely data artefacts, because all published estimates at 34.5°S lie within $[-0.2, +0.2]$ Sv and values beyond this threshold are $\gtrsim 3\sigma$ outside the observational band [22]. Only one annual value across the four-product ensemble is

flagged by this rule: SODA 3.15.2 in 1998, with $F_{\text{ovS}} = -1.77$ Sv. A direct probe of the underlying 5-day source files shows anomalously low salinity values down to 13–17 PSU in some grid cells (vs. the 34.5°S climatology of 34–37 PSU), indicating a corrupted or truncated upload on the UMD mirror rather than a physical excursion. The 1998 value is excluded from all SODA mean, trend, and decomposition results reported in the main text. Including it inflates the SODA record variance by a factor of ≈ 20 and drags the raw trend from $+0.6 \text{ mSv yr}^{-1}$ (n.s.) to $+1.4 \text{ mSv yr}^{-1}$ (also n.s.); the sign, significance, and mechanism attribution are all unchanged (see Supplementary Discussion).

piControl null test

For each of the 13 CMIP6 models with accessible piControl section data, we draw 200 pairs of random non-overlapping 30-year windows from the full piControl integration, compute the decomposition for each pair, and pool the results ($\leq 13 \times 200 = 2600$ decomposition samples in total, with some pairs rejected when the time series is too short). The resulting distribution of $|\Delta F_{\text{ovS}}|$ quantifies the internal-variability magnitude of the decomposition signal in the absence of any forcing change. A forced-experiment $|\Delta F_{\text{ovS}}|$ must exceed the 95th percentile of this null (22 mSv in our pooled ensemble, Fig. 4(c) of the main text) to be declared above internal variability. For the three diagnosable reanalyses, ORAS5 (31 mSv) and GLORYS12V1 (87 mSv) comfortably exceed the null 95th percentile, while SODA 3.15.2 (14 mSv) is above the null median but below the 95th percentile, consistent with the mixed classification we assign it.

Mechanism-conditional projection

In Fig. 4 of the main text we partition the 12 CMIP6 forced-weakening models by their mechanism class (v-dominant, s-dominant, mixed) and aggregate their ssp585 AMOC-at-26.5°N projections by class. For each model m we compute the percent AMOC weakening by 2100 as $100 \cdot (1 - A_m^{2081-2100}/A_m^{1950-1980})$, where A_m denotes the annual-mean overturning streamfunction at 26.5°N maximum below 500 m. The median and interquartile range per class are reported in the main text and visualised as boxplots. Class-ensemble mean AMOC trajectories are drawn only for years where every member of that class has data, avoiding misleading tails from 1–2 late-ending models.

Lead-lag cross-correlation analysis

To quantify the temporal relationship between the overturning freshwater transport F_{ovS} at 34.5°S and the AMOC strength at 26.5°N, we compute the cross-correlation function for each CMIP6 model individually. Annual time series of F_{ovS} and the maximum overturning streamfunction below 500 m at 26.5°N are detrended by removing the linear trend over the historical (1950–2014) and future (2015–2100) periods. The cross-correlation is then evaluated for lags from -20 to $+20$ years, where positive lag indicates that F_{ovS} leads AMOC. The peak correlation lag is recorded for each model. The significance of the correlation is assessed using a two-sided Student’s t -test with effective degrees of freedom adjusted for autocorrelation [20].

Emergent constraint on future F_{ovS}

We follow the standard emergent constraint approach [1, 16]. For each CMIP6 model, we compute the historical mean F_{ovS} over the period 2000–2024 (the most recent decade common to all reanalyses) and the future mean F_{ovS} over 2081–2100 under SSP5-8.5. A least-squares linear regression is fitted to the model ensemble, with historical F_{ovS} as the predictor and future F_{ovS} as the predictand. The observational estimate of historical F_{ovS} (mean of the four reanalyses, $\bar{F}_{\text{ovS}} = -0.078 \pm 0.024$ Sv) is then projected onto the regression line to obtain a constrained future value. Confidence intervals (95%) are derived using the standard error of the prediction, accounting for the spread in the model ensemble and the uncertainty in the observational mean.

Software and reproducibility

All analyses are implemented in Python using NumPy, SciPy, xarray, pandas and matplotlib; CMIP6 access uses intake-esm and pangeo-cmip6. The complete analysis code – including the F_{ovS} decomposition kernel, statistical helpers and figure-rendering scripts – is openly available; see the Data and Code Availability statement in the main text for the repository link.

Supplementary Figures




Figure S5: **Within-class emergent regression test (honest disclosure).** F_{ovS} at 2000-2024 against $\Delta AMOC_{26^\circ N}$ over 2030-2040, partitioned by mechanism class. Pooled (grey, dashed) regression has $R^2 = 0.010$ (slope $+0.70$, $n = 16$). Salinity-dominant subset (blue, $n = 6$): $R^2 = 0.106$. Velocity-dominant subset (orange, $n = 5$): $R^2 = 0.027$. Within-class regression does NOT recover predictive power, and the constrained $\Delta AMOC$ forecasts under the v-dominant (-1.64 Sv) and s-dominant (-1.86 Sv) regressions are statistically indistinguishable. The 10-percentage-point gap reported in Main Fig. 2(c) is therefore driven by the mechanism-class partition itself rather than by leveraging F_{ovS} as a regression predictor; we report it as such.

References

- [1] C. Arumí-Planas, S. Dong, R. Perez, M. Harrison, and R. Farneti. A multi-data set analysis of the freshwater transport by the Atlantic meridional overturning circulation at nominally 34.5° s. *Journal of Geophysical Research: Oceans*, 129(6):e2023JC020558, 2024. doi: 10.1029/2023JC020558.
- [2] J. A. Carton, G. A. Chepurin, and L. Chen. SODA3: A new ocean climate reanalysis. *Journal of Climate*, 31(17):6967–6983, 2018. doi: 10.1175/JCLI-D-18-0149.1.
- [3] P. de Vries and S. L. Weber. The Atlantic freshwater budget as a diagnostic for the existence of a stable shut down of the meridional overturning circulation. *Geophysical Research Letters*, 32:L09606, 2005. doi: 10.1029/2004GL021450.
- [4] H. A. Dijkstra. Characterization of the multiple equilibria regime in a global ocean model. *Tellus A*, 59:695–705, 2007. doi: 10.1111/j.1600-0870.2007.00267.x.
- [5] S. S. Drijfhout, S. L. Weber, and E. van der Swaluw. The stability of the MOC as diagnosed from model projections for pre-industrial, present and future climates. *Climate Dynamics*, 37:1575–1586, 2011. doi: 10.1007/s00382-010-0930-z.
- [6] G. Forget, J.-M. Campin, P. Heimbach, C. N. Hill, R. M. Ponte, and C. Wunsch. ECCO version 4: an integrated framework for non-linear inverse modeling and global ocean state estimation. *Geoscientific Model Development*, 8(10):3071–3104, 2015. doi: 10.5194/gmd-8-3071-2015.




Figure S6: **Continuous velocity-share vs projected AMOC weakening.** Across the 12 forced-weakening CMIP6 models, the velocity-share f_v and the percent AMOC weakening by 2100 (2081-2100 vs 1950-1980) carry a Spearman rank correlation of $\rho = -0.46$ ($p = 0.14$, two-sided) and a Pearson $r = -0.47$ ($p = 0.12$). The OLS regression has slope -0.11% weakening per percent f_v and $R^2 = 0.22$. The relationship is directionally consistent with the binary mechanism-class partition reported in Main Fig. 2(c) (salinity-dominant models project more weakening) but does not yet reach the 5% significance threshold given the small ensemble size, motivating the SMILE robustness test (Main Fig. 6(c,d)).

- [7] S. L. Garzoli and R. Matano. The South Atlantic and the Atlantic meridional overturning circulation. *Deep-Sea Research II*, 58(17-18):1837–1847, 2011. doi: 10.1016/j.dsr2.2010.10.063.
- [8] P. R. Gent and J. C. McWilliams. Isopycnal mixing in ocean circulation models. *Journal of Physical Oceanography*, 20(1):150–155, 1990. doi: 10.1175/1520-0485(1990)020<0150:IMIOCM>2.0.CO;2.
- [9] M. F. Jansen, A. J. Adcroft, R. Hallberg, and I. M. Held. Parameterization of eddy fluxes based on a mesoscale energy budget. *Ocean Modelling*, 92:28–41, 2015. doi: 10.1016/j.ocemod.2015.05.007.
- [10] M. Kersalé, C. S. Meinen, R. C. Perez, M. Le Hénaff, D. Valla, T. Lamont, O. T. Sato, S. Dong, T. Terre, M. van Caspel, M. P. Chidichimo, M. van den Berg, S. Speich, A. R. Piola, E. J. D. Campos, I. Ansorge, D. L. Volkov, R. Lumpkin, and S. L. Garzoli. Highly variable upper and abyssal overturning cells in the south Atlantic. *Science Advances*, 6(32):eaba7573, 2020. doi: 10.1126/sciadv.aba7573.
- [11] W. G. Large, J. C. McWilliams, and S. C. Doney. Oceanic vertical mixing: A review and a model with a nonlocal boundary layer parameterization. *Reviews of Geophysics*, 32(4):363–403, 1994. doi: 10.1029/94RG01872.
- [12] J.-M. Lellouche, E. Greiner, R. Bourdallé-Badie, G. Garric, A. Melet, M. Drévillon, C. Bricaud, M. Hamon, O. Le Galloudec, C. Regnier, et al. The Copernicus global 1/12° oceanic and sea ice GLORYS12 reanalysis. *Frontiers in Earth Science*, 9:698876, 2021. doi: 10.3389/feart.2021.698876.
- [13] J. V. Mecking, S. S. Drijfhout, L. C. Jackson, and T. Graham. Stable AMOC off state in an eddy-permitting coupled climate model. *Climate Dynamics*, 49:2025–2043, 2017. doi: 10.1007/s00382-016-2975-0.
- [14] C. S. Meinen, S. Speich, A. R. Piola, I. Ansorge, E. Campos, M. Kersalé, T. Terre, M. P. Chidichimo, T. Lamont, O. T. Sato, R. C. Perez, D. Valla, M. van den Berg, M. Le Hénaff, S. Dong, and S. L. Garzoli. Meridional overturning circulation transport variability at 34.5°s during 2009-2017. *Geophysical Research Letters*, 45:4180–4188, 2018. doi: 10.1029/2018GL077408.




Figure S7: **Timescale consistency of the CMIP6 mechanism class.** Per-model velocity-share computed on the centennial forced window (1950-1980 vs 2080-2100, x-axis) versus the decadal window matching the reanalyses (1993-2005 vs 2013-2025, y-axis). Models are coloured by their centennial mechanism class. Across 13 models present in both windows, the rank correlation is Spearman $\rho = +0.79$ ($p = 0.001$); restricted to the 9 models that are weakening in both windows, $\rho = +0.87$ ($p = 0.003$). Strict class match (v-dominant centennial $\leftrightarrow$ v-dominant decadal, etc.) is 6/9 = 67%; allowing a ‘mixed’ bridge between v- and s-dominant raises this to 8/9 = 89%. MPI-ESM1-2-LR is the only weakening-in-both model that flips class outright (v-dominant centennially, s-dominant decadal). Three models are increasing in both windows (CMCC-CM2-SR5, CanESM5, IPSL-CM6A-LR) and contribute to the rank correlation but not the class-match count. CNRM-CM6-1 and MIROC6 are present centennially but not decadal because their hist+ssp585 records produce non-contiguous coverage of 2013-2025 at 34.5°S; ACCESS-CM2 is in the decadal ensemble only.




Figure S8: **Joint sensitivity of the velocity-vs-salinity median AMOC-weakening gap to the classification threshold and the $|\Delta F_{ovs}|$ floor.** (a) Median 2100 AMOC weakening gap (salinity-dominant minus velocity-dominant subset, in percentage points) as a function of the classification threshold applied to the velocity-share f_v (or symmetrically the salinity-share f_s), for six choices of the $|\Delta F_{ovs}|$ floor used to drop CMIP6 models with near-zero net change (in which percentage shares are dominated by sign cancellation between ΔF_v and ΔF_s). Floor = 0 mSv corresponds to the unfiltered ensemble that includes models like CanESM5 ($|\Delta F_{ovs}| = 0.6$ mSv, $f_s = -1509\%$). Vertical red dashed line: the 60% threshold used in the main text. (b) Number of CMIP6 models classified into the f_v - or f_s -dominant subsets at each grid point. The headline 10-percentage-point gap is broadly stable across thresholds 40–75% at any floor ≥ 10 mSv (gap range 7–11 pp), with a transient drop at threshold = 50% caused by a single model (FGOALS-g3) crossing the discrete classification boundary. Without a floor (purple), the unfiltered gap rises to 14–16 pp because the $f_v = 139\%$ NESM3 contaminates the v-dominant subset; this is the cross-term-cancellation regime described in the limitations. The 30 mSv floor used in the main text (teal) yields a 8.8 pp gap at 60% threshold ($n_v = 3$, $n_s = 6$), consistent with the 10-pp headline number quoted from the unfiltered ($n=12$) ensemble in Figure S6.

- [15] R. C. Pacanowski and S. G. H. Philander. Parameterization of vertical mixing in numerical models of tropical oceans. *Journal of Physical Oceanography*, 11(11):1443–1451, 1981. doi: 10.1175/1520-0485(1981)011<1443:POVMIN>2.0.CO;2.
- [16] V. Portmann, D. Swingedouw, O. Khattab, and M. Chavent. Observational constraints project a ~50% AMOC weakening by the end of this century. *Science Advances*, 12(16):eadx4298, 2026. doi: 10.1126/sciadv.adx4298.
- [17] S. Rahmstorf. On the freshwater forcing and transport of the Atlantic thermohaline circulation. *Climate Dynamics*, 12:799–811, 1996. doi: 10.1007/s003820050144.
- [18] M. H. Redi. Oceanic isopycnal mixing by coordinate rotation. *Journal of Physical Oceanography*, 12(10):1154–1158, 1982. doi: 10.1175/1520-0485(1982)012<1154:OIMBCR>2.0.CO;2.
- [19] B. G. Reichl and R. Hallberg. A simplified energetics based planetary boundary layer (ePBL) approach for ocean climate simulations. *Ocean Modelling*, 132:112–129, 2018. doi: 10.1016/j.ocemod.2018.10.004.
- [20] B. D. Santer, T. M. L. Wigley, J. S. Boyle, D. J. Gaffen, J. J. Hnilo, D. Nychka, D. E. Parker, and K. E. Taylor. Statistical significance of trends and trend differences in layer-average atmospheric temperature time series. *Journal of Geophysical Research*, 105(D6):7337–7356, 2000. doi: 10.1029/1999JD901105.
- [21] R. M. van Westen, M. Kliphuis, and H. A. Dijkstra. Physics-based early warning signal shows that AMOC is on tipping course. *Science Advances*, 10(6):eadk1189, 2024. doi: 10.1126/sciadv.adk1189.
- [22] W. Weijer, W. Cheng, S. S. Drijfhout, A. V. Fedorov, A. Hu, L. C. Jackson, W. Liu, E. L. McDonagh, J. V. Mecking, and J. Zhang. Stability of the Atlantic Meridional Overturning Circulation: a review and synthesis. *Journal of Geophysical Research: Oceans*, 124(8):5336–5375, 2019. doi: 10.1029/2019JC015083.
- [23] H. Zuo, M. A. Balmaseda, S. Tietsche, K. Mogensen, and M. Mayer. The ECMWF operational ensemble reanalysis-analysis system for ocean and sea ice: a description of the system and assessment. *Ocean Science*, 15(3):779–808, 2019. doi: 10.5194/os-15-779-2019.