Source SHA-256: `c6c43f07ac3d550a72360a5cf1aee434eb45c72cf70ebe36aaccbe14d4e8864f`. Repository links identify source paths; newly committed pages may be unavailable online until publication.

Separating nearby vegetation from regional background pollen

Paper and question. Sugita, S. (2007), "Theory of quantitative reconstruction of vegetation II: all you need is LOVE," *The Holocene* 17(2), 243–257. This second Landscape Reconstruction Algorithm (LRA) theory paper, commonly cited as Sugita (2007b), asks how pollen from a smaller target basin can estimate local vegetation when part of its pollen arrived from farther away. We examine it because a local result requires a separate regional background estimate and a defensible spatial extent; a REVEALS regional fraction cannot simply be renamed local cover. Indexed publisher citation and abstract; our LOVE interface account.

Scoped conclusion. In the publisher abstract, Sugita's POLLSCAPE simulations support background-pollen prediction and report better local-cover estimates from LOVE than from pollen percentages. They also support a backward-modeling approach to the relevant source area of pollen, which remains hard to estimate for past landscapes. These are simulation claims. Our later numerical interface check used a constructed ring and no independently observed local vegetation; it does not reproduce the 2007 experiments. Publisher abstract; interface result.

What LOVE estimates and needs

A small lake or peatland receives pollen from plants near the basin plus a regional background. Pollen counts are grain shares, while vegetation cover is area share; taxon pollen productivity and dispersal separate the two. REVEALS ("Regional Estimates of VEgetation Abundance from Large Sites") estimates regional cover from large-site pollen. LOVE ("LOcal Vegetation Estimates") subtracts the background contribution and estimates cover within a relevant local source area. PPEs are relative pollen productivity estimates; fall speed, basin type and radius, and chosen extent control the modeled deposition. Later method profile.



The abstract specifies regional input cover over 10^4–10^5 km^2 from large sites of at least 10^2 ha, plus target-site counts, PPEs and a relevant local area. It provides no simulated-basin count, exact radii, numeric errors or full backward-modeling algorithm. Publisher abstract; companion REVEALS account.

A constructed interface trace, separate from the 2007 experiment

Our later LandClim author-source interface supplies 1,000 regional REVEALS draws from five pollen-supported sites, on 31 taxa, for the first time bin of grid 512. It feeds those draws, the source-aligned local `BROADOWN` pollen and declared transport inputs to one constructed 200–300 m analytical LOVE ring. `BROADOWN` also belongs to the regional cohort, so this is neither a held-out site nor an independent local-vegetation comparison. A placeholder regional-pollen argument is taxon-aligned for the interface but does not reconstruct the supplied regional-cover draws. Exact machine receipt; source and replay route.

The resulting tensor has $1{,}000 \times 1 \times 31 = 31{,}000$ finite local fraction cells. 8,312 are negative; its minimum is -0.4695300882. For a concrete taxon row, *Calluna vulgaris* has mean local output 0.6135168974, standard deviation across these draws 0.0385115602, and zero negative draws. The standard deviation is variation over this constructed input draw set, not an ecological confidence interval. Because other taxa

yield negative cover, the complete output is not a physically admissible vegetation composition. The calculation demonstrates that the regional tensor enters the local interface and its declared shape is enforced; it does not show that any inferred ring resembles real vegetation. Tensor result; Calluna row.

The separate packaged LOVE example reinforces the spatial limit: among its first ten candidate rings, only three satisfy the source's local-basin boundary, and 16 of 30 valid taxon/ring cells are negative while six exceed one. Those are fixture outcomes for a later implementation, not a test or reproduction of Sugita's 2007 POLLSCAPE finding. Nor does a finite result determine the relevant source area for an actual past basin. Exact ring rows.

Evidence boundary and reproduction

The indexed publisher abstract supplies the original theory and qualitative simulation findings. A lawful full 2007 article or original simulation scenario table was not located in the retained packet examined here, and direct publisher page retrieval did not yield full text. Thus the paper's exact simulation denominators, equations and result rows are not evaluated. The constructed interface uses a later author REVEALS release and current analytical LOVE implementation; the 2024 LOVE/LOVEoptim paper account concerns a further, separate simulated population and R tools. None supplies measured Nordic local vegetation. Publisher abstract; interface account.

The retained interface report and taxon rows are readable without private author inputs. With the intact lawful author snapshot, a fresh run writes only to an absent repository `artifacts/` directory:

sh

```
PYTHONDONTWRITEBYTECODE=1 OMP_NUM_THREADS=1 OPENBLAS_NUM_THREADS=1 \
  MKL_NUM_THREADS=1 NUMEXPR_NUM_THREADS=1 VECLIB_MAXIMUM_THREADS=1 \
  .venv/bin/python reports/science/reveals-love-interface/replay.py \
  --repository . --source-root "$BIJUX_LANDCLIM_AUTHOR_REFERENCE" \
  --output artifacts/jobs/reveals-love-interface/reproduction
```

The producer authenticates the consumed source files and rejects a 999-draw input before accepting the 1,000-draw tensor. This route replays our interface only, not the Sugita simulations. Replay method.

The unsent source request is for a lawful full Sugita (2007b) paper and original POLLSCAPE scenario definitions, target and regional basin identities/geometries, pollen counts, regional truth/REVEALS vectors, PPE and fall-speed tables, background-pollen terms, relevant-source-area search rule, local vegetation truth and uncertainty/error rows, plus usable source permissions. The unsent qualified-review request is to decide for each proposed historical target whether its regional cohort is independent and temporally comparable, its basin and local-area geometry are defensible, its taxon/parameter support is adequate, and a measured local vegetation reference exists before any accuracy claim. No independent review, original figure reproduction or colleague reading outcome is claimed.