Bijux Pollenomics: scientific evidence

Sources, method correspondence, predictions and model use

Bijux Pollenomics

29 September 2026

Source-bound research briefing

Evidence ladder

1. Archive/source custody and taxon-count mapping.
2. Author-equation and numerical implementation parity.
3. Observed-pollen held-out prediction.
4. LRA reconstruction and forward-pollen prediction.
5. Independent measured-vegetation validation: no admitted pair yet.

Each rung answers a different question; a result at one rung does not imply the next.

Evidence: `reports/README.md`; `reports/needed/README.md`

Four-country source and prediction accounting

Target	Sites	Samples	Assigned/native NISP	Scored/attempted bins
Denmark	1	91	122,998/129,768	24/24
Finland	28	1,797	1,229,019/1,260,281	435/435
Norway	52	3,544	2,018,295/2,286,020	1,084/1,178
Sweden	3	228	221,800/240,966	51/54

Closed 31-target author pollen composition. Excluded/unmapped mass remains outside the score. Zero Nordic samples are admitted to LRA by this study.

Evidence: `docs/report/regions/nordic/author-target-pollen/README.md`; `docs/report/pollen-reconstruction/README.md`

What reproduces numerically

- Author raw-to-prepared replay: 17 example files, 871 raw sample rows, 15,028 matching prepared cells.
- Python/independent author-R: 14,074 supported mean and SE values within declared tolerances.
- Archived 100–350 BP means can differ by over 10 percentage points; window and source identity are not the same.
- Archived ISTS and TBS mean arithmetic closes over all 10,500 grid-window rows, but biological classification and SE transformation remain open.

Evidence: `docs/report/pollen-reconstruction/paper-correspondence.md`

English LOVE: declared model profile

- 15 source sites, 1250–1750 BP, 30 targets; published PPE and fall speeds held fixed for retained taxa.
- Regional REVEALS plus ring-taxon-preserving LOVE; closed, nonnegative local cover.
- Uniform-start SLSQP: 300 iterations, 16,000 evaluations, objective tolerance 10^{-8}.
- 192/195 training fits complete; three failures also refused under a bounded 600-iteration check.

These are source-specific development choices, not universally optimal LRA parameters or the original R optimizer.

Evidence: `docs/report/pollen-reconstruction/author-lra-constrained/README.md`

Matched pollen comparison: result and limitation

	MAE		Hellinger	
	Direct	LRA	Direct	LRA
Training mean	.020496	.026273	.282082	.360671
Nearest	.019552	.021132	.255767	.286001
Inverse distance squared	.018579	.021088	.243673	.285415

14 paired English sites, 30 taxa; all three direct means lower. Previously exposed source examples and unequal training information limit interpretation.

Evidence: `docs/report/pollen-reconstruction/author-lra-constrained/direct-pollen/paired-hellinger.json`

Model selection follows the target and support

- Fixed spatial baselines permit clear paired interpretation on observed pollen.
- A bounded RBF candidate improved one prior-exposed author comparison on common support but reduced coverage.
- Country-excluded tuning did not establish superiority over inverse-distance-squared in the Nordic development holdouts.
- For LRA, input identity, chronology, basin geometry, PPE and denominator must be qualified before any parameter search is meaningful.

Evidence: `reports/science/pollen-prediction.md`; `reports/needed/README.md`

HYDE and vegetation: retain the boundary

- Four-country HYDE source cropland grid sums match 84 country/year/scenario totals within 8.51×10^{-11} km^2.
- This checks modeled-source internal accounting, not pollen-model benefit.
- No independent vegetation measurement pair has been admitted for prediction accuracy.

Next experiments require source semantics and a matched ecological reference, with outcomes protected from model selection.

Evidence: `docs/report/pollen-reconstruction/paper-correspondence.md`; `reports/needed/README.md`

Open source decisions

- Abborrtjärnen: LandClim-selected age model, historical basin radius, eligible denominator, PPE use and rights.
- Other Nordic cores: physical-site and sample/window lineage before LRA admission.
- Historical publication: exact source revisions and authenticated PFT/uncertainty transformation.
- Independent confirmation: new qualified cohort frozen before target inspection.

Evidence: `reports/needed/README.md`