

Supplement: Disaster Response on September 11, 2001 Through the Lens of Statistical Network Analysis

Michael Schweinberger[†]

Department of Statistics, Pennsylvania State University, University Park, PA, USA

Miruna Petrescu-Prahova

Department of Health Services, University of Washington, Seattle, WA, USA

Duy Quang Vu

Department of Mathematics and Statistics, University of Melbourne, Melbourne, Australia

We are interested in maximising the lower bound on $\log P(\mathbf{Y} = \mathbf{y})$ given by (23). We note that, despite of the fact that the family of auxiliary distributions is fully factorised, the lower bound on $\log P(\mathbf{Y} = \mathbf{y})$ is intractable, because

$$\int \left[\log \frac{P(\mathbf{Y} = \mathbf{y} \mid \boldsymbol{\delta}) p(\boldsymbol{\delta})}{q(\boldsymbol{\delta} \mid \boldsymbol{\vartheta})} \right] q(\boldsymbol{\delta} \mid \boldsymbol{\vartheta}) d\boldsymbol{\delta} = \sum_{i < j}^n E_{\boldsymbol{\vartheta}} [\log P(Y_{ij} = y_{ij} \mid \boldsymbol{\delta})] \quad (27)$$

$$+ E_{\boldsymbol{\vartheta}} [\log p(\boldsymbol{\delta})] - E_{\boldsymbol{\vartheta}} [\log q(\boldsymbol{\delta} \mid \boldsymbol{\vartheta})],$$

where

$$E_{\boldsymbol{\vartheta}} [\log P(Y_{ij} = y_{ij} \mid \boldsymbol{\delta})] = \left[\sum_{k=1}^K \vartheta_{Z_{ik}} \vartheta_{\theta_k, 1} + \sum_{l=1}^K \vartheta_{Z_{jl}} \vartheta_{\theta_l, 1} \right] y_{ij} \quad (28)$$

$$+ \sum_{k=1}^K \sum_{l=1}^K \vartheta_{Z_{ik}} \vartheta_{Z_{jl}} E_{\boldsymbol{\vartheta}} \{ -\log [1 + \exp(\theta_k + \theta_l)] \}$$

includes the intractable expectations $E_{\boldsymbol{\vartheta}} \{ -\log [1 + \exp(\theta_k + \theta_l)] \}$. To obtain a tractable lower bound on $\log P(\mathbf{Y} = \mathbf{y})$, we lower bound $E_{\boldsymbol{\vartheta}} \{ -\log [1 + \exp(\theta_k + \theta_l)] \}$ by Jensen's inequality:

$$E_{\boldsymbol{\vartheta}} \{ -\log [1 + \exp(\theta_k + \theta_l)] \} \geq -\log E_{\boldsymbol{\vartheta}} [1 + \exp(\theta_k + \theta_l)] \quad (29)$$

$$= -\log \left[1 + \exp(\vartheta_{\theta_k, 1} + \tfrac{1}{2} \vartheta_{\theta_k, 2} + \vartheta_{\theta_l, 1} + \tfrac{1}{2} \vartheta_{\theta_l, 2}) \right].$$

The lower bound on $\log P(\mathbf{Y} = \mathbf{y})$ given by (27), (28), and (29) can be maximised with respect to $\boldsymbol{\vartheta}$ by coordinate ascent algorithms.

The lower bound on $\log P(\mathbf{Y} = \mathbf{y})$ can be maximised with respect to $\vartheta_{\alpha, 1}$, $\vartheta_{\alpha, 2}$, $\vartheta_{\mu, 1}$, $\vartheta_{\mu, 2}$, $\vartheta_{\sigma^{-2}, 1}$, $\vartheta_{\sigma^{-2}, 2}$, $\vartheta_{V_k, 1}$, $\vartheta_{V_k, 2}$, $\vartheta_{Z_{ik}}$ as follows:

$$\vartheta_{\alpha, 1} = A_1 + K - 1, \quad (30)$$

$$\vartheta_{\alpha, 2} = B_1 - \sum_{k=1}^{K-1} [\Psi(\vartheta_{V_k, 2}) - \Psi(\vartheta_{V_k, 1} + \vartheta_{V_k, 2})], \quad (31)$$

[†]Email: michael.schweinberger@stat.psu.edu

$$\vartheta_{\mu,1} = \frac{MS^{-2} + E_{\boldsymbol{\vartheta}}[\sigma^{-2}] \sum_{k=1}^K E_{\boldsymbol{\vartheta}}[\theta_k]}{S^{-2} + K E_{\boldsymbol{\vartheta}}[\sigma^{-2}]}, \quad (32)$$

$$\vartheta_{\mu,2} = \frac{1}{S^{-2} + K E_{\boldsymbol{\vartheta}}[\sigma^{-2}]}, \quad (33)$$

$$\vartheta_{\sigma^{-2},1} = A_2 + \frac{K}{2}, \quad (34)$$

$$\vartheta_{\sigma^{-2},2} = B_2 + \sum_{k=1}^K \frac{\vartheta_{\theta_k,1}^2 - 2\vartheta_{\theta_k,1}\vartheta_{\mu,1} + \vartheta_{\theta_k,2}}{2} + \frac{K(\vartheta_{\mu,1}^2 + \vartheta_{\mu,2})}{2}, \quad (35)$$

$$\vartheta_{V_k,1} = 1 + \sum_{i=1}^n \vartheta_{Z_{ik}}, \quad k = 1, \dots, K-1, \quad (36)$$

$$\vartheta_{V_k,2} = E_{\boldsymbol{\vartheta}}[\alpha] + \sum_{i=1}^n \sum_{l=k+1}^K \vartheta_{Z_{il}}, \quad k = 1, \dots, K-1, \quad (37)$$

$$\begin{aligned} \log \vartheta_{Z_{ik}} &\propto \Psi(\vartheta_{V_k,1}) - \Psi(\vartheta_{V_k,1} + \vartheta_{V_k,2}) + \sum_{l=1}^{k-1} [\Psi(\vartheta_{V_l,2}) - \Psi(\vartheta_{V_l,1} + \vartheta_{V_l,2})] \\ &+ \sum_{j \neq i}^n \vartheta_{\theta_k,1} y_{ij} - \sum_{j \neq i}^n \sum_{l=1}^K \vartheta_{Z_{jl}} \log \left[1 + \exp(\vartheta_{\theta_k,1} + \frac{1}{2}\vartheta_{\theta_k,2} + \vartheta_{\theta_l,1} + \frac{1}{2}\vartheta_{\theta_l,2}) \right], \end{aligned} \quad (38)$$

where $\Psi(\cdot)$ denotes the digamma function and the expectations $E_{\boldsymbol{\vartheta}}[\alpha]$, $E_{\boldsymbol{\vartheta}}[\theta_k]$, and $E_{\boldsymbol{\vartheta}}[\sigma^{-2}]$ under the auxiliary distribution indexed by $\boldsymbol{\vartheta}$ are tractable.

The lower bound on $\log P(\mathbf{Y} = \mathbf{y})$ with respect to $\vartheta_{\theta_k,1}$ and $\vartheta_{\theta_k,2}$ can be maximised by maximising

$$\begin{aligned} &- \sum_{k=1}^K \frac{\vartheta_{\sigma^{-2},1}}{2\vartheta_{\sigma^{-2},2}} [\vartheta_{\theta_k,1}^2 - 2\vartheta_{\theta_k,1}\vartheta_{\mu,1} + \vartheta_{\theta_k,2}] + \frac{1}{2} \sum_{k=1}^K \log \vartheta_{\theta_k,2} \\ &+ \sum_{i < j}^n \left[\sum_{k=1}^K \vartheta_{Z_{ik}} \vartheta_{\theta_k,1} + \sum_{l=1}^K \vartheta_{Z_{jl}} \vartheta_{\theta_l,1} \right] y_{ij} \\ &- \sum_{i < j}^n \sum_{k=1}^K \sum_{l=1}^K \vartheta_{Z_{ik}} \vartheta_{Z_{jl}} \log \left[1 + \exp(\vartheta_{\theta_k,1} + \frac{1}{2}\vartheta_{\theta_k,2} + \vartheta_{\theta_l,1} + \frac{1}{2}\vartheta_{\theta_l,2}) \right]. \end{aligned} \quad (39)$$

Since $\vartheta_{\theta_k,2}$ are the precision parameters of the marginal auxiliary distributions of θ_k , the constraints $\vartheta_{\theta_k,2} > 0$ must be respected ($k = 1, \dots, K$). Two possible approaches are constrained maximisation subject to the constraints $\vartheta_{\theta_k,2} > 0$ and unconstrained maximisation by reparameterising $\rho_{\theta_k,2} : \vartheta_{\theta_k,2} \mapsto \log \vartheta_{\theta_k,2}$ and maximising with respect to $\rho_{\theta_k,2}$ ($k = 1, \dots, K$).