## Risk of first-time diagnosis of Alzheimer's disease in patients with type 2 diabetes (without obesity) (comparison between matched semaglutide vs other anti-diabetes medications groups)

			J			<b>0</b> 1 /
Size/Group	Exposure Group	Comparison Group	Exposure Group Cases (Overall Risk)	Comparison Group Cases(Overall Risk		HR (95% CI)
7,544	Semaglutide	Insulin	16 (0.21%)	44 (0.58%)	<b>⊢</b>	0.32 (0.18–0.57)
7,548	Semaglutide	Metformin	16 (0.21%)	38 (0.50%)	<b>⊢</b>	0.41 (0.23–0.73)
7,507	Semaglutide	DPP-4i	16 (0.21%)	32 (0.43%)	<del>  </del>	0.47 (0.26–0.86)
7,539	Semaglutide	SGLT2i	16 (0.21%)	29 (0.39%)	-	0.54 (0.29–0.99)
7,532	Semaglutide	SU	16 (0.21%)	43 (0.57%)	<b>⊢</b>	0.35 (0.20-0.62)
6,422	Semaglutide	TZD	15 (0.23%)	43 (0.67%)	<del></del>	0.33 (0.18–0.59)
7,539	Semaglutide	Other GLP-1RAs	16 (0.21%)	29 (0.39%)	-	0.54 (0.29–0.99)
					0.10 0.20 0.40 0.70 <b>Hazard Ratio (H</b>	2.0 3.00 5.00 8.00 IR)