Risk of first-time diagnosis of Alzheimer's disease in patients with type 2 diabetes (with obesity) (comparison between matched semaglutide vs other anti-diabetes medications groups)

Size/Group	Exposure Group	Comparison Group	Exposure Group Cases (Overall Risk)	Comparison Group Cases(Overall Risk)		HR (95% CI)
9,533	Semaglutide	Insulin	11 (0.12%)	33 (0.35%)	├──	0.29 (0.15–0.58)
9,530	Semaglutide	Metformin	11 (0.12%)	23 (0.24%)	⊢	0.45 (0.22–0.92)
8,224	Semaglutide	DPP-4i	11 (0.13%)	20 (0.24%)	├──■	0.50 (0.24–1.05)
7,649	Semaglutide	SGLT2i	<10 (<0.13%)	19 (0.25%)	├──	0.48 (0.22–1.03)
8,867	Semaglutide	SU	11 (0.12%)	33 (0.37%)	⊢	0.31 (0.15–0.61)
4,256	Semaglutide	TZD	<10 (<0.24%)	20 (0.47%)	⊢	0.31 (0.13–0.74)
9,448	Semaglutide	Other GLP-1RAs	11 (0.12%)	18 (0.19%)	 ■ 	0.59 (0.28–1.25)
					0.10 0.20 0.40 0.70 2.0 3.00 5.00 8.0 Hazard Ratio (HR)	¬ O