Risk of first-time diagnosis of Alzheimer's disease in patients with type 2 diabetes (age >= 60) (comparison between matched semaglutide vs other anti-diabetes medications groups)

Size/Group	Exposure Group	Comparison Group	Exposure Group Cases (Overall Risk)	Comparison Group Cases(Overall Risk		HR (95% CI)
8,225	Semaglutide	Insulin	27 (0.33%)	59 (0.72%)	⊢ ■	0.40 (0.26–0.64)
8,232	Semaglutide	Metformin	27 (0.33%)	61 (0.74%)	├──	0.44 (0.28-0.69)
8,003	Semaglutide	DPP-4i	27 (0.34%)	44 (0.55%)	├──	0.58 (0.36–0.94)
7,780	Semaglutide	SGLT2i	26 (0.33%)	30 (0.39%)		0.83 (0.49–1.40)
8,115	Semaglutide	SU	27 (0.33%)	58 (0.72%)	├──	0.45 (0.28–0.70)
6,422	Semaglutide	TZD	23 (0.36%)	43 (0.67%)	⊢	0.50 (0.30-0.83)
8,197	Semaglutide	Other GLP-1RAs	28 (0.32%)	41 (0.50%)	 ■	0.62 (0.38–1.02)
					0.10 0.20 0.40 0.70 2.0 3.00 5.00 8.00 Hazard Ratio (HR)	