Risk of overall outpatient medical encounters in patients with type 2 diabetes (comparison between matched semaglutide vs other anti-diabetes medications groups)

Size/Group	Exposure Group	Comparison Group	Exposure Group Cases (Overall Risk)	Comparison Group Cases(Overall Risk	•	95% CI)
17,087	Semaglutide	Insulin	15,296 (89.52%)	14,707 (86.07%)	0.86 (0.84	4–0.87)
17,080	Semaglutide	Metformin	15,286 (88.40%)	14,915 (87.32%)	1.03 (1.0	1–1.06)
15,878	Semaglutide	DPP-4i	14,143 (89.07%)	13,811 (86.98%)	1.02 (1.00)–1.05)
15,288	Semaglutide	SGLT2i	13,573 (88.78%)	13,246 (86.64%)	1.04 (1.02	2–1.07)
16,503	Semaglutide	SU	14,737 (89.30%)	14,224 (86.19%)	1.05 (1.02	2–1.07)
10,847	Semaglutide	TZD	9,452 (87.14%)	9,093 (83.83%)	1.08 (1.05	5–1.11)
17,029	Semaglutide	Other GLP-1RAs	15,237 (89.48%)	15,098 (88.66%)	0.99 (0.97	7–1.01)
					0.10 0.20 0.40 0.70 2.0 3.00 5.00 8.00 Hazard Ratio (HR)	