Risk of first-time diagnosis of Alzheimer's disease in men with type 2 diabetes (comparison between matched semaglutide vs other anti-diabetes medications groups)

Size/Group	Exposure Group	Comparison Group	Exposure Group Cases (Overall Risk)	Comparison Group Cases(Overall Risk		HR (95% CI)
6,895	Semaglutide	Insulin	13 (0.19%)	24 (0.35%)	├──	0.48 (0.24–0.93)
6,891	Semaglutide	Metformin	13 (0.19%)	28 (0.41%)	⊢	0.45 (0.23-0.86)
6,629	Semaglutide	DPP-4i	13 (0.20%)	31 (0.47%)		0.39 (0.20-0.74)
6,881	Semaglutide	SGLT2i	13 (0.19%)	11 (0.16%)	⊢	1.14 (0.51–2.55)
6,850	Semaglutide	SU	13 (0.19%)	30 (0.44%)	⊢	0.40 (0.21–0.77)
5,350	Semaglutide	TZD	13 (0.24%)	17 (0.32%)	 	0.71 (0.34–1.46)
6,881	Semaglutide	Other GLP-1RAs	13 (0.19%)	11 (0.16%)	 	1.14 (0.51–2.55)
					0.10 0.20 0.40 0.70 2.0 3.00 5.00 8. Hazard Ratio (HR)	□□ 00