12 month risk of opioid overdose in patients with T2DM and OUD (Comparision between matched semaglutide vs other anti-diabetes medications groups)

Size/group	Exposure group	Comparison group	cases (overall risk) (Exposure)	cases (overall risk (Comparison))	HR (95% CI)
2,568	semaglutide	insulins	40 (1.6%)	78 (3.0%)	├─■ ─	0.50 (0.34–0.73)
2,427	semaglutide	metformin	38 (1.6%)	82 (3.4%)	├	0.46 (0.32–0.68)
1,652	semaglutide	DPP-4i	28 (1.7%)	52 (3.1%)	├─■	0.53 (0.34–0.84)
2,167	semaglutide	SGLT2i	32 (1.5%)	51 (2.4%)	├	0.61 (0.40–0.96)
1,984	semaglutide	SU	29 (1.5%)	56 (2.8%)	⊢ ■	0.52 (0.33–0.81)
724	semaglutide	TZD	13 (1.8%)	16 (2.2%)	-	0.80 (0.38–1.67)
2,128	semaglutide	other GLP-1RAs	31 (1.5%)	54 (2.5%)	├──	0.58 (0.37–0.90)
					0.20 0.30 0.50 0.80 Hazard R	2.0 3.00 5.00 atio (HR)