

12-month negative control outcomes in patients with obesity and TUD (no T2D) (comparison between propensity-score matched cohorts: semaglutide vs smoking cessation medications)

Outcome	Semaglutide cohort # of cases (overall risk)	Smoking cessation medications cohort # of cases (overall risk)	HR (95% CI)
Semaglutide vs Nicotine replacement therapy (n=1,131/cohort)			
Positive test for SARS-CoV2	51 (4.5%)	45 (4.0%)	1.13 (0.76–1.69)
Nail disorders	14 (1.2%)	16 (1.4%)	0.87 (0.42–1.77)
Injuries to the ankle and foot	41 (3.6%)	38 (3.4%)	1.07 (0.69–1.66)
Injuries to the shoulder and upper arm	27 (2.4%)	28 (2.5%)	0.96 (0.56–1.63)
Semaglutide vs Bupropion/Varenicline (n=1,173/cohort)			
Positive test for SARS-CoV2	51 (4.3%)	54 (4.6%)	0.95 (0.65–1.40)
Nail disorders	17 (1.4%)	<10 (<0.9%)	1.71 (0.78–3.74)
Injuries to the ankle and foot	41 (3.5%)	45 (3.8%)	0.91 (0.60–1.39)
Injuries to the shoulder and upper arm	28 (2.4%)	34 (2.9%)	0.83 (0.50–1.37)

