

Lecture 4: *Natural selection & adaptation*

How do adaptations arise?

Required reading: Coyne - Chapter 5

1

Fitness:

- Relative genetic contribution of individuals to next generation as a result of differences in viability and fertility (= Darwinian fitness)

Selective agent:


- Any factor that causes individuals with certain phenotypes to have higher fitness than individuals with other phenotypes

2

Adaptation – has two meanings; state or process

- Any trait that contributes to fitness by making an organism better able to survive or reproduce in a given environment [as a noun]
- The evolutionary process that leads to the origin and maintenance of such traits [as a verb]

3



4




Figure 6.3
The Economy of Nature, Sixth Edition
© 2010 W.H. Freeman and Company

Ricklefs p.117

5

What would distribution look like for a discrete phenotypic trait?




Figure 6.3
The Economy of Nature, Sixth Edition
© 2010 W.H. Freeman and Company

6

Types of natural selection on phenotypic traits




Figure 6.6
The Economy of Nature, Sixth Edition
© 2010 W.H. Freeman and Company

Ricklefs p.120

Beak adapted for soft seeds



Beak adapted for hard seeds



Disruptive selection on beak size in African finches

- disruptive selection leads to character divergence and in some case may lead to speciation
- requires spatial heterogeneity or discrete resources





Figure 6.7
The Economy of Nature, Sixth Edition
© 2010 W.H. Freeman and Company


Ricklefs p.121

Beak size in Galápagos finch





9

Beak size in Galápagos finch



10

Beak size in Galapagos finch



Copyright © 2004 Pearson Prentice Hall, Inc.

11

Gall size by *Eurosta solidaginis*




EEB Prof. Art Weis

- fly *Eurosta solidaginis* lays egg inside stem of goldenrod
- larvae hatches and creates protective gall
- vulnerable to predation and parasitism

12

Gall size by *Eurosta solidaginis*



Weis & Abrahamson (1986)

13

What kind of selection do we expect to see on gall size of *Eurosta solidaginis*?

Weis & Abrahamson (1986)


14

The struggle to determine the mechanisms of selection

- Today hundreds of measurements of selection demonstrating fitness differences & evolutionary change in traits
- Fewer convincing cases that demonstrate the mechanisms (agents) of selection in natural populations – ecology is difficult!

15

Evolution of cyanide resistance - scale insects



- Alleles conferring cyanide resistance are found at low frequency in populations

- resistance spreads through the population when selection is imposed by cyanide fumigation

- this process serves as model for the evolution of many forms of resistance e.g. antibiotics, insecticides, herbicides


Figure 6.4
The Economy of Nature, Sixth Edition
© 2010 W.H. Freeman and Company

Ricklefs p.118

16

Evolution by pollution

- Evolution of industrial melanism in the peppered moth




17


The peppered moth (*Biston betularia*) and industrial melanism

- Light and dark forms of species that rest on trees
- In UK, before 1850 dark moths rare; caused by single dominant allele
- Industrial pollution blackened tree trunks near cities resulting in increase in black form
- Black variant replaces white in polluted areas; white form predominates in rural unpolluted areas
- Mechanism of selection thought to be predation by birds and differences in crypsis (camouflage) depending on the background of tree trunks

18




19



20

Kettlewell's experiments





Kettlewell (1956)

21

Doubts raised about Kettlewell's bird predation experiments

- Moths raised in the lab and this may have influenced behaviour
- Moths put on trunks at unnaturally high densities
- Moths rarely seen on trunks
- Releases conducted during the day
- Possibility of bat predation not investigated

22



Michael Majerus - Cambridge




Jerry Coyne - Chicago

*"Majerus concludes, reasonably, that all we can deduce from this story is that it is a case of rapid evolution, probably involving pollution & bird predation. I would, however, replace "probably" with "perhaps". *B. bistularia* shows the footprint of natural selection, but we have not yet seen the feet."*

Nature (1998)

23

Revising Kettlewell's experiments



Cook et al. 2012

24

Decline in frequency of the black (melanic) form after introduction of the 'Clean Air Act' in the UK in 1956

The lag in evolutionary response to changes in air pollution levels reflects the time required for forests to return to a more natural (unpolluted) state, as well as a low initial frequency of the recessive allele for typical coloration.




Figure 6.9
The Economy of Nature, Sixth Edition
© 2010 W.H. Freeman and Company

Ricklefs p.123

25

Mouse coat colour polymorphism in contrasting habitats



A

A^+/A^+ or A^+/a^-





B

a^-/a^-




26




Evolution in the lab – 'experimental evolution'

Lenski – founder of
'experimental evolution' –
30 yr. experiment with *E. coli*




> 50,000 generations
of evolution!




29

Experimental evolution with *E. coli*




12 populations propagated under
minimal glucose/citrate medium



42,000 generations in
serial culture

30

Experimental evolution with E. coli




**Samples frozen every 500 generations
allowing relative fitness of different
generations to be compared – “Fossil
record”**

- all populations rapidly increased in fitness
- similar adaptations across strains (e.g. larger cell sizes, higher maximal growth rates on glucose)
- parallel mutations at same genes
- some unique adaptations and distinct genetic changes

31

Novel adaptation in single strain after 33,000 generations!



32