

The Bitcoiner's Dilemma

Bill DeRose

June 28, 2014

I'd like to present a simplified analysis of the bidding that took place for the USMS' bitcoin this past Friday. We model the scenario as a two player zero-sum game between me (I would really like to own those coins) and a single other bidder. Both players may choose from the following set of moves

$$S = \{\text{ABOVE MARKET, AT MARKET, BELOW MARKET}\}.$$

The matrix game is then

$$G = \begin{matrix} & \begin{matrix} AM & @M & BM \end{matrix} \\ \begin{matrix} AM \\ @M \\ BM \end{matrix} & \begin{pmatrix} 0.5 & 1 & 1 \\ 0 & 0.5 & 1 \\ 0 & 0 & 0.5 \end{pmatrix} \end{matrix}$$

where the payoffs are to me, "player one". In this case my moves correspond to selecting rows so my maximin strategy is $x = [1 \ 0 \ 0]^t$. There are a few ways to arrive at this answer and we will begin with the simplest before diving a little further into the math behind it.

If we don't want to dirty our hands with the computation once we've set up our game we can use a solver readily available from [UCLA](#). Just plug in the matrix and let it spit out the answer.

However, if we are so inclined, we can find the optimal strategy ourselves by solving a linear program. Let $x' = [x_1 \ x_2 \ x_3]^t$ be the probability vector that describes my bidding strategy. That is, x_1 is the probability that I bid *above* the market price, x_2 is the probability that I bid *at* the market price, and x_3 is the probability that I bid *below* the market price. To find our optimal strategy, we must solve the following LP

$$\begin{aligned} \text{Maximize} \quad & \min\{0.5x_1, x_1 + 0.5x_2, x_1 + x_2 + 0.5x_3\} \\ \text{Subject to} \quad & x_1 + x_2 + x_3 = 1 \\ & x_1, x_2, x_3 \geq 0 \end{aligned}$$

We wish to maximize our minimum payoff, which depends on the move our opponent (“player two”) selects. The functions whose minimum we wish to maximize correspond to player two’s decision to bid above, at, or below market price, respectively. However, the min function is not linear so we must transform the LP to an equivalent one that a linear solver can handle

$$\begin{array}{ll}
 \text{Maximize} & x_4 \\
 \text{Subject to} & 0.5x_1 \geq x_4 \\
 & x_1 + 0.5x_2 \geq x_4 \\
 & x_1 + x_2 + 0.5x_3 \geq x_4 \\
 & x_1 + x_2 + x_3 = 1 \\
 & x_1, x_2, x_3 \geq 0
 \end{array}$$

Since there are no constraints on x_4 , maximizing it “pushes” its value up until x_4 is exactly equal to $\min\{0.5x_1, x_1 + 0.5x_2, x_1 + x_2 + 0.5x_3\}$. Hence we have achieved our goal of maximizing our minimum payoff. The following R code solves the LP for us

```

> library(lpSolve)
> obj <- c(0, 0, 0, 1)
> con <- matrix(c(0.5, 0, 0, -1,
+                1, 0.5, 0, -1,
+                1, 1, 0.5, -1,
+                1, 1, 1, 0), byrow = T, nrow = 4)
> rhs <- c(0, 0, 0, 1)
> dir <- c(">=", ">=", ">=", "=")
> lp("max", obj, con, dir, rhs)

```

Success: the objective function is 0.5

```

> lp("max", obj, con, dir, rhs)$solution

[1] 1.0 0.0 0.0 0.5

```

Thus the optimal values are $x_1 = 1, x_2 = 0, x_3 = 0, x_4 = 0.5$ and our optimal strategy is to always (with probability 1) bid above the market price.

An even easier way to arrive at this solution is by using [dominance](#) to eliminate possible moves. We can throw out moves two and three because I am guaranteed to always do better (i.e. my payoff is always greater) if I play move one than if I were to play move two or three. Hence, the game I see may be simplified to

$$G' = \begin{array}{c} \begin{array}{ccc} AM & @M & BM \end{array} \\ AM \left(\begin{array}{ccc} 0.5 & 1 & 1 \end{array} \right) \end{array}$$

and again my maximin strategy is $x = [1 \ 0 \ 0]^t$.

The reader is encouraged to solve the LP a third way: graphically. Start by plotting $0.5x_1$, $x_1 + 0.5x_2$, and $x_1 + x_2 + 0.5x_3$ against one another and go from there.