

Serial intervals in SARS-CoV-2 B.1.617.2 variant cases

The SARS-CoV-2 lineage B.1.617.2, also known as the **delta variant**, was declared a variant of concern by WHO on the basis of preliminary evidence suggesting faster spread relative to other circulating variants.¹ However, the epidemiological factors contributing to this difference remain unclear. In particular, an increase in observed growth rate of COVID-19 cases could be the result of a shorter generation interval (ie, the delay from one infection to the next) or an increase in the effective reproduction number, R , of an infected individual (ie, the average number of secondary cases generated by an infectious individual), or both.² Whereas a shorter generation interval would increase the speed but not the number of individual-level transmissions, a larger value of R would require both faster and wider coverage of outbreak control measures such as vaccination or physical distancing to suppress transmission.

In Singapore, whole-genome sequencing is done for respiratory samples from individuals who tested positive for SARS-CoV-2 by PCR with a cycle threshold of 30 and below. The B.1.617.2 variant was first identified in local cases on April 27, 2021. Despite high levels of adherence to mask wearing and physical distancing in the country,^{3,4} clusters of B.1.617.2 variant were detected, and some clusters displayed rapid growth of infections.

We investigated possible drivers of B.1.617.2 variant growth by studying the serial intervals (ie, onset-to-onset delay, a proxy for the generation interval) between pairs of a primary case and a secondary case occurring among household members. Exposure histories were reviewed for all household transmission pairs involving individuals infected with the B.1.617.2 variant and notified between April 27

and May 22, 2021. The B.1.617.2 variant was detected in 97% of the sequenced samples from local cases of COVID-19 identified in this period. Secondary cases with potential exposure to either more than one primary case in the household or to other cases outside the household were omitted from analysis. Households with secondary cases having different symptom onset dates were also omitted from the analysis as we were unable to rule out multiple generations of transmission.

For comparison, we identified household transmission pairs before the partial lockdown in Singapore on April 7, 2020, and applied the same exclusion criteria. This time period precedes the occurrence of the major global SARS-CoV-2 variants and most closely matches the social activity and workplace arrangements in April, 2021,⁵ when working from home was not the default. Preliminary analysis showed that the primary cases in this period had a wider range of time from symptom onset to isolation as compared to the B.1.617.2 primary cases (appendix). As such, the following sampling procedure was done to ensure that we matched the number of transmission pairs and the distribution of time from symptom onset to isolation of primary cases. For a given time from symptom onset to isolation of a B.1.617.2 primary case, we randomly sampled, with replacement, the serial intervals of primary cases in the earlier period with matching time from onset to isolation. We then fitted a skewed normal distribution to the sample of serial intervals to account for negative serial intervals arising from pre-symptomatic transmission. The process was repeated 1000 times to obtain the mean and 95% CI of the sample mean, the median, mode, and the difference of these statistics between the B.1.617.2 variant cases and those cases detected before the lockdown.

There were 32 B.1.617.2 variant household transmission pairs, and 63 household transmission pairs identified before April 7, 2020. The median serial interval of the B.1.617.2

variant cases was 3 days, whereas in cases identified before April 7, 2020, the median serial interval was 3 days (95% CI 2 to 4) after matching the time from symptom onset to isolation (figure). The mode of the serial interval was 2 days for B.1.617.2 variant cases and 2-7 days (95% CI -1 to 4) for cases detected before the lockdown. The mean, median, and mode of the serial interval distributions of B.1.617.2 variant cases and the sampled cases before the lockdown was not statistically different (appendix).

This early investigation of recent B.1.617.2 variant cases offers no evidence to support a large difference



Published Online
August 10, 2021
[https://doi.org/10.1016/S0140-6736\(21\)01697-4](https://doi.org/10.1016/S0140-6736(21)01697-4)

For more on SARS-CoV-2 variants see <https://www.who.int/en/activities/tracking-SARS-CoV-2-variants/>
See Online for appendix




Figure: Probability mass function of serial interval of SARS-CoV-2 variant B.1.617.2 cases (A), probability density function of serial interval of cases identified before the partial lockdown on April 7, 2020 (B), and empirical cumulative density function of serial intervals and estimated cumulative density function of serial intervals (C)

Most primary cases had known exposure (or exposures) outside the household and secondary cases do not have the same exposure as the primary case thereby allowing the directionality of infection to be identified. Negative serial intervals, which signify pre-symptomatic transmission, were also included in the analysis.

(ie, >1 day) in serial intervals among the samples studied, which had an exclusion criteria applied to ensure consistency. In turn, this lends support to the hypothesis that the recent rapid growth is potentially driven by an increase in the average number of secondary cases generated by a case infected with the B.1.617.2 variant. Studies with proper control of confounding factors are thus crucial to tease out the key epidemiological factors that facilitate the increased transmissibility of the B.1.617.2 variant. These factors include, but are not limited to, the viral load and shedding dynamics in individuals infected with the B.1.617.2 variant of SARS-CoV-2, the exposure settings, and the vaccination status of infected individuals. Without signs of lowered disease severity for the B.1.617.2 variant, contact tracing and testing around COVID-19 cases, along with vaccination and non-pharmaceutical interventions, continue to remain key SARS-CoV-2 outbreak control measures in the short term.

RP declares funding from the Singapore Ministry of Health. AJK declares funding from Wellcome and the National Institute of Health Research (NIHR). The views expressed in this publication are those of the authors and not necessarily those of the NIHR or the UK Department of Health and Social Care. TMM and VJL declare no competing interest. Contributors, role of funder, and members of the Centre for the Mathematical Modelling of Infectious Diseases (CMMID) COVID-19 working group are listed in the appendix. The model code and de-identified data that underlie the results reported in this Correspondence are available online at https://github.com/rachaelpong/serial_interval_covid_b.1.617.2.

***Rachael Pung, Tze Minn Mak,
CMMID COVID-19 working group,
Adam J Kucharski, Vernon J Lee
rachaelpong@hotmail.com**

Ministry of Health, Singapore (RP, VJL); Centre for the Mathematical Modelling of Infectious Diseases (RP, AJK) and Department of Infectious Disease Epidemiology (RP, AJK), London School of Hygiene & Tropical Medicine, London, UK; National Public Health Laboratory, National Centre for Infectious Diseases, Singapore (TMM); Saw Swee Hock School of Public Health, National University of Singapore, Singapore (VJL)

- 1 WHO. Weekly epidemiological update on COVID-19 - 11 May 2021. May 11, 2021. <https://www.who.int/publications/m/item/weekly-epidemiological-update-on-covid-19--11-may-2021> (accessed May 23, 2021).

- 2 Park SW, Bolker BM, Funk S, et al. Roles of generation-interval distributions in shaping relative epidemic strength, speed, and control of new SARS-CoV-2 variants. *medRxiv* 2021; published online May 5. <https://doi.org/10.1101/2021.05.03.21256545> (preprint).
- 3 Lim JM, Tun ZM, Kumar V, et al. Population anxiety and positive behaviour change during the COVID-19 epidemic: cross-sectional surveys in Singapore, China and Italy. *Influenza Other Respir Viruses* 2021; **15**: 45–55.
- 4 Sim S. More than 6 in 10 Singaporeans likely to continue good hygiene practice after Covid-19: NTU study. Jan 26, 2021. <https://www.straitstimes.com/singapore/more-than-6-in-10-singaporeans-likely-to-continue-good-hygiene-practice-after-covid-19-ntu> (accessed May 23, 2021).
- 5 Ministry of Health, Singapore. Expansion of vaccination programme; further easing of community measures. March 24, 2021. <https://www.moh.gov.sg/news-highlights/details/expansion-of-vaccination-programme-further-easing-of-community-measures> (accessed May 23, 2021).